5 Antong

SYSTEM

OF

## ANATOMY.

FROM

MONRO, WINSLOW, INNES,
And the LATEST AUTHORS.

ARRANGED,

AS NEARLY AS THE NATURE OF THE WORK WOULD ADMIT,

INTHE

Order of the Lectures delivered by the Professor of ANATOMY in the University of Edinburgh.

IN TWO VOLUMES.

ILLUSTRATED WITH COPPERPLATES.

VOL. I.



EDINBURGH:
Printed for C. ELLIOT, PARLIAMENT-SQUARE.

M. DCC. LXXXIV.

GO 403.

C2.5 S83



### ADVERTISEMENT by the EDITOR.

A 6 for several years past the Students of Medicine in this place have complained of the great expence of Anatomical Books, and of the number they were obliged to search into before they could acquire the knowledge they wanted, the editor has been advised by some of his medical friends to publish a system which should contain the principal parts of some of the latest and best authors on anatomy.

Accordingly, the following work comprehends—Monro on the Bones; Winflow on the Joints and other parts of the fresh bones; Innes on the Muscles, with his Illustration of these and of the Skeleton; Winslow on the Viscera, Blood-vessels, and Organs of the Senses; Monro, Hewson, and other late writers, on the Nerves and Lymphatic vessels: The whole arranged in the order which appeared to be the most natural, and the easiest for the student; and forming a system more complete than any that has hitherto appeared.

At first it was proposed to have published the above without any alteration: but, upon examining them more attentively, it was found necessary to throw out much superfluous matter in some parts, and to make many corrections and additions in others, so as to render the work adequate to the purposes for which it was intended.

The description of the Bones and Muscles has undergone very little alteration, as few or no discoveries relating

A z

to these parts have been made since the last editions of Monro and Innes were published. But of the Viscera, Blood-vessels, Organs of Sense, and other parts taken from Winslow, the editor has been at considerable pains and expence to render the descriptions more complete. For although, on these parts, Winslow is one of the best authors we are acquainted with, there yet appeared much room for alteration. He is often two prolix, some of his descriptions are erroneous, several of them are impersect, a few things are altogether omitted, and many of his terms, &c. are antiquated or improper. In all these respects improvements have been made, and the discoveries of laterwriters have been occasionally introduced.


Farther: As descriptions alone without figures are insufficient to give the beginner a just idea of the different
parts of which the body is composed, a number of engravings have been added, to render the work more useful and
agreeable. These were taken from authors whose works are
well known to be accurate, as Eustachius, Albinus, Cheselden, Haller, Hewson, &c. including the Eight Tables
formerly selected and published by Mr Innes.

These particulars, it is presumed, will be considered as adding greatly to the importance and utility of the Work. But, independent of these altogether, and supposing a republication of the above pieces, in the present form, had simply taken place, without correction, and without plates; even then it would have constituted an acceptable performance on the sooting of convenience and economy. Uniform, compact, and comprehensive, it is calculated to superfede a multiplicity of volumes, and to save much unnecessary expense. Of six or seven books on this subject, with

with which students were wont to be encumbered, all the parts from which they could derive any advantage are here compressed into the compass of two commodious volumes, and afforded at less than a third part of the expence.

Upon the whole, if this work shall be considered by students as an useful acquisition, the publisher cannot help informing them that they owe it chiefly to the approbation of the illustrious Professor of the Science of which it treats: For although, as already mentioned, the design was originally suggested by some gentlemen for whose judgment the publisher has the greatest deference, yet it was the above approbation alone which finally determined him to carry it into execution.

EDIN. Nov. 1783.



# CONTENTS

#### OF THE FIRST VOLUME.

### PART I.

## Of the Human Bones.

			Page.
CHAP. I. Of the Bones in ge	meral,		17
CHAP. II. Of the Skeleton,			62
Sect. I. Of the Head,		•	63
§ 1. The Cranium,	•	•	64
§ 2. Bones of the Face	•		109
Sect. II. Of the Trunk,		•	144
§ 1. The Spine,	•		ib.
§ 2. The Pelvis,	•	•	171
§ 3. The Thorax,			182
Sect. III. Of the Superior	Extremities		196
§ 1. The Shoulder,			ib.
§ 2. The Arm,	•	•	205
§ 3. The Fore-arm,		•	209
§ 4. The hand,			215
Sect. IV. Of the Inferior	Extremities,		229
§ 1. The Thigh,			ib.
§ 2. The Leg,	_		234
§ 3. The Foot,	•		244
APPENDIX. Of the Female	Skeleton.		257
	4		

### PART II.

## Description of the Human Muscles.

CHAP. I. Muscles of the Teguments of the Cranium,	267
CHAP. II. Of the Muscles of the Ear, -	269
CHAP. III. Of the Muscles of the Eye-lids,	273
	HAP.

CHAP. IV. Muscles of the Eye-ball, -	274
CHAP. V. Muscles of the Nose, -	277
CHAP. VI. Muscles of the Mouth and Lips,	278
CHAP. VII. Muscles of the Lower Jaw, -	282
CHAP. VIII. The Muscles which appear about the	
anterior Part of the Neck, -	284
CHAP. IX. Muscles situated between the Lower Jaw	
and Os Hyoides, -	286
CHAP. X. Muscles situated between the Os Hyoides	
and Trunk,	288
CHAP. XI. Muscles situated between the Lower Jaw	
and Os Hyoides laterally, -	290
CHAP. XII. Muscles situated about the Entry to the	
Fauces,	293
CHAP. XIII. Muscles situated on the posterior Part	-/3
of the Pharynx, -	295
CHAP. XIV. Muscles situated about the Glottis,	296
CHAP. XV. Muscies situated on the anterior Part of	-,-
the Abdomen,	299
CHAP. XVI. Muscles about the Male Organs of Ge-	-77
neration, -	304
CHAP. XVII. Muscles of the Anus, -	306
CHAP. XVIII. Muscles of the Female Organs of Ge-	500
neration, -	308
CHAP. XIX. Muscles situated within the Pelvis,	310
CHAP. XX. Muscles situated within the Cavity of	
the Abdomen, -	
CHAP. XXI. Muscles situated on the anterior Part	311
of the Thorax,	315
CHAP. XXII. Muscles situated between the Ribs and	2,2
within the Thorax,	316
CHAP. XXIII. Muscles situated on the anterior Part	310
of the Neck close to the Vertebræ,	318
CHAP. XXIV. Muscles situated on the posterior Part	310
of the Trunk,	220
CHAP. XXV. Muscles of the Superior Extremitics,	320
	331
CHAP. XXVI. Muscles situated on the Os Humeri,	334
[2] 2 - 15 - 15 - 15 - 15 - 15 - 15 - 15 -	HAP.

CONTENTS.	ix
CHAP. XXVII. Muscles situated on the Cubit or	
Fore-arm, CHAP. XXVIII. Muscles situated on the Hand	337
chiefly,	343
CHAP. XXIX. Muscles of the Inferior Extremities,	350
CHAP. XXX. Muscles situated on the Thigh,	355
CHAP. XXXI. Muscles situated on the Leg,	360
CHAP. XXXII. Muscles which are situated chiefly	500
on the Foot,	362
PART III.	
Explanations of the Skeleton and Muscle	s.
Table I. A front View of the Human Skeleton, with fome of the Ligaments and Cartilages	
which connect the Bones to each other, Table II. The first Layer of Muscles situated on the anterior Part of the whole Body, im- mediately under the common Teguments,	375
and tendinous Fasciæ,  Table III. The Second Layer of Muscles on the an-	379
Table IV. Represents the right Eye-ball, the fix Muscles which move it, taken out of the	382
Orbit, with the Optic Nerve, Table V. A back View of the Human Skeleton, with fome of the Ligaments and Cartilages	385
which connect the Bones, Table VI. A Back View of the Muscles which are immediately situated below the common	
Integuments,	392
Table VII. The Second Layer of the Muscles on the Back Part of the Body,	
Table VIII. The Third Layer of Muscles on the po- sterior Part of the Body, with some of	
the Ligaments and naked Bones,	397
P A	RT

### PART IV.

# An Enumeration of the Cartilages, Ligaments, &c. of the Joints.

Those of the Head, -	405
- of the Trunk, -	409
of the Sternum and Ribs,	413
of the Pelvis, -	415

4

## SYSTEM OF ANATOMY.

#### PART I.

CONTAINING THE

## ANATOMY

OFTHE


# HUMAN BONES.

BY THE LATE

ALEXANDER MONRO, M.D. AND F.R.S.

Fellow of the Royal College of Physicians, and Professor of Medicine and Anatomy in the University

of Edinburgh.



#### STUDENTS OF ANATOMY

INTHE

#### UNIVERSITY OF EDINBURGH.

GENTLEMEN,

WHEN this Ofteology was first printed in 1726, I did not know that Albinus, Winslow, and Palfyn, were to publish descriptions of the bones; otherwise my papers probably would have remained yet undelivered to the printers. I, however, flatter myself, that this essay has been of use to the gentleman who did me the honour to attend my lectures, by assisting them to understand my sense and representation of things in this fundamental part of anatomy; and that it has possibly been of more advantage to them than a more complete work from an abler hand, unless my demonstrations had been in the order and method of such an author.

This view of your improvement, Gentlemen, is a prevailing argument with me to cause this essay to be reprinted; and you cannot reasonably blame me, if I likewise acknowledge another motive for it, which more particularly relates to mysels. In a new edition an author has an opportunity of making his works more correct, complete, and consequently acceptable to the public, who may perhaps be indulgent enough to think this little treatise not altogether useless; since more reasoning on the structure and morbid phanomena of bones is to be found in it, than in the other writers, who have confined themselves almost entirely to the descriptive or proper anatomical part of the osteology.

I have here kept to the plan of the former editions, by first considering, in the order that seemed to me most natural and methodical, every thing which I thought necessary to be known concerning bones in general; and, in the second part, I have described the several bones composing the skeleton.

The bones of adults are what I principally endeavour to describe; but I have added as much of the ofteogenea as I think serviceable in the practice of physic and surgery.

That little might be omitted of what was formerly done on this subject, I have taken all the affistance I could from books; but have never afferted any anatomical fact on their authority, without confulting nature, from which all the descriptions are made: and therefore the quotations from fuch books ferve only to do justice to the authors, who have remarked any thing in the structure of the parts that was commonly omitted, and to initiate you in the history of anatomy; which I once proposed to make complete, so far as related to this subject: But not being able to procure several books, and being fenfible how many more may have never come to my knowledge, I laid aside this design, of purposeomitted many I could have inferted, and in some places I have changed an older author for a later one who has more fully or clearly described what I treated of. Beside anatomists, I have also named several other authors to confirm my reasoning by practical cases; of which it is not to be supposed my own experience could furnish a sufficient variety.

You will readily observe, that I quote no passages with a view to criticise or condemn them. This precaution of giving no offence, is very necessary in those who are sufficiently conscious of their being liable to lay themselves open to just censure; and it prevents occasions of useless wrangling, in which generally both parties are losers, and the public has

little advantage.

In this treatife I always make use of the most common name of each part, and have put the synonimous name to be met with in books at the soot of the page, that the reading might be smoother, and you might consult them at your lei-

fure to affift you in understanding different authors.

The descriptions and reasoning are blended, without which I always find young anatomists are soon disgusted with authors: Their imaginations cannot follow a long chain of descriptions, especially when they are not taught at the same time the uses which the described parts serve: Their minds must have some relaxation, by a mixture of reasoning, which never misses to strike the sancy agreeably, and raises a strong desire to understand the principles on which it depends.

The phenomena of diseases are all deduced in this essay from the structure of the parts, by way of corollaries and questions; which such an anatomical work confined me to. And this method has otherwise a good essect: For, when one meets with an useful proposition, and is obliged to employ a little thought to find out its solution, the impression it

makes is deeper, and he acquires a fondness for it as being in part his own discovery. My pupils have frequently assured me, that they could, with very small reslection, trace out the whole reasoning from which my conclusions were drawn; I hope their successors will also think this an agreeable man-

ner of being instructed.

Those gentlemen who desired I would add the lectures which I pronounce in my colleges as a commentary upon the text, where the diseases are mentioned, will, I persuade myself, excuse me for not complying with their desire, when they consider the design of this is to be a school-book, and how great the difference is between instructing youth in private, and pretending to inform the public. Art. xxiv. vol. v. of Medical Essays and Observations, published in this place, is one of these lectures which I gave as a commentary on the paragraph (p. 26.) concerning the different kinds of caries.


In this edition, I have corrected the mistakes and obscure passages which I discovered in the former, and in some places I have made the descriptions more full and exact, aiming all I could to shun unnecessary minuteness on the one hand, and a blameable inaccuracy on the other: Whether I have hit that just medium, is what you and the public must now

judge.

You have advantageous opportunities in this place of studying all parts of medicine, under the professors of its disterent branches in the University, and of seeing the practice of pharmacy, surgery, and physic, with our surgeon-apothecaries, and in the Royal Instrmary where the diseased poor are carefully treated. These your interest, and, I hope, your inclinations, will lead you, Gentlemen, so to improve, as that they become the happy means of your making a considerable sigure in your several stations. Whatever assistance is in my power towards such a desirable event, shall be given with the greatest pleasure by

Your humble fervant,

ALEX. MONRO.



### SYSTEM

OF

## ANATOMY

# PART I.

#### HUMAN BONES.

#### C H A P. I.

Of the BONES in general.

#### THE PERIOSTEUM.

BONES are covered by a membrane, named on that account Periosteum (a), which is so necessary to them, that we must examine its texture and uses before we can understand their structure. The periosseum, as well as most other membranes, can be divided into layers of fibres. The exterior ones, composed of the fibres of the muscles connected to the Vol. I.

B bones,

<sup>(</sup>a) Membrana circumoffalis, omentum offibus impositum.

bones, vary in their number, fize, and direction, and consequently occasion a very great difference in the thickness and strength of the periosteum of different bones, and even of the different parts of the same bone. The internal layers is every where nearly of a similar structure, and has its sibres in the same direction with those of the bone to which they are contiguous. Ought not then the name periosteum to be applied, strictly speaking, only to this internal layer, to which the others are

joined in an uncertain manner and number?

Some authors (b) endeavour to prove the internal layers of fibres of the periosteum to be derived from the dura mater: For, fay they, fince the membrane covering the skull is plainly a production or continuation of the dura mater, which passes out between the sutures; and fince there are muscles on the head, as well as in other parts, which might furnish a periosteum; it is needless to assign different origins to membranes which have the fame texture and uses. They add further, in proof of this doctrine, that the periosteum extends itfelf along the ligaments of the articulations from one bone to another; and therefore is continued from its origin over all the bones of the body.—While anatomists were fond of the hypothesis of all membranes being derived from one or other of the two that cover the brain, a dispute of this kind might be thought of confequence: but now that the hypothesis is neglected as useless, it is needless to examine the arguments for or against it.

Except where muscles, cartilages, or ligaments, are inserted into the periosteum, its external surface is connected to the surrounding parts by thin cellular membranes, which can easily be stretched considerably, but shorten themselves whenever the stretching force is removed. When these membranes are cut off or broken, they collapse into such a small space, that the surface of

the periofteum feems fmooth and equal.

When

<sup>(</sup>b) Havers, Ofteolog. Nov. difc. 1. p. 16.

When we attempt to tear off the periosteum from bones, we see a great number of white threads produced from the membrane into them; and, after a successful injection of the arteries with a red liquor, numerous vessels are not only seen on the periosteum (c), but most of the sibres sent from the membranes to the bone, show themselves to be vessels entering it, with the injected liquor in them; and when they are broken, by tearing off the periosteum, the surface of the bone is almost covered with red points.

The veins corresponding to these arteries are sometimes to be seen in subjects that die with their vessels full of blood; though such numerous ramifications of them, as of the arteries, can seldom be demonstrated, because sew of them naturally contain coloured liquors, and such liquors can difficultly be injected into them.

This, however, is fometimes done (d).

The great fensibility of the periosteum in the deepfeated species of paronychia, in exostoses, nodi, tophi, and gummata, from a lues venerea, or whenever this membrane is in an inflamed state, is a sufficient proof that it is well provided with nerves, though they are perhaps too small to be traced upon it; and therefore one cannot well determine, whether they are sent along with the arteries in the common way, or are derived from the tendinous sibres of the muscles expanded on the periosteum (e).

Vessels also pass through the periosteum to the marrow; of which more hereaster. And frequently muscles, ligaments, or cartilages, pierce through the periosteum,

to be inferted into the bones.

The chief uses of the periosteum are: 1. To allow B 2

(c) Ruysch. Epist. 5. tab. 5. sig. 1. 2. epist. 8. tab. 9. sig. 1. 9.
(d) Sue traité d'osteologie, traduit de l'Anglois de Mr Monro.

Note in page 9.

t

f

<sup>(</sup>e) See the dispute about the sensibility of this and of other membranes in Zimmerman Differt de irritabilit.—Act. Gotting. vol. 2.— Haller sur la nature sensible et irritable.—Whytt's physiolog. essay 2.—Remar. Differt. de sungo articulor. § 26. 34.

the muscles, when they contract or are stretched, to move and flide easily upon the bones; the smooth furface of this membrane preventing any ill effects of their friction upon each other. 2. To keep in due order and to support the vessels in their passage to the bones. 3. By being firmly braced on the bones, to affift in fetting limits to their increase, and to check their overgrowth. 4. To strengthen the conjunction of the bones with their epiphyses, ligaments, and cartilages, which are eafily separated in young creatures, when this membrane is taken away. 5. To afford convenient origin and infertion to feveral muscles which are fixed to this And laftly, to warn us when any injury membrane. is offered to the parts it covers; which, being infenfible, might otherwise be destroyed without our knowledge or endeavouring to procure a remedy.

When the cellular fubstance connecting the periosteum to the surrounding parts is destroyed, these parts
are fixed to that membrane, and lose the sliding motion
they had upon it; as we see daily in issues, or any other
tedious suppurations near a bone.—When the vessels
which go from the periosteum to the bones are broken
or eroded, a collection of liquor is made between
them, which produces a fordid ulcer or rotten bone.
This often is the case after fractures of bones and inslammations of the periosteum, or after small-pox,
measles, spotted severs, and erysipelas.—Do not the
disorders of the periosteum, coming rather along with
or soon after the cutaneous than other diseases, indicate
some similarity of structure in the periosteum and skin?

#### OF THE COMPOSITION OF BONES.

THE bones are the most hard and solid parts of the body; and, like all other parts where large vessels do not enter, are generally of a white colour; only in a living creature they are bluish, which is owing to the blood in the small vessels under their surface. The less therefore and sewer the vessels are, and the thicker and sirmer the

the bony furface covering the vessels is, the bones are whiter. Hence the bones of adults are whiter than those of children; and, in both young and old, the white colour of different bones, or of the feveral parts of the fame bone, is always in proportion to their veffels and folidities; which circumftances ought to be regarded by furgeons, when they are to judge of the condition of bones laid bare.

Bones are composed of a great many plates (f), each of which is made up of fibres or ftrings united by fmaller fibrils (g); which being irregularly disposed, and interwoven with the other larger fibres, make a reticular work—This texture is plainly feen in the bones of fœtuses, which have not their parts closely compacted; and in the bones of adults which have been burnt, long exposed to the weather, or whose composition has been made loofe by difeafes. - The chinks, which are generally made according to the directions of the larger fibres of bones that have undergone the action of fire or of the weather, show the greater strength of these than of the fibres which connect them. Numerous accurate observations of the different times in which exfoliations are made from the fides or ends of fimilar bones, might bid fair to determine what is the proportional force of cohesion in the two forts of fibres.

The plates are faid (b) to be firmly joined to each other by a great number of claviculi, or small bony processes, which, rising from the inner plates, pierce through some, and are fixed into the more external Of these nails, four kinds, viz. the perpendicular, oblique, beaded, and crooked, have been described: But in bones fitly prepared, I could only fee B 3

(f) Squamæ, bracteæ, laminæ.

<sup>(</sup>g) Malpigh. Anat. plant. & oper. posthum.
(h) Gagliard. Anatom. offium nov. invent. illustrat. cap. 1. ebf. 2.

numerous irregular processes rising out from the

plates (i).

Though the exterior part of bones is composed of firm compact plates, yet they are all more or less cavernous internally, In some (e.g. middle thin part of the scapula and os ilium) the solid sides are brought so near, that little cavity can be seen; and in others (middle of os humeri, femoris, &c.) the cavities are so large, that such bones are generally esteemed to be hollow or sistular. But the internal spongy texture is evident in young animals; and some of it may be seen to remain in those of the greatest age, when bones are cautiously opened, after they have been kept so long as to be free of the oil they contain, or after being burnt.

This fpongy cavernous internal part of bones is generally called their Cancelli or LATTICE-WORK, and is formed in the following manner. The plates are firmly joined about the middle of the bone; but as they are extended towards its ends, the more internal parts separate from the exterior, and stretch out their fibres towards the axis of the bone, where they are interwoven with the fibres of other plates that have been fent off in the fame way. Seeing the plates are thus constantly going off, the folid fides of the bones must become thinner, and the lattice-work must be thicker and stronger towards their ends. This is evident in many of them, where the folid fides of their middle are very thick, and the cancelli are scarce observable; whereas, at the ends, where their diameter is greatest, the folid walls or fides are not thicker than paper, and the cancelli are numerous, and large enough to fill up the whole space left between the fides.

The twisting and windings which these cancelli make, and the interstices which they leave, differ considerably in figure, number, and fize; and therefore form little cells, which are as different, but communicate with

<sup>(</sup>i) Malpigh. oper. posthum.

Chap. I.

with each other. Some writers (k) minutely remark these different appearances of the cancelli, after they begin to separate from the plates; and from thence distinguish them into wrinkled, perforated, and net-like.

The cancelli sustain the membranous bags of the marrow which are stretched upon them, and thereby hinder these membranous parts from being torn or removed out of their proper places in the violent motions and different postures which the bones are employed in. This support which the cancelli afford the marrow, also saves its membranes and vessels, in the lower parts of the bones, from being compressed by the weight of the marrow above.

The depressions between the sibres of the external plates of bones appear like so many surrows on their surface, into each of which the periosteum enters; by which the surface of contact, consequently the cohesion, between it and the bone, is considerably increated, and a greater number of vessels is sent from it into the bone than if it was a plain surface.

Both on the ridges and furrows, numerous little pits or orifices of canals are to be seen, by which the vessels

pass to and from the bones.

After a fuccessful injection, the arteries can be traced in their course from the pits to the plates and sibres; and, in sawing, cutting, or rasping the bones of living creatures, these vessels discover themselves by the small drops of blood which then ooze out from the most solid part of the bones. But the clearest demonstration of the intimate distribution of these small arteries, is, to observe the effect of such a tinging substance as can retain its colour, when swallowed, digested, and mixed with the blood of any living animal, and at the same time has particles small enough to be conveyed into the vessels of the bones; such is rubia tinctorum, madder-root (1): For we see the gra-

(k) Gagliard. Anat. offium, cap. 1. obf. 4. 5. 6. 7.

<sup>(1)</sup> Philosoph. Transact. num. 442. art. 8. num. 443. art 2. num. 457. art. 4. — Mem. de l'acad. des sciences. 1739, 1742.

n

dual advances which this tincture makes from the periosteum into the more internal parts of the bones, and how univerfally the distribution of the liquors is made, the whole bony fubstance being tinged. Whether the time in which this tinged liquor passes from the outer to the internal plates, till all the plates are made of its colour, and the time which the disappearing of the dye, after giving the creature no more of this fort of food, makes us think it takes to return, are the fame in which the natural liquors circulate, is uncertain; because this tinging substance may move more flowly, or may pass more quickly, than the natural liquors do.—The arteries are larger near each end than at the middle of the large bones that are much moved; because they not only serve the bony plates near the ends, but pass through them to the marrow. -As animals advance in age, the arteries of the bones become less capacious; as is evident, 1. From the bones of adults having less blood in them than those of children have. 2. From many of them becoming incapable in old age of admitting the coloured powders used in injections, which easily pass in youth. And, 2. From the bones of old creatures being more difficultly tinged with madder than those of young ones. If authors have not mistaken, the arteries of bones have fometimes become very large (m).

We may conclude, from arteries being accompanied with veins, so far as we can trace them in every other part of the body, that there are also veins in the bones; and the disappearing of the tincture of madder, after bones of living animals are coloured with it, could not be without such veins to carry it away; nay, the veins of bones can sometimes be injected, and then seen (n).

The bones of a living animal are so insensible, that they can be cut, rasped, or burnt, without putting the crea-

(n) Sue trad. d'ofteolog. p. 9.

<sup>(</sup>m) Diemerbroek Anat. lib. 9. cap. 1.—Mery, Hist. de l'acad. des sciences, 1704.

creature to pain, and the nerves distributed in their fubstance cannot be shown by diffection; from which it might be inferred that they have no nerves distributed to them: but the general tenor of nature, which beflows nerves to all other parts, should prevent our drawing such a conclusion. And if fensibility is a fure proof of nerves entering into the composition of any part, as it is generally allowed to be, we have fufficient evidence of nerves here in the bones; for the granulated red flesh which sprouts out from them, after an amputation of a limb, or performing the operation: of the trepan, or after an exfoliation, is exquifitely fenfible: and in some ulcers of bones, where the periosteum was all separated, the patient suffered racking pain, if the bone was touched with a rough instrument; nor was he free of pain after the bone was perforated (o).——The reason why the nerves of rigid hard bones become insensible, is, that all nerves must have a confiderable degree of flexibility at the part where objects are applied, otherwise it cannot be affected by their impressions. We see this illustrated in a very common analogous case, the growth of a new nail: when the former one has suppurated off, the thin membrane, which first appears, is exquisitely sensible; but gradually becomes dull in its fensation, till it can be cut or scraped, without causing pain, after it is formed into a hard nail.

From what has been faid of the vessels of bones, it is evident, that there is a constant circulation of sluids in every part of them; and that there is a perpetual waste and renewal of the particles which compose the solid sibres of bones, as well as of other parts of the body; the addition from the sluids exceeding the waste during the growth of the bones; the renewal and waste keeping pretty near par in adult middle age; and the waste exceeding the supply from the liquors in old age; as is demonstrable from their weight: for each bone increases in weight

C

as a person approaches to maturity; continues of nearly the same weight till old age begins, and then becomes lighter.—The specific gravity of the solid sides, on the contrary, increases by age; for then they become more hard, compact, and dense. In consequence of this, the bones of old people are thinner and sirmer in their sides, and have larger cavities, than those of young persons.

The vascular texture of bones must make them subject to obstructions, ecchymoses, ulcers, gangrenes, and most other diseases which the softer parts are affected with; and therefore there may be a greater variety of caries than is commonly described (p).

Hence we can account for the following appearances. Hamorrhagies from fungous flesh rifing out from the most folid part of a cut bone (q).

The regular alternate elevation and subsiding, or apparent pulsation, frequently to be seen in some of the cells of a carious bone.

Cells refembling cancelli, fometimes feen in the part of a bone, which, in a natural state, is the most folid and firm (r).

A bone, as a tube, including another bone within it (s).

On the internal furface of the folid parts of bones there are orifices of canals, which pass outwards through the plates to open into other canals that are in a longitudinal direction; from which other tranverse passages go out to terminate in other longitudinal canals: and this structure is continued through the whole substance of bones; both these kinds of canals becoming smaller gradually as they approach the outer surface (t).—

These canals are to be seen to the best advantage in a bone

<sup>(</sup>p) Edin. Medical effays and obs. vol. 5. art. 25.

<sup>(9)</sup> Medical effays, vol. 4. art. 21.

<sup>(</sup>r) Ruysch. Thef. 8. num. 8. Thef. 10. num. 176.

<sup>(</sup>s) Idem, ibid.

<sup>(</sup>t) Havers Ofteolog. Nov. p. 43.

L

r-

e-

es,

ne

of in

ıg

b-

nd ed

of

s.

le

p-

10

rt

n

25

h

1-

d

e

r

a

bone burnt till it is white. When it is broken transversely, the orifices of the longitudinal canals are in view; and when we separate the plates, the transverse ones are to be observed. Here, however, we are in danger of of believing both these forts of canals more numerous than they really are; because the holes made by the processes connecting the plates of bones have the appearance of the transverse (u), and the passages for the blood-vessels resemble the longitudinal canals. I do not know how we are to keep free of error about the transverse canals; but think we may distinguish between the two kinds of longitudinal ones: for the passages of the vessels are largest near the external surface of the bone, and every transverse section of them is circular; whereas the longitudinal canals are largest near the cancelli, and their transverse sections appear to me of a flat oval figure, which may be owing to the different momentum of the fluids conveyed in them.—The fituation of the larger longitudinal canals, and of the passages of the larger veffels, makes a bone appear more denie and compact in the middle of its folid fides, than towards its outer and inner furfaces, where it is spongy.

We see marrow contained in the larger tranverse and longitudinal canals just now described, and from thence judge that it passes also into the smaller ones. The drops of oil which we discover with a microscope every where on the surface of a recent bone fractured transversely, and the oozing of oil through the most solid bones of a skeleton, which renders them greafy and yellow, are a confirmation of the use of these canals. Of what advantage this distribution of the marrow through the substance of bones is, will be mentioned when the nature and use of this animal-oil is inquired

into.

Most bones have one or more large oblique canals formed through their sides for the passage of the medullary vessels, which are to be described afterwards.

Bones

<sup>(</sup>u) Morgagn. Advers. 2. animad. 25.

al

b

Bones exposed to a strong fire in chemical vessels, are refolved, in the same manner as the other parts of of animals, into phlegm, spirit, volatile salt, fetid oil, and a black caput mortuum. But the proportion of these principles varies according to the age, folidities, and other circumstances of bones. Young bones yield the largest proportion of phlegm; spongy bones afford most oil; and folid ones give most falt and black residuum.—Though this refiduum can scarce be changed by the force of fire while it is in close veffels; yet, when it is burnt in an open fire, the tenacious oil, to which it owes its black colour, is forced away, and a white earth is left that has little or no fixed falt in it. This earth feems to be the proper conflituent folid part of bones, and the other principles give it firmness and tenacity: for the quantity of the earth is fo great, that, after all the other principles are separated from a bone, its former shape and fize remain (v); but it is very brittle till it is moistened with water or oil, when it recovers fome tenacity.—The increase of the proportion of earth in old peoples bones, is one reason of their being more brittle than those of young people are.

Lest any imagine the falts and oils of bones, while in a natural state, to be of the same acrid kind with those obtained from them by the chemical analysis, it is to be observed, that these principles may be extracted from bones in the form of a very mild jelly, by boiling them in water.

The bones fustain and defend the other parts of the

body.

Bones are lined within, as well as covered externally, with a membrane; which is therefore commonly called Periosteum Internum.

#### INTERNAL PERIOSTEUM.

THE internal periosteum is an extremly fine membrane;

(v) Havers Ofteolog. Nov. difc. 1. p. 32.

Is,

of

nd

efe

nd

he

rd

si-

n-

t,

to

a

t.

rt

d

t,

,

y

-

-

f

e

e

1

brane; nay, frequently, it has a loofe reticular texture; and therefore it is compared by some to the arachnoid coat of the spinal marrow: so that we cannot expect to divide it into layers as we can divide the external peri-We can, however, observe its processes entering into the transverse pores of the bones, where probably they are continued to form the immediate canals for the marrow distributed through the substance of the bones; and along with them veffels are fent, as from the external periofteum, into the bone (x). These processes being of a very delicate texture, the adhesion of this membrane to the bone is fo small, that it separates commonly more easily from the bone than from the marrow which it contains: wherefore one might call it the common membrane of the marrow, rather than by the name it now has. But whether the one or the other defignation ought to be given it, is not worthy a dispute.

From the internal furface of the internal periosteum, a a great number of thin membranes are produced; which, passing across the cavity, unite with others of the same kind, and form so many distinct bags, which communicate with each other; and these again are fubdivided into communicating vesicular cells, in which the marrow is contained. Hence it is, that the marrow, when hardened, and viewed with a microscope, appears like a cluster of small pearl; and that the hardened marrow of bones, buried long under ground, or laid fome time in water and then dried, is granulous (y).—This texture is much the fame with what obtains in the other cellular parts of the body, where fat is collected; only that the cells containing the marrow are smaller than those of the tunica adiposa or cellulofa elsewhere; which probably is owing to their being inclosed in the bones, where they are not fo much

stretched or extended as in other parts.

OF

(y) Ruysch. Thesaur. 9. num. 2. et Advers. dec. 111. obs. 9.

<sup>(</sup>x) Winflow Exposition anat. des os frais, sect. 82. 83.

#### OF THE MARROW.

THE Marrow is the oily part of the blood, separated by small arteries, and depositated in these cells. Its colour and consistence may therefore vary according to the state of the vessels, and their distribution on the membranes of the cells.

The marrow, as well as the other fat of the body, chemically analized, yields, befides oil and water, a confiderable proportion of an acid liquor, but no alkali (z). This may be the reason of its being less putrescent than the blood or most other parts of animals (a); which is a necessary quality in a substance that is constantly exposed to a considerable degree of heat, and is more in

a stagnating condition than the other liquors.

Besides the arteries, which I mentioned already (p. 23.) to be sent from the bones to the marrow, there is at least one artery for each bone; several bones have more, whose principal use is to convey and secern this oily matter. After these arteries have pierced the solid side of a bone, they are divided into several branches; which soon are distributed every where on the internal periosteum, and afterwards spread their branches inwards on the medullary cells, and outwards through the tables of the bone.

The blood which remains after the fecretion of the marrow is returned by proper veins, which are collected from the membranes into one or two large trunks, to pass out at the same holes or passages at

which the artery or arteries enter.

The general rule of the small vessels decreasing in their capacities as animals advance in age, to which many phenomena in the animal-economy are owing, obtains here: for though the trunks of the medullary vessels enlarge as animals turn older, yet the small branches become smaller; as is evident from injections,

(z) Grutzmaker Dissert. de ossium medulla.—Haller Element. physiolog. lib. 4. sect. 4.

(a) Pringle Append. to camp-diseases, exper. 47.

t T.

ra-

ls.

ng

he

le-

fi-

2).

an

18

X-

in

dy

re

ve

is

id

S;

r-

es

gh

ne

1-

ce

at

n

h

5,

y

h

t.

which cannot be made to pass near so far in these vesfels of adults as of children. Hence the marrow is bloody in children, oily and balmy in middle age, and thin and watery in old people.

By experiments made on the marrow, when bones of living animals are opened or cut through (b), and from the racking pain with which suppurations within bones are frequently attended, we have sufficient proof that the membranes here are sensible, and consequently have nerves distributed to them. Hippocrates (c) might therefore say justly, that a wound penetrating into the cavity of a bone may produce a defirium.

The vessels of the marrow, wrapt up in one common coat from the periosteum, pass through the bones by proper canals; the most considerable of which are about the middle of each bone, and are very oblique. Sometimes these vessels continue at a little distance in their passage, when the canal is divided by a small bony partition or two.

From the structure of the contents of the bones, we may judge how these parts, as well as others, may be subject to oedema, phlegmon, erysipelas, scirrhus, &c. and may thence be led to a cure of each, before the common consequence, putrefaction, takes place, and frequently occasions the loss of the limb, if not of the patient.

The marrow is of very confiderable use to the bones: for by entering their transverse canals, and passing from them into the longitudinal ones, it is communicated to all the plates to soften and connect their fibres, whereby they are preserved from becoming too brittle; as we see they do in burnt bones, or those long exposed to the air, in people labouring under old age, pox, or scurvy. In all which cases, the oil is either in too little quantity,

<sup>(</sup>b) Du Verney, Memoires de l'acad. des sciences, 1700.

<sup>(</sup>c) Aphorism. § 7. aph. 24.

di

ar

X

n

ta

ti

a

n

f

t

quantity, or has its natural good qualities changed for worfe ones.

Besides this advantage which the substance of bones has from the marrow, their articulations are faid (d) to receive no less benefit from it: for it is thought that the marrow passes into the articular cavities through the holes which are in the bones near the large joints. And as a proof of this, it is alleged, that butchers, upon feeing the greater or leffer quantity of marrow in the bones of cows, can tell whether they have travelled far

or little before they were flaughtered.

When the marrow, after having ferved the uses mentioned, is reassumed into the mass of blood, (as it is continually in common with all other fecreted liquors that have not passages formed for conveying them out of the body), it corrects the too great acrimony communicated to the faline particles of our fluids by their circulation and heat, in the fame manner as lixivial falts are blunted by oil in making foap. Hence, in acute diseases, the marrow, as well as the other fat of the body, is quickly wasted, but must be immediately supplied by liquors from the veffels; feeing the cells within the bones, which have no affistance to their contraction from the pressure of the atmosphere, cannot collapse, as the tela cellularis under the skin does when the liquor in its cells is absorbed; the bones therefore are always full.

Since it is the nature of all oil to become thin and rancid when exposed long to heat, and bones have much oil in their firm hard fubstance, we may know why an ungrateful fmell and dark-coloured thin ichor, proceed more from corrupted bones than from other parts of the body; and we can understand the reason of the changes of colour which bones undergo, according to their different degrees of mortification.—Hence likewise we may learn the cause of a spina ventosa, and of the

difficulty

<sup>(</sup>d) Joan. de Muralto Vade-mecum Anat. exercit. 5. § 3. ters Ofteolog. Nov. difc. 3. p. 179.

L

or.

es

to

at

he

s.

p-

ie

ar

es

it

rs

ıt

1-

ir

ts

C

e

)-

n

ė

difficulty of curing all caries of bones proceeding from an obstruction and consequent putresaction of the marrow; and of the quick pulse, thirst, and hectic paroxysms, so often attending these diseases. These phenomena also teach us the reason of the satal prognosis taken from black setid urine in severs.

Though bones so far agree in their structure and annexed parts, yet we may observe a considerable difference among them in their magnitude, figure, situation, substance, connection, uses, &c. From which authors have taken occasion to distinguish them into asmany classes as they could enumerate of these different circumstances. But these being obvious to every perfon that looks on bones, I shall only mention one of them; which comprehends very near the whole bones of the body, and at the same time leads us to examine the most considerable variety that is to be found in the disposition of their constituent parts, and in their uses. It is this, that some bones are broad and slat, while, others are long and round.

#### THE DIFFERENT CLASSES, &c. OF BONES.

THE broad bones have thin fides, by the plates being foon and equally fent off to form the lattice-work; which therefore is thicker, and nearly of an equal form all through. By this structure they are well adapted to their uses, of affording a large enough surface for the muscles to rise from and move upon, and of defending sufficiently the parts which they inclose.

The round bones have thick strong walls in the middle, and become very thin towards their ends; which is owing to very few plates separating at their middle; where, on that account, the cancelli are so sine and small, that they are not taken notice of: but such bones are said to have a large reservoir of oil in this place. Towards their ends the lattice-work becomes very thick, and rather more complete than in the other fort of bones.—These round bones having

Vol. I. C ftrong

strong forces naturally applied to them, and being otherwise exposed to violent injuries, have need of a cylindrical figure to resist external pressure, and of a considerable quantity of oil to preserve them from becoming too brittle. Besides which, they are advantageously provided with thick sides towards their middle, where the greatest forces are applied to injure them; while their hollowness increases their diameter, and consequently their strength to resist forces applied to break them transversely (e). Thus, for instance, in estimating the proportional resistance of two cylindrical bones of unequal diameters, but consisting of an equal number of similar sibres uniformly disposed round each, it is plain,

1. That the absolute force of these two bones is equal, because they consist of equal numbers of similar

fibres.

- 2. That the absolute forces of all the fibres in each bone have the same effect in resisting any power applied to break them, as if the sum of all their forces was united in the respective centres of the transverse sections where the fractures are to be made. For, by hypothesis, the sibres being uniformly disposed in each, there is not any fibre in either bone that has not a corresponding sibre; the sum of both whose distances from the axis of revolution (about which all the parts of the bone must revolve in breaking) is equal to two semi-diameters of the bone: consequently each sibre, and all the sibres, may be regarded as resisting at the distance of one semidiameter or radius from this axis, that is, in the centre.
- 3. Since the united force of all the fibres is to be regarded as resisting at a distance from the centre of motion equal to the semidiameter, it follows, that the total resistance of all these fibres, or the strength of the bone, is proportional to its semidiameter, and consequently to its diameter.

L

g

a

e-

a-

e,

1;

ıd

to in

al

al

h,

15

ar

ch

d

23

ns

e-

re

e-

m

10

ii-

all

ce

s,

e-

0-0-

ne

e-

I

I have here taken for an example one of the most simple cases for calculating the proportional forces of bones. But, were it not too foreign to the present design, it might be universally demonstrated, that of whatever figure bones are, and in whatever manner their fibres are disposed, their strength must always be in a ratio, compounded of the area of their transverse sections, or of their quantity of bony matter, and of the distance of the centre of gravity of these sections from the centre of motion or sulcrum, on which the

bone is supposed to be broken (f).

Since, therefore, the strength of bones depends on their number of fibres, or quantity of matter, and the largeness of their diameters, one may conclude, that the part of a bone formerly fractured, and reunited by a callus, must be stronger than it was before the fracture happened; because both these advantages are obtained by a callus: which is a wife provision, fince bones are never let in fuch a good direction as they were naturally of; and then, wherever a callus is formed, there is fuch an obstruction of the vessels, that if the bone was again broken in the fame place, the offific matter could not so easily be conveyed to reunite it. This callus may indeed, for want of compression, be allowed to form into a fpongy cellular substance (g); but even in this case the strength of the bone is here increased by one or both the causes above-mentioned.

Many bones have protuberances or processes (h) rifing out from them. If a process stands out in a roundish ball, it is called caput or head.—If the head is flatted, it obtains the appellation of condyle.—A rough unequal protuberance is called tuberosity.—When a

(g) Ruysch. Thesaur. 8. n. 49. Mus. anat. thec. B. reposit 2.

n. 2.

<sup>(</sup>f) See the demonstration of this theorem by Dr Porterfield in the Edinburgh medical essays, vol. 1. art. 10.

<sup>(</sup>h) Αποφυσεις, εκφυσεις, εξοχη, προβολη, περοβλημα, Excessus, explanatio, tuberculum, gibbus, eminentia, productio, extuberantia, proctura, enascentias

process rises narrow, and then becomes large, the narrow or small part is named cervix or neck.—Long ridges of bones are called spines.—Such processes as terminate in a sharp point have the general name of coronæ (i) or coronoid bestowed on them; though most of them receive particular names from the resemblance they have, or are imagined to have, to other substances, e. g. mastoid, styloid, anchoroid, coracoid, spinal, &c.—Such processes as form brims of cavities, are called supercilia (k).

Processes serve for the advantageous origin and infertion of muscles, and render the articulations firm and

stable.

Before leaving this subject, we must remark, that much the greater number of what are called processes in adult bones, discover themselves in children to be epiphyses, or distinct bones, which are afterwards united to the other parts: such are the styloid processes of the temporal bones, processes of the vertebræ, trochanters of the thigh, &c. However, as I design to insist chiefly on the description of the adult skeleton, in which the union of these parts is so intimate that scarce any vestige remains of their former separation, I shall retain the common appellation of apophyse, or process, to all such protuberances; but shall remark the principal ones that have no just title to this name, when they occur in the description of particular bones.

On the surfaces of a great many of the bones there are cavities or depressions. If these are deep, with large brims, authors name them cotylæ (1).——If they are superficial, they obtain the designation of glenæ or glenoid. These general classes are again divided into several species:—Of which, pits are small roundish channels sunk perpendicularly into the bone;—furrows, long narrow canals formed in the surface;—nitches or

notches,

(i) Rostra, glandes.

(k) Trues, oppues, außoves, xein, Labra.

<sup>(1)</sup> Κοτυλιδες, οξυβαροι, Acetabula, pixides, buccellæ.

notches, small breaches in the bone;—finuosities, broad, but superficial depressions without brims;—fosse, large deep cavities, which are not equally surrounded by high brims;—finuses, large cavities within the substance of the bones, with small apertures;—foramina, or holes, canals that pierce quite through the substance of the bones.—When this last fort of cavity is extended any long way within a bone, the middle part retains the name of canal, and its ends are called boles.

The cavities allow the heads of bones to play in them; they lodge and defend other parts; they afford fafe passage to vessels, muscles, &c. To mention more would engage us too much in the history of particular bones, which more properly belongs to the demonstration of the skeleton, where we shall have occasion to ob-

ferve these several species of cavities.

t

e

-

e

t

e

1

To far the greater number of bones, whose ends are not joined to other bones by an immoveable articulation, there are smaller ones annexed, which afterwards become scarce distinguishable from the substance of the bone itself. These are called epiphyses or appendices (m). Some bones have one, others have two, three, or four of these appendices annexed by the means of cartilages, which are of a considerable thickness in children, but by age become thinner; the offisication proceeding from the end of the bone on one side, and from the epiphyses on the other, till at last, in adults, the place of their conjunction can scarcely be seen on the external surface; and it is only sometimes that we can then see any mark of distinction in the cancelli (n).

Several processes (e. g. trochanters of the thigh, spine of the scapula, &c.) have epiphyses; and processes frequently rise out from epiphyses; for example, at the

lower end of the femur, unna, tibia, &c. (0).

C<sub>3</sub> The

(0) Vefal. De human. corp. fabrica, lib. 1. cap. 3.

<sup>(</sup>m) Applantatio, additamentum, adnascentia, adnexum, perone.
(n) Winslow, Exposition anatomique de corps humain, traité des os secs, § 116.

The epiphyses are united chiefly to such bones as are destined for frequent and violent motion; and for this purpose they are wisely framed of a larger diameter than the bone they belong to: for, by this means, the furface of contact between the two bones of any articulation being increased, their conjunction becomes firmer, and the muscles inserted into them act with greater force by reason of their axes being further removed from the centre of motion. These advantages might indeed have been obtained by the expansion of the end of the bone itself to a thickness equal to that of the epiphyses; but then the constant separation of new plates to form so wide a cellular structure, must have left the folid fides of the bones fo thin as to yield eafily, either to the action of the muscles fixed to them and passing over them, to the weight several of them are obliged to support, or to the application of any other external force.

Several anatomists (p) thought that the epiphyses ferve other purposes; such as securing the ligaments of the articulations which rife out from between the bones and them; for, as foon as these parts are intimately joined, the ligaments infinuated betwixt them must have a much stronger connection than they could have to the smooth surface of the bones. Such an interception of the ligament between the body of the bone and its epiphyse is not to be seen; but as, at this place, the bone remains longer foft than any where elfe, and the adhesion of the periosteum and of ligaments to bones is always stronger in proportion to the bones being nearest to the consistence of those parts, which is, being foftest, the opinion of these writers, concerning the stronger connection of the ligaments where the bones and epiphyses join, is not without some foundation.

Possibly too, by the fibres of epiphyses not extending

<sup>(</sup>p) Collumb. De re anatomica, lib. 1. cap. 2.— Fallop. Expos. de offibus, cap. 14.

ing themselves so longitudinally as those of the bones, there may be less chance of the former running into each other than of the latter.

The foftness of the ends of bones may be of some advantage in the womb and at birth; after which the offification begins at different points to form epiphyses, before the offification can extend from the middle to the ends of the bones (q).

#### OF OSSIFICATION.

However folid and compact adult bones are, yet they were once cartilages, membranes, nay, a mere jelly. This needs no further proof, than repeated obfervations of embryos when diffected: And how much more tender must the bones be before that time, when neither knife nor eye is capable to discover the least rudiments of them? By degrees they become more folid, then assume the nature of griftles, and at last offify; the cohesion of their plates and sibres always increasing in proportion to their increased solidities; as is evident from the time necessary to unravel the texture of bones of people of different ages, or of dense and of spongy bones, or of the different parts of the same bone, and from the more tedious exsoliations of the bones of adults than of children.

After any part of a bone is fully offified, its fibres are extended little more in length at that part, tho' they increase there in thickness, and though their softer parts

continue to become longer (r).

As the folidity of bones increases, their periosteum more easily separates from them. When bones are membranous, the periosteum and they cannot be distinguished; they appear to be the same substance. When they are cartilages, their membrane adheres so firmly to them, that it is difficult to separate it from

(q) Haller de studio medic. p. 267. (r) Hales's Vegetable statics, p. 293.—Du Hamel, Memoires de l'acad, des sciences, 1742. them. Where the rigid bony fibres are, the periosteum is easily taken off.—Is the similarity of structure and consequent greater attraction of the membrane and substance it incloses, while they are both flexible, the cause of their greater adhesion? or is it owing to the vessels that go from the one to the other being then larger? or do both these causes combine to produce this effect? Or is the membrane or cartilage, which becomes bone asterwards, to be considered as the same substance with the periosteum (s)? and must all these plates of bones be therefore said to be layers of the periosteum hardened (t)?

The offification of bones depends principally on their vessels being so disposed, and of such diameters, as to separate a liquor, which may easily turn into a bony substance, when it is deprived of its thinner parts; as seems plain from the observation of the callous matter separated after fractures and ulcers, where part of the bone is taken out: For, in these cases, the vessels extending themselves, and the liquors added to them, are gradually formed into granulated sless, which sills up all the space where the bone is taken from, then hardens till it becomes as firm as any other part of the bone. This happens frequently, even when the ends of the diseased bone are at a considerable distance from

each other (u).

The induration of bones is also greatly assisted by their being exposed, more than any other parts, to the strong pressure of the great weights they support, to the violent contraction of the muscles fixed to them, and to the force of the parts they contain, which endeavour to make way for their own further growth. By all this pressing force, the solid sibres and vessels of bones are

. thrust

<sup>(</sup>s) Memoires des l'acad. des sciences, 1744. (t) Memoires de l'acad. des sciences, 1743.

<sup>(</sup>u) Hildan. de vuln. gravif. — Med. effays, vol. 1. art. 23.

Job a Meckren, obf. 69. — Mem. de l'acad. des feiences, 1742. — See a collection of fuch cases in Bochmer de offium callo.

I.

m

ofe

Is

or

r

f-

h

es

7-

ir

to

ıy

as er

le

C -

re

r-

ne ds

m

y

ie ie

0

0

18

e st thrust closer, and such particles of the sluids conveyed in these vessels as are sit to be united to the sibres are sooner and more sirmly incorporated with them, while the remaining sluids are forcibly driven out by the veins, to be mixed with the mass of blood. In consequence of this, the vessels gradually diminish as the bones harden. From which again we can understand one reason why the bones of young animals sooner re-unite after a fracture than those of old, and why cattle that are put too soon to hard labour, seldom are of such large size as others of the same brood who are longer kept from labour.

That the offifying of bones greatly depends on preffure, feems to be evinced from the frequent examples we meet with of other parts turning bony, when long exposed to the pressing force of the surrounding parts, or when they are subjected to the like circumstances by their own frequent and violent contraction. Witness the bones found frequently near the base of the heart in fome old men (x), and in several other creatures. Nay, the muscular substance of the heart has been offified in fuch (y), and the arteries of old men often become bony. -The cartilages of the larynx are generally offified in adults.—In beafts of burden, the cartilages between the vertebræ of the back very often change into complete bones; and, being intimately united with the vertebræ, the whole appears one continued bone: Nor is the periofteum exempted from fuch an induration (z).

To confirm this argument still farther, we may obferve, that bones begin their offification at the places where they are most exposed to these causes, viz. in the cylindrical bones, from a middle ring; and in the broad

ones.

<sup>(</sup>x) Riolan. Comment. de offib. cap. 32.—Bartholin. Hift. medic. cent. 1. hift. 50.—Ibid. cent. 2. hift. 45.

<sup>(7)</sup> Cheselden's Anatomy, book 1. introd. —Garengeot, Hist. de l'acad. des sciences, 1726.

<sup>(</sup>z) Peyer. Ephemerid. German. decur. 2. ann. 7. observ. 205.

0

tl

tl

ones, at or near their centre, from one or more diflinct points. The reason of which is, That these parts are contiguous to the bellies of the muscles annexed to the bones, where the fwelling of these moving powers is greatest. What the effects of this may be, let any judge, who view fome of the bones, as the scapula and offa ilium, which are covered with muscles on each side; how compact and thin they are in adults, where the bellies of the muscles were lodged; whereas in children they are thicker. But this being the middle part of these bones, where the greatest number of fibres is, this particular place would have been much thicker in adults, had not this forcible cause been applied, which has not had fuch effects in children, whose muscles have not been much exercised.—Besides, if we allow that all the parts of a bone are equally increased by the constant fupply of new particles, each fibre, and every particle of a fibre, endeavours to make way for its own growth, by pushing the one next to it; and consequently by far the greatest pressure is on the middle, to make the particles firm, and therefore to begin their offification Lastly, the pulsation of the medullary arteries, which enter the bones near to this middle part, may, as authors have alleged, contribute perhaps fomewhat to this induration.

From the effects of pressure only it is that we can account for the bones of old people having their sides much thinner, yet more dense and solid, while the cavities are much larger than in those of young people; and for the prints of muscles, vessels, &c. being so much more strongly marked on the surfaces of the former than of the latter, if they belong to people of near the same condition in life.—Pressure must likewise be the cause, which, in people of equal ages, makes these prints stronger in the bones of those who had much labour and exercise, than they are in people who have led an indolent inactive life.

Perhaps both the causes of offification above-men-

tioned may be affished by the nature of the climate people live in, and the food they use. Whence, in hot countries, the inhabitants sooner come to their height of stature than in the northerly cold regions: And thence seems to have arisen the common practice among the ladies, of making puppies drink brandy or spirit of wine, and of bathing them in these liquors, to prevent their growing big. Nay, it has been observed, that much use of such spirits has occasioned parts, naturally soft, to petrify in some, and to offify in other people of no great age (a).

From the foregoing account of the structure of bones, and of their offisication, we may understand the

reasons of the following phenomena.

How the natural colour of bones may be changed by fome forts of food (b).

Why the bones of some people are so long in harden-

ing, and in others never completely indurate.

Why, in such whose offisication is slow, the bones are generally thicker in proportion to their lengths, especially at their ends; as in the rickets.

How hard firm bones have become foft and pliable

by diseases (c).

Why, in some diseases, epiphyses separate from bones (d), and the ends of fractured bones come as after their fractures appeared to be cuted (e).

How

(a) Littre, Histoire de l'acad. des sciences, 1706. — Geoffroy, Mem. de l'acad. des sciences, 1706.

(b) Philosoph. transact. num. 442. art. 8. num. 443. art. 2. num. 457. art. 4. Mem. de l'acad. des sciences, 1739, 1742.

(c) Histoire de l'acad. des sciences, 1704. Mem. 1722. Gagliardi, Anatom. ossium, cap. 2. observ. 3. — Ephem. Germ. decur. 1. ann. 1. obs. 37. et schol. decur. 2. ann. 7. obs. 212. 235. decur. 3. ann. 2. obs. 3. — Philos. trans. nº 470. § 3. Ibid. vol. 48 § 4. and 44.

(d) Memoires de l'acad. des sciences, 1699. - Diemerbroek, lib. 9.

cap. 19 .- Cowper's Anat. Explic. tab. 96. fig. 1.

(e) Anion's voyage.

C

ti

T

d

b

ga

f

b

C

How bones may waste and diminish (f.)

How bones may become folid all through, without

any appearance of cancelli (g).

How nodes, tophi, and exostoses, happen after the erosion of the external plates of bones in the lues venerea, scurvy, rheumatism, and gout.

How bones exfoliate by the rifing of granulated flesh

from their furface.

How and from what callus is formed after a frac-

ture (b).

Why callus appears to be rather the continued substance of the periosteum than of the bone, while it remains soft and flexible; but seems continued with the bone after it offisses (i).

Why callus is fenfible while it is foft, but becomes

infensible when it hardens.

What occasions sometimes such difficulty in curing fractured bones; or why they never re-unite, though they are reduced, and all proper means towards a cure are used (k).————————————————Are the bones of women with child more tedious in reuniting than those of other people (l)?

Why calluses, after fractures, are sometimes very

thick and protuberant.

What difference there ought to be in the application of bandages to fractures of the bones of old and of young patients.

How bones, remaining long unreduced after a luxa-

tion,

(f) Chefelden's Anat. book 1. introd.—Hist. de l'acad. des sciences, 1700.

(g) Ruysch. Thefaur. 2. arc. 5. thes. 3. loc. 1. num. 5. thes. 9.

numb. 2. not. 3. - Bochmer de callo offium.

(b) Memoires de l'acad. des sciences, 1741. Dehtleef de ofsium callo.

(i) Mem. de l'acad. 1741.

(k) Meckren Observ. medico-chirurg. obs. 71.—Ruysch. Advers. dec. 2. § 2. Observ. anat.-chir. obs. 4.—Van Swieten in Boerhaave Aphor. § 354.

(1) Hildan. centur. 5. obs. 87. et cent. 6. obs. 68. - Philos.

tranfact. nº 494. § 21.

I.

ut

he

e-

ſh

C-

b-

e-

ne

es

e

r

y

n

1-

8

fuch

tion, may have their form so changed as to make their reduction very difficult, if not impossible (m).

Whoever is desirous to know in what time and order each bone and its several parts begin to assume a bony nature, let him consult Kerckingius (n), who gives us the delineations of abortions from three days after conception, and traces the offisication of the bones from three weeks and a month, till the time of the birth: To whom should be added Coiterus (o) and Eyssonius (p). An account of this subject might also be collected out of Ruysch's works, where some of the mistakes committed by the former authors are corrected; and several more particulars, to make the history of the osteogenea more accurate, have since been added by Nesbit (q) and Albinus (r).

I must refer to the authors now quoted for the more curious part of the human osteogeny; not having preparations enough to give such a sull history of it as is done by them. But I shall endeavour to explain the more useful and necessary part of the osteogeny, by subjoining to the description of each bone of an adult, its condition in ripe children; that is, in such as are born at the ordinary time; and shall point out what parts of each are afterwards joined in form of epiphyses. This, with the sollowing general rules, seem to me sufficient for understanding what of this subject is necessary in the practice of physic and surgery.

1. Wherever I mention any parts being cartilaginous, or their being still separable from the other parts of the bone to which they belong, I would be understood to hint, that, about seven or eight years of age,

<sup>(</sup>m) Saltzman, Obs. decur. obs. 6.—Memoires de l'acad. de chirurgie, tom. 2. p. 155.—Bochmer Instit. osteolog. § 596.

<sup>(</sup>n) Anthropograph ichnograph. et osteogenea fœtuum.

<sup>(</sup>o) De offibus fœtus abortivi.

<sup>(</sup>p) De offibus infant. cognoscend. et curand.

<sup>(</sup>q) Human ofteogeny explained.

<sup>(</sup>r) Icones offium fœtus humani; accedit osteogenez brevis hi-

Ch

it

va

de

in

ai

th

h

tl

b

a

f

fuch parts are offified and united to their proper bones, unless when it is said that they are afterwards formed

into epiphyles.

2. Such as become epiphyses are generally offissed at seven or eight years of age; but, being for the most part moistened by synovia, their external surface is still somewhat cartilaginous, and they are not yet united to their bones.

3. At eighteen or twenty years of age the epiphyses are entirely offisied, and have blended their fibres so with the body of the bone, as to make them inseparable

without violence.

The knowledge of this part of the ofteogeny I think necessary, to prevent dangerous mistakes in the cure of feveral diseases. As for example: Without this knowledge, the separation of an epiphyse might be mistaken for a fracture or luxation.—The interstice of two parts of a bone not yet joined, might be judged to be a fiffure.—A diastasis, or separation of such disjoined pieces of a bone, might be thought a fracture.—The protrusion of one piece, or its overlopping any other, could be mistaken for an excrescence or exostosis.-Such errors about the nature of a difease would give one very different indications of cure from what he would have if he really understood his patient's case: And very often the knowledge of the different inequalities on the furfaces of bones, must direct us in the execuof what is proper to be done to cure feveral of their difeases.

### OF THE CONNECTION OF BONES.

HAVING thus confidered the bones when fingle, we ought next to show the different manner of their conjunctions (s). To express these, anatomists have contrived a great number of technical terms; about the meaning, propriety, and classing of which, there has

<sup>(</sup>s) Συνταζιε, συνδεσιε, συμβολη, ομιλια, Compositio, connexio, articulatio, conjunctio, nodus, commissura, structura, compages.

unluckily been variety of opinions. Some of these terms it is necessary to retain, fince they serve to express the various circumstances of the articulations, and to understand the writers on this subject

The ARTICULATIONS are most commonly divided into three classes, viz. Symphysis, Synarthrosis, and Di-

arthrofis

I.

ed,

ed

ed

oft

ill

es

fo le

k

of

V-

n

0

a

)-

۲,

e

d

d

\$

r

I. Symphysis, which properly fignifies the concretion or growing together of parts, when used to express the articulations of bones, does not seem to comprehend, under the meaning generally given to it, any thing relating to the form or motion of the conjoined bones; but by it most authors only denote the bones to be connected by some other substance; and as there are different substances which serve this purpose, therefore they divide it into the three following species.

ing substance: thus the ribs are joined to the sternum; thus the bodies of the vertebræ are connected to each

other; as are likewise the offa pubis.

2. Synneurosis, or syndesmosis, when ligaments are the connecting bodies, as they are in all the moveable articulations.

3. Syffarcosis, when muscles are stretched from one bone to another, as they must be where there are

moveable joints.

II. The fecond class of articulations, the SYNAR-THROSIS, which is said to be the general term by which the immoveable conjunction of bones is expressed, is divided into three kinds.

1. The future (u) is that articulation where two bones are mutually indented into each other, or as if they were fewed together; and is formed by the fibres of two bones meeting while they are yet flexible and yielding, and have not come to their full extent of growth; fo that they mutually force into the interstices of each other,

till,

(u) Papn.

<sup>(</sup>t) Amphiarthrofis.

CI

fta

ra

ny it

th

at

ot

18

VC

fa

th

ti

C

T

it

11

f

t

h

C

F

(

till, meeting with fuch resistance as they are not able to overcome, they are stopped from sprouting out farther, or are reslected; and therefore these indentations are very different both in sigure and magnitude: thus the bones of the head are joined; thus epiphyses are joined to the bones, before their full connection and union with them.

Under this title of future, the harmonia of the ancients may be comprehended; fcarce any unmoved bones being joined by plain furfaces (x).

2. Gomphosis (y) is the fixing one bone into another, as a nail is fixed in a board: thus the teeth are secured in their sockets.

3. Schindylesis or ploughing (z), when a thin lamella of one bone is received into a long narrow furrow of another: thus the processus azygos of the sphenoid, and the nasal process of the ethmoid bone, are received by the vomer.

III. The third class, or DIARTHROSIS (a), is the articulation where the bones are so loosely connected as to allow large motion. This is subdivided into three kinds.

1. Enarthrofis, or the ball and focket, when a large head is received into a deep cavity; as the head of the os femoris is into the acetabulum coxendicis.

2. Arthrodia, when a round head is received into a fuperficial cavity; as in the articulation of the armbone and scapula. These two species of diarthrosis allow motion to all sides.

3. Ginglimus (b), which properly fignifies the hinge of a door or window; in it the parts of the bones mutually receive and are received, and allow of motion two ways: Workmen call it charnal.

The

<sup>(</sup>x) Vefal. observ. Fallop. examen.

<sup>(</sup>y) Conclavatio.

<sup>(</sup>z) Keil's Anat. chap. 5. fect. 3.

<sup>(</sup>a) Aπαρθρωσις, Dearticulatio, abarticulatio.

<sup>(</sup>b) Articulatio mutua.

I.

to

r,

re

ne

n-

n

n-

d

r,

d

la

of

d

y

1-

0

e

e

a

-

e

n

e

The ginglimus is generally divided into three kinds; to which some (c) give the names of contiguous (d), difant (e), and compound (f).

The first kind of ginglimus is when a bone has several protuberances and cavities, which answer to as many cavities and processes of the other bone with which it is articulated; as in the conjunction of the femur with the tibia.

The fecond species is, when a bone receives another at one end, and is received by the same bone at the other end; as in the radius and ulna.

The last fort is, when a bone receives another, and is received by a third; as in the oblique processes of the vertebræ.

When I first mentioned the articulations of bones, I faid there were different opinions concerning the use of their technical names; e. g. It has been faid, that fymphysis should be the name for the immoveable articulations, and fynarthrofis should be understood to be the conjunction of bones by some connecting medium. Those who have taken symphysis in the sense I did, of its expressing the conjunction of bones with a connecting substance, have disagreed in their definition of it; fome inferting, and others leaving out, its allowing mo-Where they have agreed in their definition, they have not been of the fame mind concerning the species of it: for several think the sysarcosis and syndesmosis applicable to fo many joints which are univerfally allowed to be classed under the diarthrofis, that it must create contusion to name them by any species of the symphysis. Few keep to fuch a general definition of the fynchondrofis as I have done; and whether they determine it to allow no motion, or an obscure or a manifest one, they bring themselves into difficulties, because there are examples VOL. I.

<sup>(</sup>c) Baker, Curf. Ofteolog. demonstr. 1.

<sup>(</sup>d) Proximus. (e) Longus.

<sup>(</sup>f) Compositus.

amples of all these three kinds.—Some again, by too nicely distinguishing obscure and manifest motions of bones, have blended the *synarthrosis* and *diarthrosis*; and from thence have branched out the different compound species of articulations that may be formed of them, so far, that they could find no examples in the body to illustrate them by.—It would be tedious to enumerate more of the jarring opinions, and it would be far more so to give a detail of the arguments used by the disputants. It is sufficient for my purpose that it is understood in what sense I take these technical terms; which I do in the following manner.

When I mention the fymphysis or fynarthrosis, or any species of them, I shall always understand them according to the explication already given of them. But though the preceding account of the diarthrosis, or articulation of moveable bones, has been almost universally received; yet seeing it does not comprehend all the moveable articulations of the body, and one of its species does not answer to any notion we can have of the conjunction of two bones, I must beg leave to

change the definitions and kinds of these joints.

I would call diarthrofis, That conjunction of bones whereby they are fitted for motion, being each covered with a fmooth cartilage, connected by one or more common ligaments, and lubricated with liquor at the conjoined parts: In which definition I have no regard to the quantity of motion which they really do perform; the motion being often confined or enlarged by some other cause not immediately depending on the frame of the two surfaces of the bones forming the particular joint which then is considered.

The first species of the diarthrosis, viz. the enarthrofis, or ball and socket, I would define more generally than above, That articulation where a round head of one bone is received into a cavity of another; and consequently, without some foreign impediment, is capable of motion to all sides. Examples of this kind I.

00

of

is;

m-

of

he

to

ıld

ed

nat

cal

or

m

m.

or

ni-

nd

of

ve

to

ies

ed

re

he

re-

do

ed

he

ar-

ro-

lly

of

n-

a-

nd

re

is

are to be seen in the articulation of the thigh-bone and offa innominata; arm-bone and scapula; astragalus and os naviculare; magnum of the wrist with the scaphoides and lunare; first bone of the thumb with the second, &c.

The fecond fort, or the arthrodia, differing from the enarthrofis, in the preceding account, only in the cavity's being more superficial, (which makes no effential difference, especially that, in the recent subject, cartilages or ligaments supply the deficiency of bone), ought, in my opinion, to be called with Vefalius (g), that articulation of two bones adapted for motion, where it is not at first fight obvious which of the two has the head or cavity, or where they are joined by plain furfaces, or nearly fo: fuch is the conjunction of the clavicle with the scapula; offa cuneiformia with the os naviculare; metatarfal bones with the offa cuneiformia, &c. From the nature of this fort of joint, it is plain, that very great motion cannot be allowed, without the bones going further out of their natural fituation than is convenient or fafe.

Ginglimus, I would reckon that articulation by the form of which the motion of the joined bones must be chiefly confined to two directions, which hinges of doors are.

The first species of this is the trochoides, when one bone turns on another, as a wheel does on its axis. Thus the first vertebra of the neck moves on the tooth-like process of the second. This is the most proper kind of ginglimus.

The fecond species should be esteemed that articulation where several prominent and hollow surfaces of two bones move on each other within the same common ligament; as in the knee, elbow, &c.

The third fort of ginglimus is, when two bones are articulated to each other at different parts, with a diffinct apparatus of the motory machines at each: fuch

D 2

(g) De corp. human. fabrica, lib. 1. cap. 4.

is the articulation of the os occipitis with the first vertebra of the neck; of any two contiguous vertebræ, by their oblique processes; of the ribs with the bodies and transverse processes of the vertebræ; of the radius with the ulna, tibia with the sibula, astragalus with the cal-

caneum, &c.

I would entirely throw out what is commonly called the third kind of ginglimus: for in examining the conjunction of a bone with two others, as in the common example of a vertebra joined with the one above and below, the connection of the middle one with each of the other two ought to be confidered separately: otherwise we might, with the same propriety, esteem the articulations that the long bones, the femur, tibia, humerus, &c. have at their different ends, as one articu-

lation: which is abfurd.

If the moveable bones are not connected and kept firm by some strong substance, they would be luxated at every motion of the joints; and if their hard rough unequal furfaces were to play on each other, their motion would not only be difficult, but the loss of substance from attrition would be great. Therefore ligaments are made to obviate the first, and cartilages to prevent the other inconveniency. But because ligaments and cartilages turn rigid, inflexible, and rough, unless they are kept moift, a fufficient quantity of proper liquors is fupplied for their lubrication, and to preserve them in a Seeing, then, these parts are so necessary flexible state. to the articulations, I shall next confider their structure, fituation, and uses, so far as they are subservient to the bones and their motions.

### OF THE LIGAMENTS.

LIGAMENTS (b) are white flexible bodies, thicker and firmer than membranes, and not so hard or firm as cartilages, without any remarkable cavity in their substance, difficultly stretched, and with little elasticity; ferving

<sup>(</sup>h) Duvdes uoi, viupoi, copulæ, vincula.

I.

e-

by

nd

th

al-

ed

n-

on

nd

of

r-

he

u-

u-

pt

ed

gh

0-

ce

its

nt

nd

ey

18

a

ry

e,

he

er

m

ir

y ;

g

ferving to connect one part to another, or to prevent the parts to which they are fixed from being removed out of that fituation which is useful and fafe.

After maceration in water, the ligaments can eafily be divided; and each ligamentous layer appears composed of fibres, the largest of which are disposed in a longitudinal direction.

The arteries of ligaments are very confpicuous after a tolerable injection, and the larger trunks of their veins are fometimes to be feen full of blood.

Such ligaments as form the fides of cavities, have numerous orifices of their arteries opening upon their internal furface, which keep it always moift. If we rub off that moisture, and then press the ligament, we can fee the liquor oozing out from small pores; and we can force thin liquors injected by the arteries into the cavities formed by ligaments.

These exhalent arteries must have corresponding abforbent veins, otherwise the cavities would foon be too full of liquor.

Ligaments, then, must be subject to the diseases common to other parts, where there is a circulation of fluids, allowance always being made for the fize of vessels, nature of the sluids, and firmness of the texture of each part.

Authors generally fay that ligaments are infensible; and confequently it may be inferred, that they have no nerves bestowed on them. But the violent racking pain felt on the least motion of a joint labouring under a rheumatism, the seat of which disease seems often to be in the ligaments, and the infufferable torture occafioned by incifions of ligaments, and by a collection of acrid matter in a joint, or by tophi in the gout, would perfuade us that they are abundantly supplied with nerves.

The ligaments which connect the moveable bones commonly rife from the conjunction of the epiphyles of the one bone, and are inferted into the same place of the

the other; or, where epiphyses are not, they come out from the cervix, and beyond the supercilia of the articulated bones; and after such a manner, in both cases, as to include the articulation in a purse or bag; with this difference, depending on their different motions, that where the motion is only to be in two directions, the ligaments are strongest on those sides towards which the bones are not moved; and when a great variety of motions is designed to be allowed, the ligaments are weaker than in the former sort of articulations, and are nearly of the same strength all round.

Part of the capfular ligaments is composed of the periosteum, continued from one bone to another, as was observed p. 18. and their internal layer is continued on the parts of the bone or cartilage which the ligament

includes (i).

Besides these common capsular ligaments of the joints, there are particular ones in several places, either for the sirmer connection of the articulated bones, or for restraining and confining the motion to some one side; such are the cross and lateral ligaments of the

knee, the round one of the thigh, &c.

From this account of the ligaments, we may conclude, that, cateris paribus, in whatever articulation the ligaments are few, long, and weak, the motion is more free and quick; but luxations happen frequently: and, on the contrary, where the ligaments are numerous, short, and strong, the motion is more confined; but such a joint is less exposed to luxations (k).—Whence we may judge how necessary it is to attend to the different ligaments, and the changes which have been made on them by a luxation, when it is to be reduced.

Ligaments also supply the place of bones in several cases to advantage. Thus the parts in the pelvis are more safely supported below by ligaments than they could

<sup>(</sup>i) Nesbit, Osteogen.—Philos. transact. No 470. § 6.
(k) Fabric. ab Aquapend. de articul. part. utilit. pars 3.

I.

ıt

i-

a-

;

)-

C-

is

1-

1-

1-

e

IS

d

it

e

r

r

e

e

could have been by bone.—The ligaments placed in the great holes of the offa innominata, and between the bones of the fore-arm and leg, afford convenient origin to muscles.—Immoveable bones are firmly connected by them; of which the conjunction of the os facrum and innominatum is an example.—They afford a focket for moveable bones to play in, as we see part of the astragalus does on the ligament stretched from the heel-bone to the scaphoid.

Numerous inconveniences may arise from too long or short, strong or weak, lax or rigid ligaments.

### OF THE CARTILAGES.

CARTILAGES (1) are folid, smooth, white, elastic substances, between the hardness of bones and ligaments, and covered with a membrane named perichondrium; which is of the same structure and use to them as the periosteum is to the bones.

Cartilages are composed of plates, which are formed of fibres, disposed much in the same way as those of bones are; as might be reasonably concluded from observing bones in a cartilaginous state before they offify, and from seeing, on the other hand, so many cartilages become bony. This may be still further confirmed by the exsoliation which cartilages are subject to as well as bones.

The perichondrium of feveral cartilages, for example those of the ribs and larynx, has arteries, which can be equally well injected with those of the periosteum; but the vessels of that membrane in other parts, e. g. the articular cartilages, are smaller, and in none of them does injection enter deep into the substance of the cartilages; nay, madder, mixed with the food of animals, does not change the colour of cartilages as it does that of bones (m).

D 4 The

<sup>(1)</sup> XovSpoi.
(m) Philof. Transact. No 442. art. 8. No 443. art. 2.

The granulated flesh which rises from the ends of metacarpal or metatarsal bones, when the cartilage exfoliates, after a singer or toe has been taken off at the first joint, is very sensible; from which the existence

of nerves in cartilages may be inferred.

While cartilages are in a natural state, it is to be remarked, first, That they have no cavity in their middle for marrow. Secondly, That their outer surface is softest, which renders them more slexible. Thirdly, That they do not appear to change their texture near so much by acids as bones do. And, lastly, That as the specific gravity of cartilages is near a third less than that of bones; so the cohesion of their several plates is not so strong as in bones: Whence cartilages laid bare in wounds or ulcers, are not only more liable to corrupt, but exsoliate much sooner than bones do.

Cartilages feem to be principally kept from oflifying, either by being subjected to alternate motions of flexion and extension, the effects of which are very different from any kind of fimple pressure; or by being constantly moistened (n). Thus the cartilages on the articulated ends of the great bones of the limbs, and the moveable ones placed between the moving bones in fome articulations, which are obliged to fuffer many and different flexions, and are plentifully moistened, fcarce ever change into bone; while those of the ribs and larynx are often offified.—The middle angular part of the cartilages of the ribs, which is constantly in an alternate state of flexion and extension, by being moved in respiration, is always the last of becoming bony.—In the larynx, the epiglottis, which is oftener bended and more moistened than the other four cartilages, feldom is offified, while the others as feldom escape it in adults.

The cartilages subservient to bones are sometimes found

Nº 457. art. 4 - Mem. de l'acad des sciences, 1739 et 1742. - Dehtleef de ossium callo.

<sup>(</sup>n) Havers Ofteolog. Nov.

found on the ends of bones which are joined to no other; but are never wanting on the ends and in the cavities of such bones as are designed for motion (o). Cartilages also are interposed between such other cartilages as cover the heads and cavities of articulated bones; nay, they are also placed between immoveable bones.

The uses of cartilages, so far as they regard bones, are, to allow, by their fmoothness, such bones as are defigned for motion, to flide eafily without detrition, while, by their flexibility, they accommodate themfelves to the feveral figures necessary in different motions, and, by their elasticity, they recover their natural position and shape as soon as the pressure is removed. This fpringy force may also affift the motion of the joint to be more expeditious, and may render fhocks in running, jumping, &c. lefs .- To these cartilages we chiefly owe the fecurity of the moveable articulations: for, without them, the bony fibres would fprout out, and intimately coalefce with the adjoining bone; whence a true anchylofis must necessarily follow; which never fails to happen when the cartilages are, eroded by acrid matter, or offified from want of motion or defect of liquor, as we fee often happens after wounds of the joints, paidarthrocace, scrophula, and spina ventosa, or from old age, and long immobility of joints (p).—Hence we may know what the annihilation is, which is faid to be made of the head of a bone, and of the cavity for lodging it, after an unreduced The moveable cartilages interpofed in fracture (q). joints ferve to make the motions both freer and more fafe than they would otherwise be.—Those placed on the ends of bones that are not articulated, as on

(0) Cels. de re medic. lib. 8. cap. 1.

<sup>(</sup>p) Columb. de re anat. lib. 15.—Deslandes, Hist. de l'acad. des sciences, 1716—Phil. transact. N° 215.—Ibid. N° 461. sect. 16.

<sup>(</sup>q) Hildan. de ichor. et melicer. acri Celfi, cap. 5.—Ruysch. Thes. 8. N° 103.—Saltzman, in Act. Petropolit. tom. 3. p. 275.

C

lo

b

01

m

fa

b

n

m

d

0

li 0

ta

fe

W

m

fu

ci

CI

W

ti

li

fe

ir

p

a

0

the spine of the os ilium, base of the scapula, &c. serve to prevent the bony fibres from growing out too far.— Cartilages fometimes ferve as ligaments, either to fasten together bones that are immoveably joined, fuch are the cartilages between the os facrum and offa ilium, the offa pubis, &c. or to connect bones that enjoy manifest motion, as those do which are placed between the bodies of the true vertebræ, &c. --- Cartilages very often do the office of bones to greater advantage than these last could; as in the cartilages of the ribs, those which supply brims to cavities, &c.

Too great thickness or thinness, length or shortness, hardness or suppleness, of cartilages, may therefore

cause great disorders in the body.

### OF THE SYNOVIA.

THE liquor which principally ferves to moisten the ligaments and cartilages of the articulations is supplied by glands, which are commonly fituated in the joint, after such a manner as to be gently pressed, but not destroyed by its motion. By this means, when there is the greatest necessity for this liquor, that is, when the most frequent motions are performed, the greatest quantity of it must be separated. These glands are foft and pappy, but not friable: in some of the large joints they are of the conglomerate kind, or a great number of small glandules are wrapt up in one common membrane. Their excretory ducts are long, and hang loofe, like fo many fringes, within the articulation; which, by its motion and pressure, prevents obstructions in the body of the gland or its excretories, and promotes the return of this liquor, when fit to be taken up by the absorbent vessels, which must be in the joints, as well as in the other cavities of the body; and, at the same time, the pressure on the excretory ducts hinders a superfluous unnecessary secretion, while the fimbriated disposition of these excretories does not al-

low

L

C

n

c

1,

1-

n

es

1-

e

s,

e

ie

)-

le

ıt

n s,

e

ls

le

a

le

1-

ts

s,

e

e

1,

ts

e

lw low any of the secreted liquor to be pushed back again

by these canals towards the glands (r).

Very often these fountains of slimy liquor appear only as a net-work of vessels.—Frequently they are almost concealed by cellular membranes containing the sat—and sometimes small simple mucous solliculi may be seen (s).

The different joints have these organs in different numbers and fizes: the conglomerate ones do not vary much, especially as to situation, in the similar joints of different bodies; but the others are more uncertain.

Upon pressing any of these glands with the singer, one can squeeze out of their excretories a mucilaginous liquor, which somewhat resembles the white of an egg or serum of the blood; but it is manifestly salt to the taste. It does not cogulate by acids or by heat, as the serum does; but by the latter turns first thinner, and, when evaporated, leaves only a thin salt film.

The quantity of this mucilage, constantly supplied, must be very considerable, since we see what a plentiful troublesome discharge of glary matter follows a wound or ulcer of any joint: of which liquor the mu-

cilage is a confiderable part.

The vessels which supply liquors for making the secretion of this mucilage, and the veins which bring back the blood remaining after the secretion, are to be seen without any preparation; and, after a tolerable injection of the arteries, the glands are covered with them.

In a found state, we are not conscious of any sensibility in those glands: but, in some cases which I have seen, when they inslame and suppurate, the most racking pain is selt in them; a melancholy, though a sure proof that they have nerves.

These mucilaginous glands are commonly lodged in a cellular substance; which is also to be observed in other parts of the bag formed by the ligaments of the

arti-

(s) Morgagn. Adversar. 2. animad. 23.

<sup>(</sup>r) Cowper, Anatom. explicat. tab. 79. lit. E. E.

ru

th

it

bo

to

ti

al

cl

e

to

fo

te

W

h

e

i

P

d

t

E

t

r

articulation; and contains a fatty matter, that must necessarily be attenuated, and forced through the including membranes into the cavity of the joint, by the pressure which it suffers from the moving bones.

If, then, the oil is conveyed from this cellular fubstance; and if the attenuated marrow passes from the cancelli of the bones by the large pores near their ends, or in their cavities, and fweats through the cartilages there into the articulations; which it may, when affifted by the constant heat and action of the body, more easily do, than when it escapes through the compact substance of the bones in a skeleton: If, I say, this oil is fent to a joint, and is incorporated with the mucilage, and with the fine lymph that is constantly oozing out at the extremities of the small arteries distributed to the ligaments, one of the fittest liniments imaginable must be produced; for the mucus diluted by the lymph contributes greatly to its lubricity, and the oil preserves it from hardening. How well fuch a mixture ferves the purpose it is designed for, Boyle (t) tells us he experienced in working his air-pump; for the fucker could be moved with much less force after being moistened with water and oil, than when he used either one or other of these liquors: And I believe every one, at first view, will allow the diluted mucilage to be much preferable to simple water. The synovia (u), as this liquor, composed of oil, mucilage, and lymph, is commonly now called, while in a found state, effectually preferves all the parts concerned in the articulations foft and flexible, and makes them flide eafily on each other, by which their mutual detrition and overheating is prevented, in the manner daily practifed in coach and cart wheels by befinearing them with greafe and tar.

After the liquor of the articulations becomes too thin and unferviceable, by being constantly pounded and rub-

<sup>(</sup>t) Physico-mechanic. experim.

<sup>(</sup>u) Muζa, mucus, axungia.

rt I.

ne-

clu-

the

ub-

the

ds,

ges

ift-

ore

act

oil

ıci-

ing

ted

ble

iph

ves

ves

ex-

cer

ist-

ne

at

ich

li-

m-

re-

nd

by

nt-

art

in

nd

h-

rubbed between the moving bones, it is reassumed into the mass of blood by the absorbent vessels.

When the fynovia is not rubbed betwixt the bones, And fometimes, when the head of a it inspissates. bone has been long out of its cavity, this liquor is faid to fill up the place of the bone, and hinder its reduction; or if a joint continues long unmoved, it is also faid to cement the bones, and occasion a true anchylofis (x).——If the fynovia becomes too acrid, it erodes the cartilages and bones; as frequently happens to those who labour under the lues venerea, scurvy, fcrophula, or spina ventosa.——If this liquor is separated in too small quantity, the joint becomes stiff; and, when with difficulty it is moved, a crackling noise is heard, as people advanced in years frequently experience (y).——If the mucilage and lymph are deposited in too great quantity, and the absorbent vessels do not perform their office fufficiently, they may occasion a dropfy of the joints (z).—From this fame cause also the ligaments are often fo much relaxed, as to make the conjunction of the bones very weak: Thence arise the luxations from an internal cause, which are easily reduced, but difficultly cured (a).——Frequently, when fuch a fuperfluous quantity of this liquor is pent up, it becomes very acrid, and occasions a great train of bad fymptoms; fuch as fwelling and pain of the joints, long finuous ulcers, and fiftulæ, rotten bones, immobility of the joints, marcor and atrophia of the whole body, hectic fevers, &c. (b).—From a depravity in the blood, or diseases in the organs that furnish the fynovia of the joints, it may be greatly changed from its natural state; it may be purulent after inflammation,

<sup>(</sup>x) Pare, Chirurgie, livre 15. chap. 18. et livre 16. chap. 5.

<sup>(</sup>y) Galen de usu part. lib. 12. cap. 2.—Fabric. ab Aquapend. de articul. part. utilitat. pars 3.—Bartholin. hist. medic. cent. 3.

<sup>(</sup>z) Hildan. de ichore et melicer. acri Celfi.

<sup>(</sup>a) Hippocrat. de locis in homine, sect. 14. et de articul.

<sup>(</sup>b) Hildan. de ichore et meliceria acri Celsi.

mation, mucous in the white swellings, gelatinous in the rheumatism, chalky from the gout, &c.: hence a great variety of disorders in the joints (c):

(c) See Reimar, Differt. de fungo articulor.

# C H A P. II.

## Of the SKELETON.

Though any dry substance may be called skeleton, yet, among anatomists, this word is universally understood to signify the bones of animals connected together, after the teguments, muscles, bowels, glands,

nerves, and veffels, are taken away (a).

A skeleton is said to be a natural one when the bones are kept together by their own ligaments; and it is called artificial when the bones are joined with wire, or any other substance which is not part of the creature to which they belonged. Small fubjects, and fuch whose bones are not fully offished, are commonly prepared the first way; because, were all their parts divided, the nicest artist could not rejoin them, by reason of their smallness, and of the separation of their unoffified parts; whereas the bones of large adult animals are foonest and most conveniently cleaned when single, and are easily restored to and kept in their natural situation.—Sometimes the skeleton of the same animal is prepared in both these ways; that is, the smaller bones are kept together by their natural ligaments, and the larger ones are connected by wires or some such substances.

Be-

Ch

(cr

tha

lar the pe fal ac ax

> ne fa

> > w fu

B

b

t

d

C

t L

in

e a

ly

d

s,

e

d

h

Ĉ

d

ý

I

Before we proceed to the division and particular description of the skeleton, it is worth while to remark, that, when the bones are put into their natural fituation, scarce any one of them is placed in a perpendicular bearing to another; though the fabric composed of them is fo contrived, that, in an erect posture, a perpendicular line, from their common centre of gravity, falls in the middle of their common base (b). On this account, we can support ourselves as firmly as if the axis of all the bones had been a straight line perpendicular to the horizon; and we have much greater quickness, ease, and strength, in several of the most necesfary motions we perform. It is true, indeed, that wherever the bones, on which any part of our body is fustained, decline from a straight line, the force required in the muscles to counteract the gravity of that part is greater than otherwise it needed to have been: But then this is effectually provided for in fuch places, by the number and strength of the muscles. So long, therefore, as we remain in the same posture, a considerable number of muscles must be in a constant state of contraction; which we know, both from reason and experience, must soon create an uneasy sensation. This we call being weary of one posture: An inconvenience that we should not have had in standing erect, if the bearing of all the bones to each other had been perpendicular; but it is more than compensated by the advantages above-mentioned.

The human skeleton is generally divided into the HEAD, the TRUNK, the SUPERIOR and the INFERIOR

EXTREMITIES.

## SECT. I. OF THE HEAD.

BY the HEAD is meant all that spheroidal part which is placed above the first bone of the neck. It therefore comprehends the cranium and bones of the face.

§ 1. The

<sup>(</sup>b) Cowper, Anat. of human bodies, explic. of tab. 87, 88.

Cl

are

viz ad

gr

to

is

di

ta

ar

th

fi

0

to

tl

C

n

f

P

### 6 1. The CRANIUM.

THE cranium (c), helmet, or brain-case, consists of several pieces, which form a vaulted cavity, for lodging and defending the brain and cerebellum, with their

membranes, veffels, and nerves.

The cavity of the cranium is proportioned to its contents. Hence such a variety of its size is observed in different subjects; and hence it is neither so broad nor so deep at its fore-part, in which the anterior lobes of the brain are lodged, as it is behind, where the large posterior lobes of the brain, and the whole cerebellum, are contained.

The roundish figure of the skull, which makes it more capacious, and better able to defend its contents from external injuries, is chiefly owing to the equal pressure of these contained parts as they grow and increase before it is entirely offisied.—It is to be obferved, however, that the fides of the cranium are depreffed below a spherical surface by the strong temporal muscles, whose action hinders here the uniform protrufion of the bones, which is more equally performed in other parts where no fuch large muscles are. dren, whose muscles have not acted much, and consequently have not had great effects on the bones, this depression is not so remarkable; and therefore their heads are much rounder than in adults. These natural causes, differently disposed in different people, produce a great variety in the shapes of skulls, which is still increafed by the different management of the heads of children when very young: So that one may know a Turk's skull by its globular figure, a German's by its breadth and flatness of the occiput, Dutch and English by their oblong shapes, &c. (d). Two advantages

(d) Vefal. lib. 1. c. 5.

<sup>(</sup>c) Κογχος, κυτος, κωδεια, σκαφιον, calva, calvaria, cerebri galea, theca et olla capitis, testa capitis, soutella capitis.

are reaped from this flatness of the sides of the cranium, viz. the enlargement of our sphere of vision, and more advantageous situation of our ears, for receiving a greater quantity of sound, and for being less exposed

to injuries.

rt I.

of

ing

eir

on-

in

nor

of

rge

m,

it

nts

ual

in-

b-

le-

ral

u-

in

il-

fe-

nis

eir

ral

ce

n-

of

a

its

es

re

a,

The external furface of the upper part of the cranium is very smooth and equal, being only covered with the periosteum, (common to all the bones, but in the skull distinguished by the name of perioranium), the thin frontal and occipital muscles, their tendinous aponeurosis, and with the common teguments of the body; while the external surface of its lower part has numerous risings, depressions, and holes, which afford convenient origin and insertion to the muscles that are connected to it, and allow safe passage for the vessels and nerves

that run through and near it.

The internal furface of the upper part of the skull is commonly fmooth, except where the vessels of the dura mater have made furrows in it, while the bones were foft.—Surgeons should be cautious when they trepan here, left, in fawing or raifing the bone where fuch furrows are, they wound these vessels.——In the upper part of the internal furface of feveral skulls, there are likewise pits of different magnitudes and figures, which feem to be formed by fome parts of the brain being more luxuriant and prominent than others. these pits are, the skull is so much thinner than any where elfe, that it is often rendered diaphanous, the two tables being closely compacted without a diploë; the want of which is supplied by vessels going from the dura mater into a great many small holes observable in the pits. Thefe veffels are larger, and much more confpicuous than any others that are fent from the dura mater to the skull; as evidently appears from the drops of blood they pour out, when the skull is raised from the dura mater in a recent subject; and therefore they may furnish a sufficient quantity of liquors necessary to prevent the brittleness of this thin part. The know-VOL. I.

C

os

ai

tl

0

fi

P

Sa

fi

n

r

fi

n

t

ledge of these pits should teach surgeons to saw cautiously and slowly through the external table of the skull, when they are performing the operation of the trepan; since, in a patient whose cranium has these pits, the dura mater and brain may be injured, before the instrument has pierced near the ordinary thickness of a table of the skull.—The internal base of the skull is extremely unequal, for lodging the several parts and appendices of the brain and cerebellum, and allowing passage and defence to the vessels and nerves that go into or come out from these parts.

The bones of the cranium are composed of two tables and intermediate cancelli, commonly called their diploë (e). The external table is thickest; the inner, from its thinness and consequent brittleness, has got the name of vitrea. Whence we may see the reason of those mischievous consequences which so often attend a collection of matter in the diploë, either from an external or internal cause, before any sign of such a collection appears in the teguments that cover the part

of the skull where it is lodged (f).

The diploë has much the fame texture and uses in the

skull, as the cancelli have in other bones.

The diploë of several old subjects is so obliterated, that scarce any vestige of it can be seen; neither is it observable in some of the hard craggy bones at the base of the skull. Hence an useful caution to surgeons who trust to the bleeding, want of resistance, and change of sound, as certain marks in the operation of the trepan, for knowing when their instrument has sawed through the first table, and reached the diploë (g). In other people, the diploë becomes of a monstrous thickness, while the tables of the skull are thinner than paper.

The cranium confilts of eight bones, fix of which are faid to be proper, and the other two are reckoned

com-

<sup>(</sup>e) Meditullium, commissura.

<sup>(</sup>f) Boneti Sepulchret. anat. lib. 1. § 1. obf. 96.—103. (g) Bartholin. Anat. reform. lib. 4. cap. 4.

tI.

au-

the the

efe

ore

ess

the

rts w-

hat

wo

led

in-

nas

ea-

en

m

na

art

he

d,

it

ife ho

ge

e-

ed

In

k-

r.

ch

ed

n-

This

common to it and to the face.—The fix proper are the os frontis, two offa parietalia, two offa temporum, and the os occipitis.—The common are the os ethmoides and sphenoides!

The os frontis forms the whole fore-part of the vault; the two offa parietalia form the upper and middle part of it; the offa temporum compose the lower part of the sides; the os occipitis makes the whole hinder part, and some of the base; the os ethmoides is placed in the fore-part of the base, and the os sphenoides is in the middle of it.

### THE SUTURES.

THE above bones are joined to each other by five futures; the names of which are the coronal, lambdoidal, faggital, and two fquamous.

The coronal (b) future is extended over the head, from within an inch or so of the external canthus of one eye, to the like distance from the other; which being near the place where the ancients wore their vittæ, coronæ, or garlands, this suture has hence got its name.

—Though the indentations of this suture are conspicuous in its upper part, yet an inch or more of its end on each side has none of them; for it is squamous and smooth there.

The lambdoidal (i) future begins some way below, and farther back than the vertex or crown of the head, whence its two legs are stretched obliquely downwards, and to each side in form of the Greek letter A, and are now generally said to extend themselves to the base of the skull: but formerly anatomists (k) reckoned the proper lambdoidal suture to terminate at the squamous sutures; and what is extended at an angle down from that on each side, where the indentations are less conspicuous than in the upper part of the suture, they called additamentum sutura lambdoidis (l).

(b) Στιφαναία, Arcualis, puppis.

(i) Laudæ, proræ, hypfyloides. (k) Vesal. Anat. lib. 1. cap. 6.

(1) Lambdoides harmonialis, lambdoides inferior, occipitis corona.

Ch

mi

ter

lan

pla

Ho

the

no

tha

me

the

are

le

di

th

ve

fc

no

tu

Te

ne

le

or

la

u

01

th

ar

Ik

in

u

1

This future is fometimes very irregular, being made up of a great many small sutures, which surround so many little bones that are generally larger and more conspicuous on the external surface of the skull than internally. These bones are generally called triquetra or Wormiana: but some other name ought to be given them, for they are not always of a triangular figure; and older anatomists (m) than Olaus Wormius (n) have described them.—The specific virtue which these bones were once thought to have in the cure of the epilepfy (o), is not now afcribed to them; and anatomists generally agree, that their formation is owing to a greater number of points than ordinary of offification in the skull, or to the ordinary bones of the cranium not extending their offification far enough or foon enough; in which case, the unossified interstice between such bones begins a separate offification in one or more points; from which the offification is extended to form as many distinct bones as there were points that are indented into the large ordinary bones, and into each other. Probably those children who have a large opening in this place at their birth, will have the largest offa triquetra.——To confirm this account of the formation of these little bones, we may remark, that such bones are fometimes feen in other futures, as well as in the lambdoidal (p); and they are fometimes in one table of the fkull, and not in the other (q).

The fagittat future (r) is placed longitudinally in the middle

(111) Eustach. Offium examen — Bauhin. Theat. Anat. lib. 3. cap. 5. — Paaw in Hippocrat. de vulner. cap. p. 56.

(n) Mufæum, lib. 3. c. 26.

(0) Bauhin, et Paaw ibid. Bartholin. Anat. reform. lib. 4.

cap. 5 .- Hildan Epiftol. 65.

(p) See examples in Vefal. lib. 1. cap. 6. fig. 4.—Paaw in Hippocrat. de cap. vuln.—Bartholin. Hist. Anat. cent. 1. hist. 51.—Rusch. Mus. Anat.—Sue Trad. d'osteolog. p. 47.

(9) Hunald. in Mem. de l'acad. des sciences, 1730.

(r) 'Paßsousne, osidaia, inizioyvora, Inftar virgæ, nervalis, inftar teli, inftar veru, fecundum capitis longitudinem prorepens, conjungens, columnalis, recta, acualis.

t I.

de

fo

ore

an

ra

en

e;

ve

les

P-

e-

ter

he

X-

in

es

8;

ny

n-

-1

in

e-

on

es

he

of

ne

le

3.

4.

in.

ar

11.

middle of the upper part of the skull, and commonly terminates at the middle of the coronal and of the lambdoidal sutures; between which it is said to be placed, as an arrow is between the string and bow.—However, this suture is frequently continued through the middle of the os frontis, down to the root of the nose; which, some (s) say, oftener happens in women than men; but others (t) allege, that is to be met with more frequently in male skulls than in semale: Among the skulls which I have seen thus divided, the semale are the most numerous.—Several (u) have delineated and described the sagittal suture, sometimes dividing the occipital bone as far down as the great hole through which the medulla spinalis passes. This I never saw.

In some old skulls that are in my possession, there is scarce a vestige of any of the three sutures which I have now described. In other heads, one or two of the sutures only disappear; but I never could discover any reason for thinking them disposed in such different manners in skulls of different shapes, as some ancients allege they are (x).

The fquamous agglutinations, or false sutures (y), are one on each side, a little above the ear, of a semicircular sigure, formed by the overlopping (like one scale upon another) of the upper part of the temporal bones on the lower part of the parietal, where, in both bones, there are a great many small risings and surrows, which are indented into each other; though these inequalities do not appear till the bones are separated. In some skulls, indeed, the indentations here are as conspicuous

de offib. cap. 8.

E 3

<sup>(1)</sup> Vefal. lib. 1. cap. 6. et in epitome.

in Celf. de re medic. cap. 1.—Laurent. Hist. Anat. lib. 2. cap. 161

<sup>(</sup>x) Hippocrat. de vulner. capitis, § 1. — Galen de offib. et de usu part. lib. 9. cap. 17.

<sup>(</sup>у) лежьбоный, продходимата, кротата, Temporales, corticales, mendofæ; harmonialis, commissuræ in unguem.

C

th

fp

fp

m

ft

0

is th

b

P

n

d

P

0

b

externally as in other futures (z); and what is commonly called the posterior part of this squamous suture, always has the evident serrated form; and therefore is reckoned by some (a) a distinct suture, under the name of additamentum posterius suturæ squamosæ.—I have seen two squamous sutures on the same temple, with a

femicircular piece of bone between them (b).

We ought here to remark, that the true squamous fort of future is not confined to the conjunction of the temporal and parietal bones, but is made use of to join all the edges of the bones on which each temporal muscle is placed (c): For the two parts of the sphenoidal future which are continued from the anterior end of the common fquamous future just now described, of which one runs perpendicularly downwards, and the other horizontally forwards, and also the lower part of the coronal future already taken notice, may all be justly faid to pertain to the fquamous future.—The manner how I imagine this fort of future is formed at these places is, that, by the action of the strong temporal muscles on one side, and by the pressure of the brain on the other, the bones are made fo thin, that they have not large enough furfaces opposed to each other to stop the extension of their fibres in length, and thus to cause the common ferrated appearance of futures explained in p. 47. but the narrow edge of the one bone flides over The fquamous form is also more convethe other. nient here; because such thin edges of bones, when accurately applied one to another, have fcarce any rough furface, to obstruct or hurt the muscle in its contraction; which is still further provided for, by the manner of laying these edges on each other; for, in viewing their

(a) Albin. de offib. § 54.

(b) Sue Trad. d'ofteolog. p. 48.

<sup>(</sup>z) Columb. de re anat. lib. 1. cap. 4.—Dionis, Anat. 3. de-monstrat. des os.

<sup>(</sup>c) Vefal. Anat. lib. 1. cap. 6.—Winflow, Mem. de l'acad. deq sciences, 1720.

ot

their outfide, we fee the temporal bones covering the fphenoidal and parietal, and this last supporting the fphenoidal, while both mount on the frontal: From which disposition it is evident, that, while the temporal muscle is contracting, which is the only time it presses ftrongly in its motion on the bones, its fibres flide eafily over the external edges. Another advantage still in this is, that all this bony part is made stronger by the bones thus supporting each other.

The bones of the skull are joined to those of the face, by schyndeless and sutures.—The schyndeless is in the partition of the nofe. The futures faid to be common to the cranium and face are five, viz. the ethmoidal, fphenoidal, transverse, and two zygomatic. Parts, however, of these sutures are at the junction of

only the bones of the skull.

The ethmoidal and sphenoidal sutures surround the bones of these names; and in some places help to make up other futures, particularly the fquamous and tranfverse; and in other parts there is but one future common

to these two bones.

art I,

om-

ure,

re is

ame

nave

th a

lous

the

join

oral

noiend

, of

the

t of

uft-

anele

oral

on ave

top

ule lin

ver

ve-

ac-

gh ac-

ier

ng eir

de-

deq

The transverse suture is extended quite cross the face, from the external canthus of one orbit to the same place of the other, by finking from the canthus down the outfide of the orbit to its bottom; then mounting upon its infide, it is continued by the root of the nofe down the internal part of the other orbit, and rifes up again on its outfide to the other canthus. It may be here remarked, that there are some interruptions of this future in the course I have described; for the bones are not contiguous every where, but are separated, to leave holes and apertures, to be mentioned hereafter.

The zygomatic futures are one on each fide, being fhort and flanting from above obliquely downwards and backwards, to join a process of the cheek-bone to one of the temporal bones, which advance towards the tace; fo that the two processes thus united, form a fort

E 4

C

ir

ir

ti

16

ti

t

O

a

P

C

of bridge or jugum, under which the temporal muscle passes; on which account the processes, and suture join-

ing them, have been called zygomatic.

It must be observed, that the indentations of the sutures do not appear on the infide of the cranium by much so strong as on the outside; but the bones seem almost joined in a straight line: Nay, in some skulls, the internal furface is found entire, while the futures are manifest without; which may possibly be owing to the less extent of the concave than of the convex furface of the cranium, whereby the fibres of the internal fide would be stretched farther out at the edges of the bones than the exterior ones, if they were not refisted. The refistances are; the fibres of the opposite bone, the parts within the skull, and the diploë: of which the last being the weakest, the most advanced fibres or serræ run into it, and leave the contiguous edges equal, and more ready to unite; whereas the ferræ of the external table have space enough for their admission between the fibres of the opposite bone, and therefore remain of the indented form, and are less liable to the concretion whereby the futures are obliterated (d),—By this mechanism, there is no risk of the sharp points of the bones growing inwards, fince the external ferræ of each of the conjoined bones rest upon the internal fmooth-edged table of the other; and external forces applied to these parts are strongly resisted, because the futures cannot yield, unless the serrated edges of the one bone, and the plain internal plate of the other, are broken (e).

The advantages of the futures of the cranium are these: 1. That this capsula is more easily formed and extended into a spherical sigure, than if it had been one continued bone. 2. That the bones which are at some distance from each other at birth, might then yield, and allow to the head a change of shape, ac-

commodated

<sup>(</sup>d) Hunald. Memoires de l'acad. des sciences, 1730. (e) Winslaw, Memoires de l'acad. des sciences, 1720.

rt I.

fcle

oin-

fu-

by

em

ills,

to ur-

nal

ed.

the last

rræ

er-

ain

re-

By

of

of

nal

ces

he

he

ire

nd

en

at

en

ed

commodated to the passage it is engaged in. Whence, in hard labour of child-bed, the bones of the cranium. instead of being only brought into contact, are sometimes made to mount one upon the other. 3. It is alleged, that, through the futures, there is a transpiration of steams from the brain, which was the old doctrine; or fome communication of the veffels without. and of those within the skull, larger here than in any other part of the cranium, according to fome moderns; and therefore cucuphæ, fomentations, cataplasms, cephalic plasters, blisters, are applied, and iffues are eroded or cut in the head, at those places where the sutures are longest in forming, and where the connection of the bones is afterwards loofest, for the cure of a phrenitis, mania, inveterate headach, epilepfy, apoplexy, and other diseases of the head. The favourers of the doctrine of transpiration, or communication of veffels at the futures, endeavour to support it by observations of persons subject to headachs which caused death, from the futures being too closely united (f). 4. That the dura mater may be more firmly fuspended by its processes, which infinuate themselves into this conjunction of the bones: for doing this equally, and where the greatest necessity of adhesion is, the sutures are disposed at nearly equal distances; and the large refervoirs of blood, the finuses, are under or near them. 5. That fractures might be prevented from reaching fo far as they would in a continued bony fubstance. 6. That the connection at the futures being capable of yielding, the bones might be allowed to separate; which has given great relief to patients from the violent fymptoms which they had before this separation happened (g). And it feems reasonable to believe, that the opening of the futures was of great benefit to feveral others who were rather judged to have been

<sup>(</sup>f) Columb. de re anat. lib. 1. cap. 5. — Verduc. Nouvelle Offeologie, chap. 14—Dionis, Anat. 3. demonstr des os. (g) Ephemerid, Germanic, dec. 1. ann. 4. et 5. observ. 33.

C

b

hurt by it (b): for we must think, that the consequences of such a force acting upon the brain, as was capable of thrusting the bones as under, must have been fatal, unless it had been thus yielded to.

Having gone through the general structure of the cranium, I now proceed to examine each bone of which that brain-case consists, in the order in which I first named them.

### OS FRONTIS.

THE os frontis (i) has its name from its being the only bone of that part of the face we call the forehead, though it reaches a good deal farther. It has fome refemblance in shape to the shell of the concha bivalvis, commonly called the cockle; for the greatest part of it is convex externally, and concave internally, with a serrated circular edge; while the smaller part has processes and depressions, which make it of an irregular figure.

The external furface of the os frontis is smooth at its upper convex part; but several processes and cavities are observable below: for at each angle of each orbit, the bone juts out to form four processes, two internal, and as many external; which, from this situation, may well enough be named angular. Between the internal and external angular processes on each side, an arched ridge is extended, on which the eye-brows are placed.

—Very little above the internal end of each of these superciliary ridges a protuberance may be remarked, in most skulls, where there are large cavities, called sinuses, within the bone; of which hereafter.—Betwixt the internal angular processes a small process rises, which forms some

(i) Μετώπη, Βρεγμα, Coronale, inverecundum, puppis, sensus communis, sincipitis.

<sup>(</sup>b) Ephemerid. Germ. dec. 2. ann. 9. obs. 230. Ibid. cent. 10. obs. 31.—Vander Linden Medicin.-phys. cap. 8. art. 4. § 16.—Hildan. Observ. cent. 1. obs. 1. cent. 2. obs. 7.—Bauhin. Theat. Anat. lib. 3. cap. 6.—Pechlin, Observ. lib. 2. obs. 39.

t I.

en-

ca-

en

he

ch

rst

he

d,

e-

is,

it

er-

es

its

es it,

ıl,

ay

al

ed

d.

ſe

in

5,

n-

18

e

0.

t.

na

In each of the orbitar processes, behind the middle of the superciliary ridges, a considerable sinuosity is observed, where the glandula innominata Galeni or lachrymalis is lodged.—Behind each internal angular process, a small pit may be remarked, where the cartilaginous pully of the musculus obliquus major of the eye is fixed.—Betwixt the two orbitar processes, there is a large discontinuation of the bone, into which the cribrisorm part of the os ethmoides is incased.—The frontal bone frequently has little caverns formed in it here where it is joined to the ethmoid bone.—Behind each external angular process, the surface of the frontal bone is considerably depressed where part of the temporal muscle is placed.

The foramina, or holes, observable on the external surface of the frontal bone, are three in each side.—
One in each superciliary ridge, a little removed from its middle towards the nose; through which a twig of the ophthalmic branch of the fifth pair of nerves passes out of the orbit, with a small artery from the internal carotid, to be distributed to the teguments and muscles of the forehead.—These vessels, in some skulls, make surrows in the os frontis, especially in the bones of children, as has also been observed of another considerable vessel of this bone near its middle (k); and

derable vessel of this bone near its middle (k); and therefore we ought to beware of transverse incisions on either side of the os frontis, which might either open

<sup>(</sup>k) Ruysch. Mus. Anat. theca D. reposit. 4. No. 3.

Ch

Me

tha

for

nu

ey

or

on

ho

nu ra

ne

po

no

it

X th

0

te

these vessels, or hurt the nerves, while they are yet in part within the bone: for when veffels are thus wounded, it is difficult to stop the hæmorrhagy, because the adhesion of a part of the artery to the bone hinders its contraction, and consequently styptics can have little effect; the fides of the furrow keep off compressing fubstances from the artery; and we would wish to shun cauteries or escharotics, because they make the bone carious; and nerves, when thus hurt, fometimes produce violent symptoms.—But to return to the superciliary foramina, we must remark, that often, instead of a hole, a notch only is to be feen: nay, in some skulls, scarce a vestige even of this is left; in others, both hole and notch are observable, when the nerve and artery run separately. Frequently a hole is found on one fide, and a notch on the other; at other times we fee two holes, or there is a common hole without, and two diffinct entries internally. The reason of this variety of a hole, notch, depression, or smoothness, in the superciliary ridge, is the different length and tension of the nerves and veffels; the shorter they are, the more they are funk into the bone as it grows.—Near the middle of the infide of each orbit, hard by or in the transverse suture, there is a small hole for the passage of the nafal twig of the first branch of the fifth pair of nerves, and of a branch of the opthalmic artery. This hole is fometimes entirely formed in the os frontis; in other skulls, the sides of it are composed of this last bone and of the os planum. It is commonly known by the name of orbitarium internum, though anterius should be added, because of the next, which is commonly omitted.—This, which may be called orbitarium internum posterius, is such another as the former; only fmaller, and about an inch deeper in the orbit: Through it a small branch of the ocular artery passes to the nose. -Besides these six, there are a great number of small holes observable on the outer surface of this bone, particularly in the two protuberances above the eye-brows. Molt

I.

in

d-

1e

ts le

g

m

e

• 6

r-

le

s,

e

S

1,

S

n

n

C

e

e

0

f

Most of these penetrate no further than the sinuses, or than the diploë if the sinuses are wanting; though sometimes I have seen this bone so personated by a vast number of these small holes, that, placed between the eye and a clear light, it appeared like a sieve.—In the orbit of the generality of skeletons, we may observe one, two, or more holes, which allow a passage to a hog's bristle through the skull. The place, size, and number of these, are however uncertain: they generally serve for the transmission of small arteries or nerves.

The internal furface of the os frontis is concave, except at the orbitar processes, which are convex, to support the anterior lobes of the brain. This furface is not fo fmooth as the external; for the larger branches of the arteries of the dura mater make some furrows in its fides and back parts. The finuofities from the luxuriant rifings of the brain, mentioned when describing the general structure of the cranium, are often very observable on its upper part; and its lower and foreparts are marked with the contortions of the anterior lobes of the brain.—Through the middle of this internal furface, where always in children, and fometimes in old people, the bone is divided, either a ridge stands out, to which the upper edge of the falx is fastened, or a furrow runs, in which the upper fide of the fuperior longitudinal finus is lodged; on both thefe accounts, chirurgical authors justly forbid the application of the trepan here.—The reason of this difference in skulls is alleged by some authors to be this, That in thin skulls the ridge strengthens the bones, and in thick ones there is no occasion for it. To this way of accounting for this phenomenon, it may justly be objected, that, generally, very thick skulls have a large spine here, and frequently thin ones have only a furrow. Perhaps this variety may be owing to the different times of complete offification of those parts in different subjects: for if the two fides of this bone meet before they arrive

C

vi

T

in

in

ha

in

he

th

th

ot fe

th

ſe

th

is

ra

h

la

u

th

th

tu

m

fi

tl

fi

n

n

n

ir

ti

n

0

arrive at their utmost extent of growth, they unite very firmly, and all their fibres endeavour to stretch themselves out where the least resistance is, that is, between the hemispheres of the brain. To support this reasoning, we may remark, that those adults whose frontal bone is divided by the sagittal suture, never have a

ridge in this place.

Immediately at the root of this ridge or furrow, there is a small hole, which sometimes pierces through the first table, and, in other sculls, opens into the superior sinus of the ethmoid bone within the nose. In it a little process of the falx is lodged, and a small artery, and sometimes a vein, runs (1); and the superior longitudinal sinus begins here.—This hole, however, is often not entirely proper to the os frontis: for, in several skulls, the lower part is formed in the upper part of the base of the crista galli, which is a process of the ethmoid bone (m).

The os frontis is composed of two tables, and an intermediate diploë, as the other bones of the cranium are, and in a middle degree of thickness between the os occipitis and the parietal bones; and is pretty equally dense all through, except at the orbitar processes, where, by the action of the eye on one side, and pressure of the lobes of the brain on the other, it is made extremely thin and diaphanous, and the meditullium is entirely obliterated. Since in this place there is so weak a defence for the brain, the reason appears why fencers

esteem a push in the eye mortal (n).

The diploë is also exhausted in that part above the eye-brows, where the two tables of the bone separate, by the external being protruded outwards, to form two large cavities, called finus frontales.—These are divided

(1) Morgagn. Adversar. 6. animad. 31.

<sup>(</sup>n) Ingrass. Comment. in Galen. de ossib. cap. 1. comment. 8.
(n) Ruysch. Observ. Anat.-chir. observ. 54—Diemerbroeck.
Anat. lib. 3. cap. 10.—Bonet. Sepulch. Anat. lib. 4. § 3. observ.
17.

t.

ry

n-

en

1-

al

a

re

e

or

a

Y s

1-

is

-

rt

e

1-

n

e

y

f

y

S

e

vided by a middle perpendicular bony partition.-Their capacities in the fame subject are seldom equal; in some the right, in others the left is largest.——And in different bones their fize is as inconstant: nay, I have examined fome where they were entirely wanting; which oftener happens in fuch as have a flat forehead, and whose fagittal future is continued down to the nose, than in others (o).——In some skulls, besides the large perpendicular *septum*, there are several bony pillars, or short partitions, found in each sinus: in others these are wanting.——For the most part the septum is entire; at other times it is discontinued, and the two finuses communicate.—When the finuses are feen in fuch skulls as have the frontal bone divided by the fagittal future, the partition dividing these cavities is evidently composed of two plates, which easily separate.—Each finus commonly opens by a roundish small hole, at the inner and lower part of the internal angular processes, into a finus formed in the nose, at the upper and back part of the os unguis; near to which there are also some other small sinuses of this bone (p), the greater part of which open separately near the septum narium, and often they terminate in the same common canal with the large ones.

In a natural and found state, these cavities are of confiderable advantage; for the organ of smelling being thus enlarged, the effluvia of odorous bodies more difficultly escape it; and their impressions being more numerous, are therefore stronger, and affect the organ more. That odorous particles may be applied to the membrane of the sinuses, is evident from the pain selt in this part of the fore-head, when the essuring of volatile spirits, or of strong aromatics, are drawn up into the nose by a quick inspiration.—These, and the other cavities which open into the nose, increase the sound of our voice, and render it more melodious, by serving as

(0) Fallop. Exposit. de offib. cap. 13.

<sup>(</sup>p) Cowper in Drake's Anthropolog. book 3. chap. 10.

bouring under a coryza, or stoppage of the nose, from any other cause, when they are by the vulgar, though falsely, said to speak through their nose, have such a disagreeable harsh voice.—The liquor separated in the membrane of these sinuses drills down upon the mem-

brane of the nose to keep it moift.

From the description of these sinuses, it is evident, how useless, nay, how pernicious, it must be to apply a trepan on this part of the skull: for this instrument, instead of piercing into the cavity of the cranium, would reach no further than the finuses; or, if the inner table was perforated, any extravafated blood that happened to be within the skull, would not be discharged outwardly, but would fall into the finuses, there to stagnate, corrupt, and stimulate the sensible membranes; from which also there would be such a constant flow of glairy mucus, as would retard, if not hinder a cure, and would make the fore degenerate into an incurable fif-Besides, as it would be almost impossible in this case to prevent the air, passing through the nose, from having constant access to the dura mater or brain; such a corruption would be brought on these parts as would be attended with great danger. Further, in respiration, the air rushing violently into these cavities of the os frontis, and passing through the external orifice, whenever it was not well covered and defended, would not only prevent the clofing up of the external orifice, but might otherwise bring on bad consequences (q)—The membrane lining these sinuses is so sensible, that inflammations of it must create violent torture (r); and worms, or other infects crawling there, must give great uneafiness (s)

The

CI

to

th

ro

18

A

ar

to

or

lit

ar

tu

tie

lo

ea

CI

al

tr

n

la

0

h

11

tl

t

t

1

1

<sup>(</sup>q) Paaw de offibus, pars 1. cap. 7.—Palfyne Anatom.-chir. traité 4. chap. 15. Nouvelle Ofteologie, partie 2. chap. 3.

<sup>(</sup>r) Fernel. Patholog. lib. 5. cap. 7.—-Saltzman Decur. observ.

<sup>(</sup>s) Fernel. Patholog. lib. 5. cap. 7.—Bartholin. Epistol. Medic. cent. 2. epist. 74:—Hist. de l'acad. des sciences, 1708 & 1733.

I.

la-

m

gh

if-

he

n-

ıt,

a

n-

ld

le

ed

t-

g-

5;

of

nd

f-

is

m

h

d

n,

SC

1-

ot

ıt

e

1-

d

ıt

C

r.

v.

The upper circular part of the os frontis is joined to the offa parietalia, from one temple to the other, by the coronal future. From the termination of the coronal future to the external angular processes, this bone is connected to the sphenoid by the sphenoidal suture. At the external canthi of the eyes, its angular processes are joined by the transverse suture to the osla malarum, to which it adheres one-third down the outfide of the orbits; whence to the bottom of these cavities, and a little up on their internal fides, these orbitar processes are connected to the fphenoidal bone by that fame future.—In some few skulls, however, a discontinuation of these two bones appears at the upper part of the long flit, near the bottom of the orbit. - On the infide of each orbit, the orbitar process is indented between the cribriform part of the ethinoid bone, and the os planum and unguis.—The transverse suture afterwards joins the frontal bone to the fuperior nafal processes of the offa maxillaria fuperiora, and to the nafal bones. And, laftly, its nafal process is connected to the nafal lamella of the ethmoid bone.

The frontal bone ferves to defend and support the anterior lobes of the brain. It forms a considerable part of the cavities that contain the globes of the eyes, helps to make up the septum narium, organ of smelling, &c. From the description of the several parts, the other uses of this bone are evident.

In a ripe child, the frontal bone is divided through the middle; the fuperciliary holes are not formed; often a small round piece of each orbitar process, behind the superciliary ridge, is not offissed; and there is no sinus to be seen within its substance.

## OSSA PARIETALIA.

EACH of the two Ossa Parietalia (t), or bones ferving as walls to the encephalon, is an irregular fquare;

<sup>(</sup>t) Kopupec, Paria fyncipitis, verticis, arcualia, nervalia, cogitationis, rationis, bregmatis, madefactionis.

C

u

tl

li

li

d

18

ir

fi

V

ua

r

fo

10

fi

tl

(

tl

0

t

18

a

p

p

0

i

t

The

fquare; its upper and fore fides being longer than the one behind or below. The inferior fide is a coneave arch; the middle part receiving the upper round part of the temporal bone.—The angle formed by this upper fide and the fore one, is fo extended, as to

have the appearance of a process.

The external furface of each os parietale is convex. Upon it, somewhat below the middle height of the bone, there is a transverse arched ridge, of a whiter colour generally than any other part of the bone; from which, in bones that have strong prints of muscles, we fee a great many converging furrows, like formany radii drawn from a circumference towards a centre. From this ridge of each bone the temporal muscle rises: and, by the prefiure of its fibres, occasions the furrows just now mentioned.—Below these we observe, near the semicircular edges, a great many rifings and depressions, which are joined to like inequalities on the infide of the temporal bone, to form the fquamous future. The temporal bone may therefore ferve here as a buttrefs, to prevent the lower fide of the parietal from starting outwards when its upper part is pressed or struck (u).

Near the upper fides of these bones, towards the hind part, is a small hole in each, through which a vein passes from the teguments of the head to the longitudinal sinus. Sometimes I have seen a branch of the temporal artery pass through this hole, to be distributed to the upper part of the salx, and to the dura mater at its sides, where it had frequent anastomoses with the branches of the arteries derived from the external carotids, which commonly have the name of the arteries of the dura mater, and with the branches of the internal carotids which serve the salx.—In several skulls, one of the ossa parietalia has not this hole: In others, there are two in one bone; and in some, not one in cither. Most frequently this hole is through both tables; at other times the external table only is perforated.—

(u) Hunauld. in Mem. de l'acad. des sciences 1730.

The knowledge of the course of these vessels may be of use to surgeons when they make any incision near this part of the head; lest, if the vessels are rashly cut near the hole, they shrink within the substance of the bone, and so cause an obstinate hamorrhagy, which neither

ligatures nor medicines can stop.

Ĩ.

n

1-

d

y

0

X.

10

er

m

ve

a-

m

d,

W

r-

ch

n-

0-

e-

It-

nd

af-

al

0-

to

at

he

a-

es

11-

is,

rs,

ci-

S;

he

On the inner concave furface of the parietal bones, we fee a great many deep furrows, disposed somewhat like the branches of frees: the furrows are largest and deepest at the lower edge of each os parietale, especially near its anterior angle, where fometimes a full canal is formed. They afterwards divide into small furrows, in their progress upwards.——In some skulls a large furrow begins at the hole near the upper edge, and divides into branches, which join with those which come upwards; showing the communications of the upper and lower veffels of the dura mater.—In these furrows we frequently fee passages into the diploë; and fometimes I have observed canals going off, which allowed a finall probe to pass some inches into the bony fubitance. Some (x) tell us, that they have observed these canals piercing the bone towards the occiput.— On the infide of the upper edge of the offa parietalia, there is a large finuofity, frequently larger in the bone of one fide than of the other, where the upper part of the falx is fastened, and the superior longitudinal sinus is lodged.—Generally part of the lateral finuses makes a depression near the angle, formed by the lower and posterior sides of these bones; and the pits made by the prominent parts of the brain are to be feen in no part of the skull more frequent, or more considerable, than in the internal furface of the parietal bones.

The offa parietalia are amongst the thinnest bones of the cranium; but enjoy the general structure of two tables and diploë the completest, and are the most

equal and fmooth.

These bones are joined at their fore-fide to the os

<sup>(</sup>x) Cowper. Anatom. explic. of tab. 90. fig 2.

y

d

C

ri

1

tl

fi

ti

2

W

n

19

ti

p

tl

P

0

16

13

u

h

la

frontis by the coronal future; at their long inferior angles, to the sphenoid bone, by part of of the suture of this name; at their lower edge, to the offa temporum, by the squamous suture, and its posterior additamentum; behind, to the os occipitis, or ossa triquetra, by the sambdoid suture; and above, to one another, by the sagittal suture.

They have no particular uses besides those mentioned in the description of their several parts, except what are included in the account of the general structure of

the cranium.

In a child born at the full time, none of the fides of this bone are completed; and there never is a hole in

the offified part of it near to the fagittal future.

The large unoffified ligamentous part of the cranium observable between the parietal bones and the middle of the divided os frontis of new-born children, called by the vulgar the open of the head, was imagined by the ancients to serve for the evacuation of the superfluous moisture of the brain: and therefore they named it bregma (y), or the fountain; sometimes adding the epithet pulsatilis, or beating, on account of the pulsation of the brain felt through this slexible ligamento-cartilaginous substance. Hence very frequently the parietal bones are called offa bregmatis.

The upper middle part of the head of a child, in a natural birth, being what presents itself first at the os uteri (z), an accoucheur may reach the bregma with his singer, when the os uteri is a little opened. If the bregma is stretched, and the pulsation of the brain is selt through it, the child is certainly alive: but if it is shrivelled and flaccid, without any observable pulsation in it, there is some reason to suspect the child to be very weak, or dead. Those who practise midwisery should therefore examine the state of the bregma accurately.

All

(y) Palpitans vertex, foliolum, folium, triangularis lacuna.

<sup>(</sup>z) Burton's Midwifery, § 51. Smellie's Midwifery, book 1. chap. 1. § 5.

T.

or

re

0-

a-

e-

-C

nat

of

of

in

m

lle

11-

Dy

u-

ed

he

a-

0-

a-

a

OS

th

he

18

18

on be

el y

u-

All

· le

All the bregma is generally offified before feven years of age. Several authors (a) fay, they have obferved it unoffified in adults; and physicians, who order the application of medicines at the meeting of the coronal and fagittal sutures, seem yet to think that a derivation of noxious humours from the encephalon is more easily procured at this part than any other of the skull, and that medicines have a greater effect here than elsewhere in the internal disorders of the head.

# OSSA TEMPORUM.

TEMPORUM (b), so named, say authors, from the hair's first becoming gray on the temples, and thus discovering peoples ages, are each of them equal and fmooth above, with a very thin femicircular edge; which, from the manner of its connection with the neighbouring bones, is distinguished by the name of os fquamofum ——Behind this, the upper part of the temporal bone is thicker, and more unequal; and is fometimes described as a distinct part, under the name of pars mamillaris (c).—Towards the base of the skull, the temporal bone appears very irregular and unequal; and this part, instead of being broad, and placed perpendicularly, as the others are, is contracted into an oblong very hard substance, extended horizontally forwards and inwards, which in its progress becomes smaller, and is commonly called os petrofum.

Three external processes of each temporal bone are generally described.—The first, placed at the lower and hind part of the bone, from its resemblance to a nipple, is called massoides, or mamillaris. It is not folid; but within is composed of cancelli, or small cells, which have a communication with the large cavity of the ear,

(a) Bartholin. Anat. reform. lib. 4 cap. 6. —— Diemerbroek, Anat. lib. 9. cap. 6. — Kerkring. Ofteogen cap. 2.

(b) Κοςτατων, κοςσων, κοςςων, λεπιδοειδη, πολυειδη, λιποειδη, Temporalia, lapidofa, mendofa, dura, arcualia, tympanum, armalia, faxea, paratalia.

(c) Albin. de offib. feet. 26.

CI

fo

th

m of

ce

th

th

in

ce

ha

Z

lo

m

18

be

W

in

b

in

th

C

v fi

f

0

fe

2

the drum; and therefore founds, being multiplied in this vaulted labyrinth, are increased before they are applied to the immediate organ of hearing. maltoid process the stenomastoideus muscle is inserted; and to its back part, where the furface is rough, the trachelomaftoideus and part of the splenius are fixed.— About an inch farther forward, the fecond process begins to rife out from the bone; and having its origin continued obliquely downwards and forwards for fome way, it becomes fmaller, and is stretched forwards to join with the os malæ; they together forming the bony jugum, under which the temporal muscle passes. Hence this process has been named zygomatic (d). edge has the strong aponeurosis of the temporal muscle fixed into it; and its lower part gives rife to a share of the maffeter.—The fore-part of the base of this process is an oblong tubercle, which, in a recent subject, is covered with a fmooth polished cartilage, continued from that which lines the cavity immediately behind this tubercle.—From the under craggy part of the os temporum, the third process stands out obliquely The shape of it is generally faid to resemble the ancient stylos scriptorius; and therefore it is called the styloid process (e). Some authors (f) however contend, that it ought to be named steloid, from its being more like to a pillar. Several muscles have their origin from this process, and borrow one half of their name from it; as stylo-glossus, stylo-byoideus, stylo-pharyngeus; to it a ligament of the os hyoideus is fometimes fixed; and another is extended from it to the infide of the angle of the lower jaw. This process is often, even in adults, not entirely offified, but is ligamentous at its root, and fometimes is composed of two or three diitinct pieces.—Round the root of it, especially at the

(d) Kayxpoc, Paris, ansæ ossium temporum, ossa arcualia, paria, jugalia, conjugalia.

(e) Γραιτοιίδη. βιλονοτίδη, πλημτρον, Os calaminum, fagittale, clavale, acuale, calcar capitis.

(f) Galen, De usu part. lib. 2. cap. 4.—Fallop. Observ. anatom.

I

in

re

he

d;

e-

in

ne to

ny

ce

er

of

ois

ed nd

he

le

ed

n-

ıg

ir

11-

es

of

en

ts i-

ne

e-

11-

e,

n.

fore-part, there is a remarkable rifing of the os petrofum, which some have esteemed a process; and, from the appearance it makes with the stylisorm, have named it vaginalis.—Others again have, under the name of auditory process, reckoned among the external processes that semicircula rridge, which, running between the root of the mastoid and zygomatic processes, forms the under part of the external meatus auditorius.

The finuofities or depressions on the external surface of each os temporum are these: ——A long fosfa at the inner and back part of the root of the mammary procels, where the polterior head of the digastric muscle has its origin.——Immediately before the root of the zygomatic process, a confiderable hollow is left for lodging the crotaphite muscle.—Between the zygomatic, auditory, and vaginal processes, a large cavity is formed; through the middle of which, from top to bottom, a fiffure is observable, into which part of the ligament that fecures the articulation of the lower jaw with this bone is fixed. The fore-part of the cavity being lined with the fame cartilage which covers the tubercle before it, receives the condyle of the jaw; and in the back-part a small share of the parotid gland, and a cellular fatty substance, are lodged.—At the inside of the root of the styloid apophyse, there is a thimble-like cavity, where the beginning of the internal jugular vein, or end of the lateral finus, is lodged.—And as the finules of the two fides are frequently of unequal fize, fo one of these cavities is as often larger than the other (g) --- Round the external meatus auditorius, several finushties are formed for receiving the cartilages and ligaments of the ear, and for their firm adhefion.

The holes that commonly appear on the outfide of each of these bones, and are proper to each of them, are five.—The first, situated between the zygomatic and mastoid processes, is the orifice of a large funnel-

(g) Hunauld. in Mem de l'acad. des sciences, 1730.

C

tl

a

f

n

t

t

b

2

t

I

t

t

like canal, which leads to the organ of hearing; therefore is called meatus auditorius externus (b).——The fecond gives passage to the portio dura of the seventh pair of nerves; and from its fituation between the mastoid and styloid processes, is called foramen stylomastoideum (i).—Some way before, and to the inside of the styloid process, is the third hole; the canal from which runs first upwards, then forwards, and receives into it the internal carotid artery, and the beginning of the intercostal nerve; where this canal is about to make the turn forwards, one, or fometimes two, very fmall holes go off towards the cavity of the ear, called tympanum: Through these Valsalva (k) affirms the proper artery or arteries of that cavity are fent.—On the anterior edge of this bone, near the former, a fourth hole is observable, being the orifice of a canal which runs outwards and backwards, in a horizontal direction, till it terminates in the tympanum. This, in the recent subject, is continued forward and inward, from the parts which I mentioned just now as its orifice in the skeleton, to the side of the nostrils; being partly cartilaginous, and partly ligamentous. The whole canal is named Iter a palato ad aurem, or Eustachian tube. -On the external fide of the bony part of this canal, and a-top of the chink, in the cavity that receives the condyle of the lower jaw, is the course of the little nerve faid commonly to be reflected from the lingual branch of the fifth pair, till it enters the tympanum, to run across this cavity, and to have the name of chorda tympani.—The fifth hole is very uncertain, appearing fometimes behind the mastoid process; sometimes it is common to the temporal and occipital bones; and in feveral skulls there is no such hole. The use of it, when found, is for the transmission of a vein from the external teguments to the lateral finus: But, in some fub-

(i) Aquæductus Fallopii.

<sup>(</sup>b) Hogos THE axons, OTH TWY WTWY, feneftra aurium.

<sup>(</sup>k) De aure humana, cap. 2. § 22. et tab. 7. fig. 1.

t I.

ere-

fe-

nth

na-

2:10-

of of

om

ves

gof

to

ery

led

ro-

the

rth

ich

ecthe

om

in

rtly

ca-

ibe.

al,

the

ttle

ual

to

rda

ar-

nes

ind

it,

he

ıb-

fubjects, a branch of the occipital artery passes thro's this hole, to serve the back-part of the dura mater; in others, I have seen two or three such holes: but they are oftener wanting than found. And we may, once for all, in general remark, that the largeness, number, situation, and existence of all such holes as for the most part allow only a passage for veins from without to the internal receptacles, are very uncertain.

The internal furface of the offa temporum is unequal; the upper circular edge of the fquamous part having numerous small ridges and furrows for its conjunction with the parietal bones; and the rest of it is irregularly marked with the convolutions of the middle part of the brain, and with furrows made by the branches of the arteries of the dura mater.

From the under part of this internal furface, a larger transverse hard craggy protuberance runs horizontally inwards and forwards, with a sharp edge above, and two slat sides, one facing obliquely forwards and outwards, and the other as much backwards and inwards. To the ridge between these two sides, the large lateral process of the dura mater is fixed.

Sometimes a small bone, akin to the sesamoid, is found between the small end of this petrous process and the sphenoid bone (1).

Towards the back-part of the infide of the os temporum, a large deep fossa is conspicuous, where the lateral sinus lies; and frequently on the top of the petrous ridge, a surrow may be observed, where a small sinus is situated-

The internal proper foramina of each of these bones are, first, the internal meatus auditorius in the posterior plain side of the petrous process. This hole soon divides into two; one of which is the beginning of the aqueduct of Fallopius; the other ends in several very small

<sup>(1)</sup> Riolan. Comment. de offib. cap. 32.—Winflow, Exposi-

fmall canals (m) that allow a passage to the branches of the portio mollis of the seventh pair of nerves, into the vestibule and cochlea. Through it also an artery is fent, to be distributed to the organ of hearing.—The fecond hole, which is on the anterior plain fide of the craggy process, gives passage to a reflected branch of the second branch of the fifth pair of nerves, which joins the portio dura of the auditory nerve, while it is in the aqueduct (n), small branches of blood-vessels accompanying the nerves, or paffing through fmaller holes near this one.—The passage of the cutaneous vein into the lateral finus, or of a branch of the occipital artery, is feen about the middle of the large fossa for that finus; and the orifice of the canal of the carotid artery is evident at the under part of the point of the petrous process.

Besides these proper holes of the temporal bones which appear on their external and internal surfaces, there are two others in each side that are common to this bone, and to the occipital and sphenoidal bones; which shall be mentioned afterwards in the description

of these bones.

The upper round part of the squamous bones is thin, but equal; while the low petrous part is thick and strong, but irregular and unequal, having the distinction of tables and diploë confounded, with several cavities, processes, and bones within its substance, which are parts of the organ of hearing. That a clear idea may be had of this beautiful, but intricate organ, anatomists generally choose to demonstrate all its parts together. I think the method good; and therefore, since it would be improper to insert a complete treatise on the ear here, shall omit the description of the parts contained within the os petrosum of the skeleton.

The temporal bones are joined above to the parietal bones by the squamous sutures and their po-

sterior

Ci

fte

th

ZY

by

th

110

m

CC

fre

ve

th

of

th

ye

m

WA

ty

ci

lo

n

C

t

a

V

ì

(n) Ibid. cap. 3. § 10.

<sup>(</sup>m) Valsalva de aure humana, cap. 3. § 11.

t I.

of

is he

he

of

in

ler

ein

ar-

at

ry

us

les

es,

to

S;

on

n,

nd

a-

ch

ea

la

0-

ce

on

rts

a-

or

flerior additamenta: Before, to the sphenoid bone, by the suture of that name; to the cheek-bones, by the zygomatic sutures: Behind, to the occipital bone, by the lambdoid suture and its additaments; and they are articulated with the lower jaw in the manner which shall be described when this bone is examined.

The purposes which these two bones serve, are easily collected, from the general use of the cranium, and from what has been said in the description of their serveral parts.

In an infant, a small fissure is to be observed between the thin upper part and the lower craggy part of each of these bones; which points out the recent union of these parts.—Neither mastoid nor styloid processes are yet to be seen.—Instead of a bony funnel-like external meatus auditorius, there is only a smooth bony ring, within which the membrane of the drum is fastened—At the entry of the Eustachian tube, the side of the tympanum is not completed.—A little more outward than the internal auditory canal, there is a deep pit, over the upper part of whose orisice the interior semi-circular canal of the ear is stretched; and some way below this, the posterior semicircular canal also appears manifestly.

#### OS OCCIPITIS.

Os occipitis (0), so called from its situation, is convex on the outside, and concave internally. Its sigure is an irregular square, or rather rhomboid; of which the angle above is generally a little rounded; the two lateral angles are more finished, but obtuse; and the lower one is stretched forward in form of a wedge, and thence is called by some the cuneiform process.—If one would, however, be very nice in observing the several turns which the edges of the os occipitis

<sup>(0)</sup> Ivov, Basilare, proræ, memoriæ, pixidis, sibrosum, nervosum,

Ch

ing

fau

wh

ple

an

fu

ap

is

wl

an

fic

th

In

be

ex

bo

pi

ar

h

tl

ai

W

m

h

fi

r

tl

V

n

ti

i

P

F

f

ing

pitis make, five or feven fides, and as many angles of

this bone, might be described.

The external furface is convex, except at the cuneiform apophyse, where it is flatted. At the base of this triangular process, on each fide of the great hole, but more advauced forwards than the middle of it, the large oblong protuberances, named the condyles, appear, to serve for the articulation of this bone with the first vertebra of the neck. The smooth surface of each of these condyloid processes is longest from behind forwards, where, by their oblique fituation, they come much nearer to each other than they are at their back-part. Their inner fides are lower than the external, by which they are prevented from fliding to either fide out of the cavities of the first vertebra (p). In fome subjects, each of these plain smooth surfaces seems to be divided by a fmall rifing in its middle; and the lower edge of each condyle, next the great foramen, is discontinued about the middle, by an intervening notch: Whence fome (q) allege, that each of these apophyses is made up of two protuberances. Round their root a finall depression and spongy roughness is observable, where the ligaments for furrounding and fecuring their articulations adhere.—Though the motion of the head is performed on the condyles, yet the centre of gravity of that globe does not fall between them, but is a good way further forward; from which mechanism it is evident, that the muscles which pull the head back must be in a constant state of contraction; which is stronger than the natural contraction of the proper flexors, else the head would always fall forwards, as it does when a man is afleep, or labours under a palfy, as well as in infants, where the weight of the head far exceeds the proportional strength of these muscles. This seeming disadvantageous situation of the condyles is, however, of good use to us, by allow-

<sup>(</sup>p) Galen. de usu part. lib. 12. cap. 7. (q) Diemerbroek, Anat. lib. 9. cap. 6.

ing fufficient space for the cavities of the mouth and fauces, and for lodging a fufficient number of muscles, which commonly ferve for other uses; but may at pleasure be directed to act on the head, and then have an advantageous lever to act with, fo as to be able to fultain a confiderable weight appended, or other force

applied, to pull the head back.

t I,

s of

nei-

this

but

the

ap-

the

ot

be-

ney

eir

ex-

ei-

In

ms

the

en,

ng

ele

eir

er-

cu-

of

tre

m,

re-

the

n;

he

ds,

a

he

ese

of W-

ng

Somewhat more externally than the condyles, there is a fmall rifing and femilunated hollow in each fide, which make part of the holes common to the occipital and petrous bones.—Immediately behind this, on each fide, a scabrous ridge is extended from the middle of the condyle towards the root of the mastoid process. Into this ridge the musculus lateralis, commonly ascribed to Fallopius, is inferted.—About the middle of the external convex furface, a large arch runs cross the bone; from the upper lateral parts of which the occipital muscles have their rise; to its middle the trapezii are attached; and half-way between this and the great hole, a leffer arch is extended.—In the hollows between the middle of these arches the complexi are inserted and in the depressions more external and further forward than these, the splenii are inserted. Between the middle of the leffer arch and the great hole, the little hollow marks of the recti minores appear; and on each fide of these the sleshy insertions of the obliqui superiores and recti majores make depressions.—Through the middle of the two arches a small sharp spine is placed, which ferves as some fort of partition between the muscles of different sides, or rather is owing to the action of the mufcles depressing the bone on each side of it, while this part is free from their compression. These prints of the muscles on this bone are very strong and plain in some subjects, but are not so distinct in others. All round the great foramen, the edges are unequal, for the firmer adhesion of the strong circular ligament which goes thence to the first vertebra. One end of each lateral or moderator ligament of the head, is fixed

Ch

lim

Ro

pre

twe

of

int

the

the

wl

fw

fir

its

an

th

ne

W

ag

ni

ar

fic

th

er

th

th

e

th

tr

11

fi

to a rough furface at the fore-part of each condyle, and the perpendicular one is connected to a rough part of the edge of the great hole between the two condyles. Immediately before the condyles, two little depressions are made in the external furface of the cuneiform process, for the insertion of the recti anteriores minores muscles, which are unjustly ascribed to Cowper: And still further forward, near the sphenoid bone, are two other fuch depressions, for the reception of the recti anteriores majores. When we consider the fize of the prints of muscles on the occipital bone, before and behind its condyles, and at the fame time compare their distances from these centres of motion of the head, we must see how much stronger the muscles are which pull the head backwards, than those are which bend it forward; and how much greater force the former acquire by the long lever they act with, than the latter which are inferted so near the condyles. This great force in the extensor muscles is altogether necessary, that they might not only keep the head from falling forward in an erect posture, but that they might support it when we bow forward in the most necessary offices of social life, when the weight of the head comes to act at right angles on the vertebræ of the neck, and obtains a long lever to act with.

On the inner furface of the os occipitis we fee two ridges; one standing perpendicular, the other running horizontally across the first. The upper part of the perpendicular limb of the cross, to which the falx is fixed, is hollowed in the middle, or often on one side, for the reception of the superior longitudinal sinus; and the lower part of it has the small or third process of the dura mater fastened to it, and is sometimes hollowed by the occipital sinus. Each side of the horizontal limb is made hollow by the lateral sinuses inclosed in the transverse process of the dura mater; the fossa in the ripht side being generally a continuation of the one made by the longitudinal sinus in the perpendicular limb,

t I.

and

of

les.

ons

ro.

res

and

wo

an-

the

be-

eir

we

oull

or-

ire

ich

in

ey

in

en

ial

ht

ng

WO

ng

he

is le,

nd

he

ed

nb

he he

ne

ar

b,

simb, and therefore is larger than the left one (r)—Round the middle of the cross there are four large depressions, separated by its limbs; the two upper ones being formed by the back part of the brain, and the two lower ones by the cerebellum.—Farther forward than the last-mentioned depressions, is the lower part of the sossa for the lateral sinus on each side.—The inner surface of the cuneiform apophyse is made concave for the reception of the medulla oblongata, and of the basilar artery.—A surrow is made on each side, near the edges of this process, by a sinus of the dura mater, which empties itself into the lateral sinus (s).

The holes of this bone are commonly five proper, and two common to it and to the temporal bones——The first of the proper holes, called for amen magnum (t) from its fize, is immediately behind the wedge-like process; and allows a passage to the medulla oblongata, nervi accessorii, to the vertebral arteries, and sometimes to the vertebral veins.——At each fide of this great hole, near its fore-part, and immediately above the condyles, we always find a hole, fometimes two, which foon unite again into one that opens externally; through these the ninth pair of nerves go out of the skull.—The fourth and fifth holes pierce from behind the condyle of each fide into the fossæ of the lateral sinuses; they serve for the passage of the cervical veins to these finuses. en one of these holes is wanting, sometimes both, when the veins pals through the great foramen.—Besides these five, we frequently meet with other holes near the edges of this bone, for the transmission of veins; but their number and diameter are very uncertain. two common foramina are the large irregular holes, one m each fide, between the fides of the cuneiform procels and the edges of the petrous bones. In a recent subject, a strong membrane runs cross from one side to

<sup>(</sup>r) Morgagn. Advers. anat. 6. animad, 1,

<sup>(1)</sup> Albin. de offib. fect. 65. (1) Rachitidis, Medullæ spinalis.

C

be

V

fi

fc

de

of

fh

th

uj

th

lit

T

al

fr

CE

fo

T

a

ni

th

th

et

CO

as

th

re

bo

id

pr

fe.

its

to the other of each of these holes: in some heads I have seen this membrane offisied, or a bony partition dividing each hole; and in the greater number of adult skulls, there is a small sharp-pointed process stands out from the os petrosum, and a more obtuse rising in the occipital bone, between which the partition is stretched. Behind this partition, where the largest space is lest, the lateral sinus has its passage; and before it the eighth pair of nerves and accessorius make their exit out of the skull; and some authors say, an artery passes through this hole to be bestowed on the dura mater.

The occipital bone is among the thickest of the cranium, though unequally so; for it is stronger above, where it has no other defence than the common teguments, than it is below; where, being pressed by the lobes of the brain and cerebellum on one side, and by the action of the muscles on the other, it is so very thin as to be diaphanous in many skulls: but then these muscles ward off injuries, and the ridges and spines, which are frequent here, make it sufficiently strong to

refift ordinary forces.

The tables and diploë are tolerably distinct in this bone, except where it is so thin as to become diaphanous.

The occipital bone is joined above to the offa parietalia and triquetra, when present, by the lambdoid suture;—laterally to the temporal bones, by the additamenta of the lambdoid suture;—below to the sphenoid bone, by the end of its cuneiform process, in the same way that epiphyses and their bones are joined: For in children a ligamentous cartilage is interposed between the occipital and sphenoid bones, which gradually turns thinner as each of the bones advances, till their sibres at last run into each other; and about sixteen or eighteen years of age, the union of these two bones becomes so intimate, that a separation cannot be made without violence.—The os occipitis is joined by a double articulation to the first vertebra of the neck, each condyle being

being received into a superior oblique process of that vertebra. What motion is allowed here we shall confider afterwards, where the vertebræ are described.

The uses of this bone appear from the preceding de-

scription, and therefore need not be repeated.

An infant, born at the full time, has this bone divided, by unoffified cartilages, into four parts. - The first of these is larger than the other three, is of a triangular shape, and constitutes all the part of the bone above the great foramen. Generally fiffures appear in the upper part and fides of this triangular bone, when all the cartilage is separated by maceration; and sometimes little distinct bones are seen towards the edges of it — The fecond and third pieces of this bone are exactly alike, and fituated on each fide of the great foramen, from which very near the whole condyles are produced; and they are extended forwards almost to the fore-part of the hole for the ninth pair of nerves.— The fourth piece is the cuneiform process, which forms a small share of the great hole, and of those for the ninth pair of nerves and of the condyles; betwixt it and the fphenoid bone, a cartilage is interposed.

Of the eight bones which belong to the cranium, there are only two which are not yet described, viz. the ethmoid and sphenoid. These we already mentioned, in complaisance to the generality of writers on this subject, as bones common to the cranium and face, because they enter into the composition of both; but the same reason might equally be used for calling the frontal bone a common one too. I shall, however, pass any idle dispute about the propriety of ranging them, and proceed to examine the structure of the bones them-

felves.

t.

s I

on

alt

ut

he

ed.

he

th

he

gh

a-

re,

u-

he

by

ry

efe

es,

to

nis

la-

ie-

u-

ta-

oid

ne

in

en

ns

at en

fo

ut

ti-

yle

ng

### OS ETHMOIDES.

Os ETHMOIDES (u), or the fieve-like bone, has got its name from the great number of small holes with Vol. I.

G which

(u) Cribriforme, eneryouens, spongiforme, cristatum.

(

t

ti

1

0

fe

t

f

u

h

t

which that part of it first taken notice of is pierced. When this bone is entire, the figure of it is not easily described; but, by a detail of its several parts, some idea may be afforded of the whole: and therefore I shall distinguish it into the cribriform lamella with its process, the nasal lamella, cellulæ, and offa spongiosa.

The thin horizontal lamella is all (except its backpart) pierced obliquely by a great number of finall holes, through which the filaments of the olfactory In a recent subject, these holes are so nerves pals. closely lined by the dura mater, that they are much le's conspicuous than in the skeleton. From the middle of the internal fide of this plate, a thick process rifes upwards; and being highest at the fore-part, gradually becomes lower as it is extended backwards. fome refemblance which this process was imagined to have to a cock's comb, it has been called crista galli (x). The falx is connected to its ridge, and to the unperforated part of the cribriform plate. When the crista is broke, its base is sometimes found to be hollow, with its cavity opening into the nose (y). Immediately before the highest part of this process, is the blind hole of the frontal bone, which, as was formerly remarked, is often in a good measure formed by a notch in the fore-part of the root of the crista.

From the middle of the outer furface of the cribriform lamella, a thin folid plate is extended downwards
and forwards, having the fame common base with the
crista galli. Generally it is not exactly perpendicular,
but is inclined to one side or other, and therefore divides the cavity of the nose unequally. Its inclination
to one side, and slexure in the middle, is sometimes so
great, that it fills up a large share of one of the nostrils,
and has been mistaken for a polypus there. It is thin
at its rise, and rather still thinner in its middle; yet
afterwards, towards its lower edge, it becomes thicker,

that

<sup>(</sup>x) Verruca prædura, feptum offis fpongiofi. (y) Palfyn. Anat.-chir. tr. 4. chap. 15.

ı.

y

le

I

ts

<-

Ill

ry

fo

ch

le

es

ly

m

to

21-

he

he

ol-

ic-

he

rly

a

ri-

ds

he

ar,

di-

on

fo

ils,

nin

yer

er,

hat

are

that its conjunction with the bones and middle cartilage of the nose might be firmer.

At a little distance from each side of this external process, a cellular and spongy bony substance depends from the cribriform plate. The number and sigure of the cells in this irregular process of each side, are very uncertain, and not to be represented in words; only the cells open into each other, and into the cavity of the nose: The uppermost, which are below the aperture of the frontal sinuses, are formed like sunnels. The outer surface of these cells is smooth and plain, where this bone at sists in composing the orbit; at which place, on each side, it has got the name of os planum; on the upper edge of which a small notch or two may sometimes be observed, which go to the formation of the internal orbitar holes; as was remarked in the description of the frontal bone.

Below the cells of each fide, a thin plate is extended inwards; and then bending down, it becomes thick and of a fpongy texture. This fpongy part is triangular, with a straight upper edge placed horizontally, an anterior one slanting from above, downwards and forwards, and with a pendulous convex one below. The upper and lower edges terminate in a sharp point behind. The side of this pendulous spongy part next to the septum narium is convex, and its external side is concave. These two processes of the ethmoid bone have got the name of ossa spongiosa, or turbinata superiora, from their substance, sigure, and situation.

All the prominences, cavities, and meanders of this ethmoid bone, are covered with a continuation of the membrane of the nostrils in a recent subject. Its horizontal cribriform plate is lodged between the orbitar processes of the frontal bone, to which it is joined by the ethmoid suture, except at the back-part, where it is connected with the cuneiform bone by a suture common to both these bones, though it is generally esteemed part of the sphenoidal.—Where the ossa plana

G 2

0

p

f

fi

p

C

f

V

k

h

t

P

8

b

t

f

i

K

are contiguous to the frontal bone within the orbit, their conjunction is reckoned part of the transverse suture.—Farther forward than the offa plana, the cells are covered by the offa unguis: which are not only contiguous to these cells, but cannot be separated from them without breaking the bony fubstance; and therefore, in propriety, those bones ought to be demonstrated as part of the ethmoid bone.—Below the offa unguis and plana, these cells and offa spongiosa are overlopped by the maxillary bones. The cellular part of each palate bone is contiguous to each os planum and cells The lower edge of the nafal perpendicular plate is received into the furrow of the vomer.— Its posterior edge is joined to the fore-part of the processus azygos of the sphenoid bone. Its upper edge joins the nasal process of the frontal and nasal bones; and its anterior one is connected to the middle cartilage of the nofe.

From all which, the uses of this bone are evident, viz. to sustain the anterior lobes of the brain; to give passage to the olfactory nerves, and attachment to the falx; to enlarge the organ of smelling, by allowing the membrane of the nose a great extent; to straiten the passage of the air through the nose, by leaving only a narrow winding canal, on the sensible membranous sides of which the substances conveyed along with the air must strike; to form part of the orbit of the eyes and septum narium; while all its parts are so light as not to be in hazard of separating by their weight; and they are so thin, as to form a large surface, without occupying much space. This brittle substance, however, is sufficiently protected from external injuries by the firm bones which cover it.

If this bone is seized on by any corroding matter, we may easily conceive what destruction may ensue. Hence it is, that an ozæna is dissicult to cure; and that in violent scurvies, or in the lues venerea, the fabric of the nose, the eyes, and life itself, are in danger.

The

The fituation of the nafal plate may show us how dangerous a fracture of the bones of the nose may be, when made by a force applied to their middle fore-part, of a person in whom this nasal plate is perpendicular.

The ethmoid bone of ripe children is divided into two by a perpendicular cartilage, which, when offified, is the crista galli and nasal plate; but its other parts are offished and complete.

## OS SPHENOIDES.

Os sphenoides (z), or wedge-like bone, so called because of its situation in the middle of the bones of the cranium and face, is of such an irregular sigure, that I know not any thing to which it may be likened, unless, perhaps, it bear some faint resemblance to a bat

with its wings extended.

T.

,

l-

y

n

-

d

is

d

l.

1-

)-

e

e

t,

re

e

e

e

a

15

e

es

18

d c-

r,

e

r,

2.

d

1-

r.

e

When we view the external furface of the os fphenoides, two or three remarkable processes from each fide of it may be observed, which are all of them again fubdivided. The first pair is the two large lateral processes or wings; the upper part of each of which is called the temporal process, because they join with the temporal bones in forming the temples, and the feat for some share of the crotaphite muscles. That part of the wings which juts out towards the infide, fomewhat lower than the temporal apophyses, and is smooth and hollowed, where it makes up part of the orbit, is thence named orbitar processes. Behind the edge, feparating these two processes, there is often a small groove, made by a branch of the superior maxillary nerve, in its passage to the temporal muscle. The lowest and back part of each wing, which runs out fharp to meet the offa petrofa, has been styled the spinous process: from near the point of which a sharp-pointed process is frequently produced downwards, which some call styhform, that affords origin to the ptery-staphylinus exter-G

<sup>(</sup>z) Cuneiforme, \*\* nultiforme, paxillum, cribratum palati, colatorii, cavilla, basilare.

CI

VO

bu

fo

of

pa

fo

te

u

p

th

th

tl

te

t

nus muscle. From this styloid process a very small groove is extended along the edge of the bone to the hollow at the root of the internal plate of the following processes, which forms part of the Eustachian tube (a). The fecond pair of external processes of the cuneiform bone, is the two which stand out almost perpendicular to the base of the skull. Each of them has two plates, and a middle fossa facing backwards, and should, to carry on our comparison, be likened to the bat's legs; but are commonly faid to refemble the wings of that creature; and therefore are named pterygoid or aliform (b) processes. The external plates are broadest, and the internal are longest. From each fide of the external plates the pterygoid muscles take their rife. the root of each internal plate, a fmall hollow may be remarked, where the mufculus ptery-staphylinus internus or circumflexus palati rifes, and fome share of the cartilaginous end of the Eustachian tube rests; and, at the lower end of the fame plate, is a hook-like rifing or process, round which the tendon of the last-named muscle plays, as on a pulley. From the edge of the external plates some small sharp spikes stand out; but their number and bulk are uncertain,—To these another pair may be added, to wit, the little triangular thin process, which comes from each fide of the body of the fphenoid bone, where the pterygoid processes are rising from it, and are extended over the lower part of the aperture of the finus as far as to join the ethmoid bone, while their body hangs down into the nares (c). Befides these pairs of processes, there is a sharp ridge which stands out from the middle of its base: Because it wants a fellow, it may be called processus azygos. lower part of this process, where it is received into the

<sup>(</sup>a) Winflow, Exposition anatomique du corps humain, traité des os secs, § 233.

<sup>(</sup>b) Naviculares.
(c) Albin. Tab. Off. 5. fig. 2. 6. A. A. Bertin, Mem. de l'acad. des sciences, 1744.—Sue, planche viii. fig. 2, 3, 4, 5, 6.

I.

Ill

le

g

).

m

ar

s,

to

; at

112

d

r-

lt

e

r-

at

g

-

ir

r

n e

e

S

h

t

e

e

6

vomer, is thick, and often not quite perpendicular, but inclining more to one fide than the other. The fore-part of this process, where it joins the nasal plate of the os ethmoides, is thin and straight. These two parts have been described as two distinct processes by some.

The depressions, finuosities, and fosse, on the external furface of this sphenoid bone, may be reckoned up to a great number, viz. two on the temporal apophyles, where the crotaphite muscles lodge.—I'wo on the orbitar processes, to make way for the globes of the eyes.—Two between the temporal and fpinous processes, for receiving the temporal bones —— Two between the plates of the pterygoid processes, where the mufculi pterygoidei interni and ptery-staphylini interni are placed. Two between the pterygoid and orbitar processes, for forming the holes common to this and to the cheek and maxillary bones. the lower ends of the aliform processes, which the palate bones enter into.—Two at the roots of the temporal and pterygoid processes, where the largest share of the external pterygoid muscles have their rife. - Two at the fides of the processus azygos, for forming part of the nofe, &c.

What I described under the name of temporal and spinous processes on the outside of the skull, are likewise seen on its inside, where they are concave, for receiving part of the brain; and commonly three apophyses on the internal surface of the sphenoid bone are only mentioned.—Two rising broad from the fore-part of its body, become smaller as they are extended obliquely backwards—The third standing on a long transverse base, near the back-part of the body of this bone, rises nearly erect, and of an equal breadth, terminating often in a little knob on each side. The three are called clinoid, from some resemblance which they were thought to have to the supporters of a bed. Sometimes one or both the anterior clinoid processes are joined to the sides

G 4

th

in

C

tr

el

w

r

tl

i

t

of the posterior one, or the body of the bone itself. From the roots of the anterior clinoid processes the bone is extended on each side outwards and forwards, till it ends in a sharp point, which may have the name of the transverse spinous process.—Between, but a little farther back than the two anterior clinoid processes, we see a protuberance considerably smaller than the posterior clinoid process, but of its shape.—Another process from between the transverse processes, often forces itself forwards into the os ethmoides.

Within the skull, there are two finuofities in the internal part of each wing of the sphenoid bone, for receiving the middle part of the brain.—One between the transverse spinous processes, for lodging the part of the brain where the crura medullæ oblongatæ are.—Immediately before the third or middle clinoid process, a fingle pit generally may be remarked, from which a fossa goes out on each side to the holes through which the optic nerves pass. The pit is formed by the conjoined optic nerves; and in the fossa these nerves are lodged, as they run divided within the kull. that third protuberance and the posterior clinoid process, the larger pit for the glandula pituitaria may be remarked. This cavity, because of its resemblance to a Turkish saddle, is always described under the name of fella turcica, or ephippium.—On the fides of the posterior clinoid process a fossa may be remarked, that stretches upwards, then is continued forwards along the fides of the fella turcica, near to the clinoid anterior processes, where a pit on each side is made. These fossæ point out the course of the two internal carotid arteries, after they have entered the skull. Besides all these, several other fosse may be observed, leading to the feveral holes, and imprinted by the nerves and blood-veffels.

The boles on each fide of the os sphenoides are six proper, and three common.— The first is the round one immediately below the anterior clinoid processes, for

the

t I.

om

ex-

in

erse

ck

tu-

bid

oe-

or-

n-

e-

he

he

e-

a

a

ch

n-

re

en

0-

e

to

ie

)-

at

le

r

c

d

11

d

the passage of the optic nerve, and of the branch of the internal carotid artery that is fent to the eye.—The fecond is the foramen lacerum, or large flit between the transverse spinous and orbitar processes: the interior end of which flit is large; and, as it is extended outwards, it becomes narrower. The outer end of it is formed in the os frontis; and therefore this might be reckoned among the common foramina. Through it the third, fourth, the first branch of the fifth, and the greater share of the fixth pair of nerves, and an artery from the internal carotid, go into the orbit. times a small branch of the external carotid enters near its end, to be distributed to the dura mater (d), and a vein, some call it the venous duct, or Nuck's aquaduct, returns through it to the cavernous finus.—The third hole, fituated a little below the one just now described, is called rotundum, from its shape. It allows passage to the fecond branch of the fifth pair of nerves, or superior maxillary nerve, into the bottom of the orbit.-The fourth is the foramen ovale, about half an inch behind the round hole. Through it the third branch of the fifth pair, or inferior maxillary nerve, goes out; and fometimes a vein from the dura mater passes out here (e).—Very near the point of the spinous process is the fifth hole of this bone: it is small and round, for a passage to the largest artery of the dura mater, which often is accompanied with a vein—The fixth proper hole (f) cannot be well feen, till the cuneiform bone is separated from all the other bones of the cranium; for one end of it is hid by a small protuberance of the internal plate of the pterygoid process, and by the point of the processus petrosus of the temporal bone. Its ca-

<sup>(</sup>d) Winflow, Exposition anatomique du corps humain, traité des arteries, § 60. et de la tete, § 26.

<sup>(</sup>e) Ingraff. Commentar. in Galen. de offib lib. 1. comment. 8. (f) Vefal. Anat. lib. 1. cap. 12.—Eustach. tab. 46. fig. 13. & 16.—Vidus Vidius, Anat. lib. 2. cap. 2. explicat. tab. 5. & tab. 6. fig. 8, 9, 10. lit. O.

Ch

pa

pa

pe

its

ar

an

at

by

la

th

th

tv

th

01

63

of

b

n

th

fu

P

ai

b

P

e

tl

nal is extended above the inner plate of the pterygoid process; and where it opens into the cavity of the nose, it is concealed by the thin laminous part of the palatebone. Through it a considerable branch of the second branch of the fifth pair of nerves is reslected.—Often, in the middle of the sella turcica, a small hole or two pierce as far as the cellular substance of the bone; and sometimes at the sides of this sella, one or more small holes penetrate into the sphenoidal sinuses. These observations afforded some anatomists (g) an argument of weight in their days in defence of Galen (h), who afferted the descent of the pituita that way into the sinuses below.

The first of the common holes is that unequal fissure at the fide of the fella turcica, between the extreme point of the os petrofum and the spinous process of the cunei-This hole only appears after the bones form bone. are boiled; for in a recent subject its back-part is covered by a thin bony plate that lies over the internal carotid artery, and further forward it is filled with a cartilaginous ligament, under which the cartilaginous part of the Eustachian tube is placed: It was by this pasfage that the ancients believed the flimy matter was conveyed from the eumunctory of the brain, the glandula pituitaria, to the fauces —— The fecond commou hole is the large discontinuation of the external fide of the orbit, left between the orbitar processes of the cuneiform bone, the os maxillare, malæ, and palati. In this large hole the fat for lubricating the globe of the eye and temporal muscle is lodged, and branches of the fuperior maxillary nerve, with fmall arteries from the carotid and veins, pass. The third hole is formed between the base of this bone and the root of the orbitar process of the palate-bone of each fide. this a branch of the external carotid artery, and of the fecond branch of the fifth pair of nerves, are allowed a paffage

(b) Galen. de usu part. lib. 9. cap. 1.

<sup>(</sup>g) Jac. Sylv. Calumniæ fecundæ amolitio.—Laurent. Hist. Anat. lib. 2. quest. 11.

I.

id le,

nd

n,

NO

nd

all

er-

of

ed

w.

at

nt ei-

es

o-

ir.

af-

as

nou

of

u-

In

he

ot

m

n-

he

o'

he

la

ge

at.

passage to the nostrils, and a returning vein accompanies them. Sometimes, however, this hole is proper to the palate-bone, being entirely formed out of its substance.

Under the fella turcica, and fome way farther forward, but within the fubstance of the sphenoid bone, are two finuses, separated by a bony plate. Each of them is lined with a membrane, and opens into the upper and back part of each nostril by a round hole, which is This hole is not formed only at their upper fore-part. by the os fphenoides, which has an aperture near as large as any transverse section of the sinus, but also by the palate-bones, which are applied to the fore-part of these sinuses, and close them up, that hole only excepted which was already mentioned. Frequently the two finules are of unequal dimensions, and sometimes there is only one large cavity, with an opening into These cavities are likewise said (i) to be one nostril. extended fometimes as far back as the great foramen of the occipital bone. In other subjects they are not to be found, when the bone is composed of large cells (k). Some (1) mention a cavity within the partition of the fi-The fphenoidal finuses serve nuses; but it is small. the fame uses as the frontal do.

As this bone is extremely ragged and unequal, so its substance is of very different thickness, being in some places diaphanous; in others it is of a middle thickness, and its middle back-part surpasses the greatest share of the cranium in thickness.

The os sphenoides is joined by its wings to the parietal bones above, to the os frontis and ossa malarum before, to the temporal bones behind;—by the forepart of its body and spinous processes, to the frontal and ethmoid bones;—by its back-part, behind the two sinuses, to the occipital, where it looks like a bone with the epiphyses taken off, and, as was formerly observed

in

<sup>(</sup>i) Albin. de offib. § 39. (k) Vefal. lib. 1. cap. 6.

<sup>(1)</sup> Id. ibid.

Ch

the

by

giv

W

mo

cu

m

no

th

it

fu

ar

n

tr

0

ri

i

1

in the description of the occipital bone, it cannot be separated without violence in adults;—to the palatebones, by the ends of the pterygoid processes, and still more by the fore-part of the internal plates of the pterygoid processes, and of the sinuses;—to the maxillary bones, by the fore-part of the external pterygoid plates;—to the vomer and nasal plate of the os ethmoides, by the processus azygos. All these conjunctions, except the last, which is a schindylesis, are said to be by the suture proper to this bone; though it is at first sight evident, that several other sutures, as the transverse, ethmoidal, &c. are consounded with it.

We see now how this bone is joined to all the bones of the cranium, and to most of the upper jaw; and therefore obtained the name of the wedge-like bone.

The uses are so blended with the description, as to leave nothing new to be added concerning them.

The sphenoidal bone is almost complete in a sætus of nine months; only the great alæ separate after maceration from the body of the bone.—The processus azygos is very large and hollow;—the thin triangular processes are not offissed;—the internal surface of the body is unequal and porous;—the sinuses do not

appear.

Whoever is acquainted with each bone of the cranium, can, without difficulty, examine them as they stand united, so as to know the shapes, sizes, distances, &c. of their several parts, and the forms, capacities, &c. of the cavities formed by them; which is of great use towards understanding the anatomy of the parts contiguous to, contained within, or connected to them. Such a review is necessary, after considering each class of bones. Thus the orbits, nostrils, mouth, face, head, spine, thorax, pelvis, trunk, extremities, and skeleton, ought likewise to be examined.

rt I.

fe-

atestill

ote-

lary

es;

ex-

by

irst

nf-

the

w;

ne.

to

tus

na-

Tus

u-

of

ot

ra-

ey

n-

a-

of

he

to

ng

h,

s,

he

# § 2. Bones of the FACE.

THE face is the irregular pile of bones composing the fore and under part of the head, which is divided by authors into the upper and lower maxillæ or jaws.

The fuperior maxilla (m) is the common defignation given to the upper immoveable share of the face; tho, if we would follow Celfus (n), we should apply the word maxilla to the lower jaw only, and the name mala to this upper jaw. In complaisance to prevailing custom, he shall, however, use the terms as now commonly employed. The shape of the superior jaw cannot easily be expressed; nor is it necessary, provided the shape and situation of all the bones which compose it are described. It is bounded above by the transverse suture, behind by the fore-part of the sphenoid bone, and below by the mouth.

The upper jaw consists of six bones on each side; of a thirteenth bone which has no fellow, placed in the middle; and of sixteen teeth. The thirteen bones are, two ossa nasi, two ossa unguis, two ossa malarum, two ossa maxillaria, two ossa palati, two ossa spongiosa infe-

riora, and the vomer.

The offa nasi are placed at the upper part of the nose;
—ossa unguis are at the internal canthi of the orbits;
—ossa malarum form the prominence of the cheeks;—
ossa maxillaria form the side of the nose, with the whole lower and fore part of the upper jaw, and the greatest share of the roof of the mouth;—ossa palati are situated at the back-part of the palate, nostrils, and orbit;—ossa spongiosa are seen in the lower part of the nares;—and the vomer helps to separate these two cavities.

The bones of the upper jaw are joined to the bones of the skull by the schindylesis and sutures already described as common to the cranium and face, and they

are

<sup>(1)</sup> Σιαγων, γινος, mandibula.

<sup>(</sup>m) Lib. 8. cap. 1.

Cha

of

of t

rat

eac

jec

or we

tur be

har

iku

for

ne

int

Th

bo

pr

6

al

th

CO

fh

th

ro

fi

W

ef

ti

are connected to each other by gomphosis and fifteen futures.

The gomphosis only is where the teeth are fixed in their fockets, and the schindyless is only where the edges

of the vomer are joined to other bones.

The futures are generally distinguished by numbers, which have been differently applied; and therefore I join those (n) who prefer the giving names to each, which may be easily contrived from their situation, or from the bones which they connect.

The first is the anterior nasal (0), which is straight, and placed longitudinally in the middle fore-part of

the nofe.

The second and third are the lateral nasal (p), which are at each side of the nose, and almost parallel to the first suture.

Each of the two lacrymal is almost semicircular, and

is placed round the lacrymal groove.

The fixth and feventh are the internal orbitar: each of which is extended obliquely from the middle of the lower fide of an orbit to the edge of its base.

The two external orbitars are continued, each from the end of the internal orbitar to the under and fore-

part of the cheek.

The tenth is the mystachial, which reaches only from the lower part of the septum narium to between the two middle dentes incisores.

The longitudinal palate (q) future stretches from the middle of the foremost teeth through the middle of all the palate.

The tranverse palate (r) runs across the palate, nearer

the back than the fore-part of it.

Each

<sup>(</sup>n) Vander Linden. Medicin. physiolog. cap. 13. art. 2. § 10.

—Rolfinc. Anat. lib. 2. cap. 25.—Schenk. Schol. part. § ult.
par. 2. cap. 5.

<sup>(0)</sup> Nafalis recta. (p) Nafalis obliqua.

 <sup>(</sup>q) Laquearis, palataria recta.
 (r) Arcuata, palatina postica.

Each of the two palato-maxillary is at the back-part of the fide of each nostril.

The fifteenth is the spinous, which is in the middle of the lower part of the nostrils. This may perhaps be

rather thought a double schindylesis.

rt I.

een

in

ges

ers,

e I

ch,

or

ht,

of

ich

the

and

ach

the

om

re-

cm

wo

the

all

rer

ich

ult.

The connection of the offa spongiosa to the side of each nostril, is so much by a membrane in young subjects, by a fort of hook and afterwards by concretion or union of substance in adults, that I did not know well how to rank it: But if any chooses to call it a suture, the addition of two transverse nasal sutures may be made to those above named.

These sutures of the face (formerly called harmoniae) have not such conspicuous indentations as those of the skull have; the bones here not having substance enough for forming large indentations, and there being less necessity for security against external injuries, or any internal protruding force, than in the cranium.—

These sutures often disappear in old people, by the bones running into each other; which can do little prejudice, because the principal use of the bones being so numerous here, is to allow them to be extended into a proper form.

It is evident, from the manner of the conjunction of these bones, that they can have no motion, except in

common with the cranium.

The purposes which this pile of bones serves, will be shown in the description which I am to give of each of them.

## OSSA NASI.

Ossa NASI, so named from their situation at the root of the nose, are each of an irregular oblong square figure, being broadest at their lower end, narrowest a a little higher than their middle, and becoming somewhat larger at the top, where they are ragged and thickest, and have a curvature forwards, that their connection with the frontal bone might be stronger.—These bones

Cha

thof

upo

con

hin

mo

can

par

tha

fift

for

tha

inf

a f

its

car

or

ge

op

ot

CO

ve

of

pi

th

ti

b

C

C

fi

e

bones are convex externally, and thereby better refift any violence from without; and they are concave in-

ternally, for enlarging the cavity of the nofe.

The lower edge of these bones is unequal, and is stretched outwards and backwards, to join the cartilages of the nostrils.—Their anterior side is thick, especially above, and unequal, that their conjunction to each other might be stronger; and a small rising may be remarked on their inner edge, where they are sustained by the septum narium.—Their posterior side, at its upper half, has externally a depression, where it is overlopped some way by the maxillary bones, while its lower half covers these bones: By which contrivance, they do not yield easily to pressure applied to their forepart or sides.

A finall hole is frequently to be observed on their external furface, into which two, three, or four holes, which appear internally, terminate for the transmission of small veins; sometimes the holes go on further than

the cancelli of the bones.

The nasal bones are firm and solid, with very few cells or cancelli in them; the thin substance of which

they confift not requiring much marrow.

They are joined above to the frontal bone, by the middle of the transverse survey.—behind, to the maxillary bones, by the lateral nasal sutures;—below, to the cartilages of the nose;—before, to one another, by the anterior nasal suture;—internally, to the septum narium.

The use of these bones is to cover and defend the root

of the nofe.

In an infant the nafal bones are proportionally shorter, and less thick at their upper part, than in an adult; but are otherwise complete.

# OSSA UNGUIS, OR LACRYMALIA.

Ossa unguis, or LACRYMALIA, are so named, because their figure and magnitude are something near to those tI.

fift

in-

is

la-

oe-

ch

re-

ed

its

is

its

e,

e-

ir

s,

n

in

W

h

e

0

t

those of a nail of one's finger, and because the tears pa's upon them into the nose.

Their external surface is composed of two smooth concavities and a middle ridge. The depression behind forms a small share of the orbit for the eye- all to move on; and the one before is a deep perpendicular canal, or fossa, larger above than below, containing part of the lacrymal fac and duct. This is the part that ought to be pierced in the great operation for the fistula lacrymalis.——This fossa of the bone is cribriform, or has a great number of small holes through it, that the filaments from the membrane which lines it. infinuating themselves into these holes, might prevent a feparation of the membrane, and fecure the bone in its natural fituation.—The ridge between these two cavities of the os unguis is the proper boundary of the orbit at its internal canthus; and beyond which furgeons should not proceed backwards in performing operations here. The internal or posterior surface of this bone confifts of a furrow in the middle of two convexities.

The fubstance of the os unguis is as thin as paper, and very brittle; which is the reason that those bones are often wanting in skeletons, and need little force to

pierce them in living fubjects.

Each of these bones is joined, above, to the frontal bone, by part of the transverse suture;—behind, to the os planum of the ethmoid bone, by the same suture;—before, and below, to the maxillary bone, by the lacrymal suture—Internally, the ossa unguis cover some of the sinus ethmoidales; nay, are really continuous with the bony lamellæ which make up the sides of these cells; so that they are as much part of the ethmoid bone as the ossa plana.

These unguisorm bones compose the anterior internal parts of the orbits, lodge a share of the lacrymal sac and duct, and cover the ethmoid cells.—Their situation and tender substance make a rash operator in

H

Ch

and

mu

fur

mo

fio

int

for

pro

flit

lar

bu

th

ve

fe

in

ill

la

tl

l

danger of destroying a considerable share of the organ of smelling, when he is performing the operation of the sistual lacrymalis; but when these bones are hurt, they cast off without much difficulty, and consequently the wound is soon cured, unless the patient labours under a general cacoethes, or there is a predisposition in the bones to caries; in which case, a large train of bad symptoms follow, or at best the cure proves tedious.

These bones are fully formed in a new-born child.

### OSSA MALARUM.

Ossa Malarum (s) was the name given by Celsus, as was already remarked, to all the upper jaw; but is now appropriated to the prominent square bones which form the cheek on each side.—Before, their surface is convex and smooth; backward, it is unequal and concave, for lodging part of the crotaphyte muscles.

The four angles of each of these bones have been reckoned processes by some. The one at the external canthus of the orbit, called the superior orbitar process, is the longest and thickest.——The second terminates near the middle of the lower edge of the orbit in a sharp point, and is named the inferior orbitar process.—The third, placed near the lower part of the cheek, and thence called maxillary, is the shortest, and nearest to a right angle.—The fourth, which is called zygomatic, because it is extended backwards to the zygoma of the temporal bone, ends in a point, and has one fide straight, and the other slopping. --- Between the two orbitar angles there is a concave arch, which makes about a third of the external circumference of the orbit, from which a fifth process is extended backwards within the orbit, to form near one third of that cavity; and hence it may be called the internal orbitar process.—From the lower edge of each of the offa malarum, which is between the maxillary and zygomatic processes, the masseter muscle takes its origin; and

<sup>(</sup>s) Jugalia vel zygomatica, hypopia, fubocularia.

rt I.

gan

the

hey

the

der

the

bad

d.

lus,

t is

ich

ace

ind

een

ex-

tar

nd

or-

tar

of

ft,

is

to

nd Be-

h,

e-

d.

rd

al

ne

d

and from the exterior part of the zygomatic process, the musculus distortor oris rises; in both which places the surface of the bone is rough.

On the external furface of each cheek-bone, one or more small boles are commonly found, for the transmission of small nerves or blood-vessels from, and sometimes into, the orbit.—On the internal surface are the holes for the passage of the nutritious vessels of these bones.

—A notch on the outside of the internal orbitar process of each of these bones assists to form the great slit common to this bone and to the sphenoid, maxillary, and palate bones.

The fubstance of these bones is, in proportion to their bulk, thick, hard, and solid, with some cancelli.

Each of the offa malarum is joined, by its superior and internal orbitar processes, to the os frontis, and to the orbitar process of the sphenoid bone, by the transverse suture. By the edge between the internal and inferior orbitar processes, to the maxillary bone, by the internal orbitar suture. By the side between the maxillary and inferior orbitar process, again to the maxillary bone, by the external orbitar suture. By the zygomatic process, to the os temporum, by the zygomatic suture.

The cheek-bones are entire, and fully offified in all their parts, in infants.

## OSSA MAXILLARIA SUPERIORA.

Ossa Maxillaria Superiora, are the largest bones, and constitute the far greater part, of the upper jaw, which has appropriated the name of maxillaria to them. The figure of one of them, or of the two when joined, is so irregular, that words can scarce give an idea of it.

The processes of each os maxillare may be reckoned seven. The first is the long nasal one at its upper and fore part, which is broad below, and turns smaller, as it rises upwards, to make the side of the nose. At the

H 2

Ch

nu

po

pro

bic

a b

3.

cu

ftr

the

is

pre

Th

nu

m

nu

for

an

6.

th

T

ftr

di

fo

m

fic

th

pa

al

ri

fu

d

ri

th

h

root of this, a transverse ridge may be observed within the nostrils, which supports the fore-part of the upper edge of the os spongiosum inferius. The second is produced backwards and outwards, from the root of the nasal process, to form the lower side of the orbit; and therefore may be called orbitar. The edge of this orbitar process, and the ridge of the nasal one, which is continued from it, make a confiderable portion of the external circumference of the orbit. From the proper orbitar process, a very rough triangular surface is extended downwards and outwards, to be connected to the cheek-bone; and therefore may be called the molar process, from the lowest protuberant part of which fome share of the masseter muscle takes its rise. Behind the orbitar process, a large tuberosity or bulge of the bone appears, which is esteemed the fourth process. On the internal part of this we often meet with a ridge, almost of the same height with that in the nafal process, which runs transversely, and is covered by a fimilar ridge of the palate-bone, on which the backpart of the upper edge of the os spongiosum inferius rests.—The convex back-part of this tuberosity is rough for the origin of part of the external pterygoid muscle (t); and more internally is scabrous, where the palate and fphenoid bones are joined to it.—That fpongy protuberance (u) at the lower circumference of this bone, where the fockets for the teeth are formed, is reckoned the fifth. The fixth is the horizontal plate, which forms the greater part of the base of the nostrils, and roof of the mouth: i's upper furface, which belongs to the nostrils, is very smooth; but the other below is arched and rough, for the stronger adhesion of the membrane of the mouth, which is stretched upon it; and in chewing, speaking, &c might otherwise be liable to be feparated. The feventh rifes like a spine from the inner edge of the last, and forms a small part of the partition of the nostrils.

The

<sup>(</sup>t) Albin. de offib. § 79.

t I.

hin

per

ro-

the

ind

or-

is

the

per

ex-

to

10-

ch

Be-

of

ro-

ith

na-

by

k-

ius

is

oid

he

n-

his

18

te,

ls,

e-

e-

ot

on

be

ne

rt

10

The depressions in each maxillary bone are, I. A finuofity behind the orbitar process, made by the temporal muicle. 2. A pit immediately before the fame process, where the origin of the musculus elevator labiorum communis, and elevator labii superioris, with a branch of the fifth pair of nerves, are lodged fecurely. 3. The hollow arch of the palate. 4. The semicircular great notch, or entry to the lower part of the nostrils, betwixt the root of the nasal process and spine of the palate plate.—Below this, the fore-part of the bone is flatted, or fometimes hollowed, by the musculus depreflor labit superioris. 5. Sockets for the teeth (x): The number of these sockets is uncertain; for the same number of teeth is not in all people, and the four backmost teeth of each fide of each jaw vary greatly in their number of roots; and when the teeth of a living perfon fall out or are taken away, the fockets fill up with an offeous net-work, which becomes folid afterwards. 6. The lacrymal fofia in the nafal process, which affifts the os unguis to form a passage for the lacrymal duct. The part of the bone forming this fossa is so firm and firong, that a furgeon scarce can perforate it with the ordinary instruments for the fistula lacrymalis; and therefore ought to avoid it in doing this operation.——Immediately on the outside of this, there is a small depresfion, from which the inferior or leffer oblique muscle of the eye has its origin (y). 7. The canal on the upper part of the great tuberofity within the orbit, which is almost a complete hole; in this a branch of the superior maxillary nerve passes. Besides these, the superior furface of the great bulge is concave, to receive the under part of the eye. Immediately above the transverse ridge in the nasal process, a small hollow is formed by the os spongiosum. In some subjects, the nasal process has a fmall round pit above the lacrymal duct, where the H 3

(v) Winflow, Exposition anatomique des os secs, § 276.

<sup>(</sup>x) Βοδρια, ολμισκοι. Alveoli, fossulæ, mortariola, fræna, locelli, cavæ, prassepiola, loculamenta.

Cha

at tl

plac

in:

leav

whi

mor

rat

app

in :

the

inf

are

win

oft

po Th

vic

wh

in Th

us

th

fe

ta

fu of

di

bi

h

ř

the little tendon or ligament of the orbicular muscle of the eye-lids is inserted. It is this tendon, and not the tendon of the larger oblique muscle of the eye, which there is some hazard of cutting in the operation of the

fiftula lacrymalis.

The holes of this bone are two proper and two common, which are always to be found; besides several others, whose magnitude, number, &c. are uncertain. The first of the proper is the external orbitar, immediately below the orbit, by which the infra-orbitar branch of the fecond branch of the fifth pair of nerves, and a small artery, come out, after having passed in the canal, at the botom of the orbit, described numb. 7. of the depressions. This hole is often double, and that when the nerve has happened to split before it has escaped from the bone. The second is the foramen incisivum, just behind the fore-teeth; which, at its under part, is one irregular hole common to both the maxillary bones when they are joined; but, as it afcends, foon divides into two, three, or fometimes more holes; fome of which open into each nostril. Through them small arteries and veins, and a twig of the fecond branch of the fifth pair of nerves, pass, and make a communication between, or join the lining coats of the nofe and mouth. In some subjects, Steno's duct may be traced some way on the fide of these passages next to the nose, and small orifices may be observed opening into the mouth.

The first common hole is that which appears at the inner side of the back-part of the tuberosity and of the sockets of the teeth; and is formed by a fossa in this bone, and a corresponding one in the os palati: through it a nerve, which is a branch of the second branch of the sisth pair, runs to the palate. The other common hole is the great slit in the outside of the orbit, described already as the second common hole of the

Iphenoid bone.

On the nasal process often holes may be observed for the passage of vessels to the substance of the bones; and,

,

rt L

e of

the

nich

the

omeral

ain.

itar

ves.

the

of hat

sca-

ifi-

art,

ary

oon

me

nall of

ion

ith.

vav

nall

at

ind

iti:

ond her

bit,

the

for

nd,

at

at the back-part of each tuberofity, several foramina are placed, for the transmission of nerves to the cavity within: but these are uncertain.

All the body of the maxillary bone is hollow, and leaves a large finus a-kin to the frontal and sphenoid, which is commonly, but unjustly, called antrum Highmorianum (z). When the os maxillare is fingle, or feparated from all the other bones of a skeleton, its antrum appears to have a large aperture into the nostrils; but, in a recent subject, it is so covered at its back-part by the palate bone, in the middle by the os spongiosum inferius, before by a strong membrane, that one or fometimes two holes, fcarce larger than a crow-quill, are only left at the upper part; which, after a short winding progress, open into the nostrils between the two offa spongiosa.—At the bottom of this cavity, we may often observe some protuberances, in which the small points of the roots of the teeth are contained (a). This cavern and the fockets of the teeth are often divided by the interpolition only of a very thin bony plate, which is liable to be eroded by acrid matter collected in the antrum, or to be broke in drawing a tooth (b). The fymptoms of a collection of matter here naturally led us to the practice of pulling out the teeth, and piercing through this plate into the antrum, to procure an evacuation of the collected matter; by which confiderable fervice is frequently done (c).

The maxillary finuses have the same uses as the frontal and sphenoidal; and the situation of the sinuses is such, that the liquor drilling from them, from the cells of the ethmoid and palate bones, and from the lacrymal ducts, may always moisten all the parts of the membrane of the nares in the different situations which the

head is in.

H 4 Though

(z) Genæ.

(a) Highmore, Difquis. Anat. lib. 3. part. 2. cap. 1.

(b) Highmore, ibid.

<sup>(</sup>c) Cowper in Drake's Anthropol. book 3. chap. 10,—Medical Effays and Observ. vol. 5. art. 30.

Part I,

Ch

ed

or

the

or

tra

tio

th

ho

wi

fo

th

at

fc

b

0

iı

P

b

0

6

Though the membranes which line the frontal, sphenoidal, and maxillary sinuses, are continuations of the one which covers the bones within the nose; yet they are much thinner than it is, and have so much smaller vessels, that the injection which makes the membrane of the nose red all over, fills only some sew vessels of the makes, and scarce is observed in the frontal and sphenoidal. Are not the larger vessels intended for a more pleatiful secretion of a viscid liquor to defend the membrane from the effects of the perstatus, which is constantly through the nose? Are not the membranes which have the smallest vessels, cateris paribus, the most sensible? Are not many phanomena of smelling, instammations of these parts, megrim, polypi, &c. depending on this structure of these membranes?

The substance of the ossa maxillaria is compact and firm, except at the inserior processes, in which the

teeth are lodged, where it is very fpongy.

The maxillary bones are joined, above, by the upper ends of their nafal processes to the os frontis, by the transverse suture;—at the sides of these processes, to the offa unguis, by the lacrymal futures;—to the nafal bones, by the lateral nafal futures;—by their orbitar processes, to the cheek-bones, by the external orbitar futures;—by the internal fides of the internal orbitar processes, to the osla plana, by part of the ethmoidal future; --- by the back part of the tuberofities, to the palate-bones, by the future palato-maxillares; -----by the posterior edges of their palatine lamellæ, to the offa nafal spines, to the vomer, by the spinous suture; by their fockets, to the teeth, by gomphosis; -----by the internal edge of the palate-plate, to one another, by the longitudinal palate future, on the upper and fore-part of which a furrow is left for receiving the cartilage which forms the partition of the nostrils; between the fore-part of the nostrils and mouth, to each other, by the mystachial suture :- sometimes they are connect-

ed

ed to the offa spongiofa inferiora, by a plain concretion or union of lubitance.

These bones form the greater part of the nose and of the roof of the mouth, and a confiderable share of the They contain fixteen teeth, give rife to mulcles, transmission to nerves, &c. as mentioned in the descrip-

tion of their leveral parts.

rt I,

he-

the

hey

ller

e of the

ntal

tor end

ich

nes

tor ım-

ing

and the

per the

the

afal tar

tar

itar

dal

the -by

offa

eir

the

the

art

ige

een

er,

a-

ed

In each of the maxillary bones of a new-born child, the external orbitar process is hollow, with remarkable holes in it; ——there are five fockets for the teeth, of which the two posterior are very large, and, when divided by a fecond crofs partition, make the number of fockets fix (d). The palate-plate is cribriform about The great tuberofity is not formed;—inthe middle. flead of the antrum, there is only an oblong depression at the fide of the nostrils.

### OSSA PALATI.

Ossa Palati are commonly described as two small fquare bones, at the back-part of the palate or roof of the mouth, though they are of much greater extent, being continued up the back-part of the nostrils to the orbit (e). Each palate-bone may therefore be divided into four parts, the palate fquare-bone, the pterygoid

process, nasal lamella, and orbitar process.

The square-bone is unequally concave, for enlarging both the mouth and cavity of the nofe. The upper part of its internal edge rifes in a spine, after the same manner as the palate-plate of the maxillary bone does, to be joined with the vomer. Its anterior edge is unequally ragged, for its firmer connection with the palate-process of the os maxillare. The internal edge is thicker than the rest, and of an equal surface, for its conjunction with its

(d) Albin. Osteogen. tab. 5. fig. 45. Ungebav. de dentit. fecund. jun. § 1.

<sup>(</sup>e) Eustach. tab. 47. fig. 1, 3, 6, 7, 8. - Vidus Vidius, de Anat. lib. 2. cap. 2 explicat. tab. 6. fig. 19. - Winflow, Memoires de l'acad. des sciences, 1720.

CI

th

of

ne

ta

bo

of th

pl

A h

la

P

p

it

n

u 18

a

its fellow of the other fide. Behind, this bone is fomewhat in form of a crescent and thick, for the firm connection of the velum pendulum palati; the internal point being produced backwards, to afford origin to the palato-staphylinus or azygos muscle. This square bone is well diffinguished from the pterygoid process by a perpendicular fosfa, which, applied to such another in the maxillary bone, forms a passage for the palatine branch of the fifth pair of nerves; and by another fmall hole behind this, through which a twig of the

fame nerve passes.

The pterygoid process is somewhat triangular, having a broad bate, and ending fmaller above. The backpart of this process has three fossæ formed in it; the two lateral receive the ends of the two plates of the fphenoid bone, that are commonly compared to a bat's wing; the middle fossa makes up a part of what is commonly called the fossa pterygoidea; the fore-fide of this palatine pterygoid process is an irregular concave, where it receives the back-part of the great tuberofity of the maxillary bone. Frequently feveral small holes may be observed in this triangular process, particularly one near the middle of its base, which a little above communicates with the common and proper holes of this bone already taken notice of.

The nafal lamella of this bone is extremely thin and brittle, and rifes upwards from the upper fide of the external edge of the square bone, and from the narrow extremity of the pterygoid process; where it is so weak, and at the same time so firmly fixed to the maxillary bone, as to be very liable to be broken in separating the bones. From the part where the plate rifes, it runs up broad on the infide of the tuberofity of the maxillary bone, to form a confiderable share of the sides of the maxillary finus, and to close up the space between the fphenoid and the great bulge of the maxillary bone, where there would otherwife be a large flit opening into the nostrils (f). From the middle internal fide of thes rt I.

is

rm

nal

to

are

no-

pa-

ner

the

ing

khe

he

it's

mhis

ere

he

ay

ne

mhis

nd

he

k,

ry

ng ns

il-

of

en

e,

naf

15

this thin plate, a cross ridge, placed on such another of the maxillary bone, is extended; on it the back-part of the os spongiosum inserius rests. Along the outside of this plate, the perpendicular sossa made by the palatenerve is observable.

At the upper part of this nafal plate, the palate-bone divides into two processes, which I already named orbitar;—between which and the body of the sphenoid bone, that hole is formed which I mentioned as the last of the holes common to the sphenoid bone. Sometimes this hole is wholly formed in the os palati, by a crofs plate going from the one orbitar process to the other. A nerve, artery, and vein, belonging to the nostrils, pass here.—The anterior of the two orbitar processes is the largest, and has its fore-part contiguous to the backpart of the maxillary finus, and its upper furface appears in the bottom of the orbit, behind the backpart of the os maxillare and planum. It has cells behind, refembling those of the ethmoid bone, to which it is contiguous; it is placed on the aperture of the finus sphenoidalis, so as to leave only a round hole at its upper fore-part.—The other part of the orbitar process is extended along the internal fide of the upper backpart of the maxillary tuberofity, to the base of the sphenoid bone, between the root of the processus azygos and the pterygoid process.

The palate square part of this plate-bone, and its pterygoid process, are firm and strong, with some cancelli; but the nasal-plate and orbitar processes are very thin and brittle.

The palate-bones are joined to the maxillary, by the fore-edge of the palate square bone, by the transverse palate suture:—By their thin nasal plates, and part of their orbitar processes, to the same bones, by the palatomaxillares sutures:—By their pterygoid processes, and back-part of the nasal plates, to the alæ vespertilionum, by the sphenoid suture:—By the transverse ridges of the nasal palates, to the ossa spongiosa inferiora, by con-

Cha

fpo

eac

fma

the

but

ber

ipo ver

un

and

gre

elt

tha

pa

be

fu

in

bo

th

th

n

It

1

tact; hence frequently there is an intimate union of the fubilance of these bones in old skulls:—By the orbitar processes, to the ossa plana and cellulæ ethmoideæ, by the ethmoid suture: —To the body of the sphenoid bone, by the sphenoid suture:—By the internal edge of the square bones, to each other by the longitudinal palate suture; and by their nasal spines, to the vomer, by the spinous suture.

The palate-bones form part of the palate, nostrils, orbits, and fossæ pterygoideæ; and they cover part of the finus maxillares, sphenoidales, and ethmoidei.

These bones are very complete in a new-born infant, the nasal plates being then thicker and stronger than in adults; but the orbitar processes have not the

cells which appear in the bones of adults.

When we are acquainted with the history of these bones, the reason is evident, why the eyes are so much affected in ulcers of the palate, as to be often attended with blindness, which frequently happens in an ill-managed lues venerea; or why, on the other hand, the palate suffers from an ægylops (g).

### OSSA TURBINATA.

Ossa Turbinata, or spongiosa inferiora, resemble the superior ossa spongiosa in shape and substance, but have their anterior and upper edges contiguous to the transverse ridges of the nasal processes of the maxillary and palate bones.——From their upper straight edge, two small processes stand out: The posterior, which is the broadest, descends to cover some of the antrum Highmorianum; the anterior rises up to join the os unguis, and to make part of the lacrymal duct.

Below the spongy bones already mentioned, there are sometimes two others, one in each nostril, which seem to be a production of the sides of the maxillary sinus turned downwards (b). When this third sort of

fpongy

(g) Hossman in Fphemerid. Germ. cent. 1 and 2. obs. 135. (b) Cowper in Drake's Anthropolog. book 3. chap. 10. I.

he

r-

01-

he

n-

)e

to

S.

of

1.

er

ne

fe.

h

d

1-

l,

ıt

y

,

S

1

1

fpongy bones is found, the middle one of the three in each nostril is the largest, and the lowest is the smallest. -Befides all these, there are often several other fmall bones standing out into the nostrils, that, from their shape, might also deserve the name of turbinata; but are uncertain in their bulk, figuation, and number (i).

The names of these bones sufficiently declare their fpongy substance, which has no firm external plate covering it.

They are joined to the offa maxillaria, palati, and unguis, in all fubjects, by a firm union of fubstance; and as this happens also frequently in people of no great age, fome (k) are of opinion, that they should be elleemed part of the palate-bones; others (1) think, that fince their upper edge is continued by a plate to a part of the as ethmoides, they ought to be esteemed to be a part of this bone.

Their use is, to straigen the nostrils, to afford a large furface for extending the organ of smelling, to cover part of the antra maxillaria, and to assist in forming the under part of the lacrymal ducts, the orifices of which

into the nose are concealed by these bones.

The offa turbinata are nearly complete in a newborn infant.

### VOMER.

VOMER, or bone refembling a plough-share, is the thirteenth of the upper jaw, without a fellow, forming the lower and back parts of the partition of the note (m).

The figure of this bone is an irregular rhomboid. Its fides are flat and fmooth. Its posterior edge appears

(k) d ibid. cap. 65,7.

(1) Hunauld. in Memoires de l'acad. des sciences, 1730.

<sup>(</sup>i) Santorin. Observat. Anatomicæ, cap. 5. sect. 9.

<sup>(</sup>m) Columb de re anat. lib. 1. cap. 8. — Fallop. Observat. Anatom.

Ch

vor

late

wa

fire

the

litt

fwe

the

on

fol

it i

bo

an

be

fu

of

.br

pl

W

ft

fu

10

th

f

V

f

t

in an oblique direction at the back-part of the nostrils. The upper one is firmly united to the base of the sphenoid bone, and to the nafal plate of the ethmoid; and, when it can be got separated, is hollow for receiving the processus azygos of the sphenoid. The anterior edge has a long furrow in it, where the middle cartilage of the nose enters. The lower edge is firmly united to the nafal fpines of the maxillary and palate bones. These edges of this bone are much thicker than its middle, which is as thin as the thinnest paper; by which, and the firm union or connection this bone has above and below, it can very feldom be separated entire in adults: but in a child it is much more eafily feparated entire, and its structure is more distinctly seen; wherefore I shall examine all its parts in such a subject.

Its fituation is not always perpendicular, but often inclined and bended to one fide, as well as the nafal

plate of the ethmoid bone.

The vomer is convex at its upper part; and then is straight, as it is extended downwards and forwards, where it is composed of two plates; the edges of which have a great number of small processes, disposed somewhat like the teeth of a faw, but more irregularly, and feveral of them are reflected back. Between these plates a deep fossa is left, which, so far as the top of the curvature, is wide, and has strong sides, for receiving the processus azygos of the sphenoid bone. Beyond the arch forwards, the fossa is narrower and shallower gradually to the point of the bone, receiving for fome way the nafal lamella ethmoidea; which, after the offification is complete, is so closely united to the vomer by the little processes piercing into its substance, as to prevent any feparation: on which account it has been esteemed by some (n) a part of the ethmoid bone. The middle cartilage of the nose fills up what remains of the fossa at its fore-part. The posterior edge of the

<sup>(</sup>n) Lieutaud. Essais anatomiques, sect. 1. l'os ethmoide.

tI.

ils.

ne-

nd,

ng

ior

rti-

ni-

es.

its

by

as

en-

fe-

n;

ıb-

en

fal

is

ds,

ich

ne-

nd

efe of

ei-

Be-

al-

for

he

10-

as

en he

of

he

er,

vomer, which appears above the back-part of the palate-bones, is broader above; but as it descends forwards, becomes thinner, though it is still solid and firm. The lower edge of this bone, which rests on the nasal spine of the palate and maxillary bones, has a little surrow on each side of a small middle ridge, answering to the spines of the bones of different sides, and the interstices between them. This edge and the upper one meet in the pointed fore-end of this bone.

The body of the vomer has a smooth surface, and folid, but thin substance; and towards its sides, where it is thickest, some cancelli may be observed when the bone is broken.

It is joined above to the sphenoid and ethmoid bones, and to the middle cartilage of the nose by schindyless; below, to the maxillary and palate bones by the spinous suture.

The vomer divides the nostrils; enlarges the organ of smelling, by allowing place for expanding the membrane of the nose on its sides; and sustains the palate-plates of the maxillary and palate bones, which otherwise might be in hazard of being pressed into the nostrils; while the vomer is secured from shuffling to one side or other by the double schindyless, by which it is joined to the bones above and below.

THESE, then, are all the bones which compose the upper jaw, except the teeth, which are so much akin to those of the lower jaw, that I choose to make one description serve for both, in which the differences observable in them shall be remarked, after the second part of the sace, the lower jaw, is examined; because the structure of the teeth cannot be well understood, until the case in which they are set is explained.

## MAXILLA INFERIOR.

MAXILLA INFERIOR (0), the lower jaw, confifts, only

(0) Tivos, riayav, Mandibula, facies.

only of one moveable bone, and fixteen teeth incafed into it.

This bone, which is somewhat of the figure of the Greek letter v, is situated at the lower part of the face, so as its convex middle part is forwards, and its legs are stretched back. It is commonly divided into the chin, sides, and processes.—The chin is the middle forepart; the extent of which to each side is marked on the external surface by the holes observable there, and internally by the beginning of an oblique ridge. Beyond these the sides appear; and are continued till the bone, by bending upwards, begins to form the processes.

On the fore-part of the chin, a transverse ridge appears in the middle; on each side of which the musculi quadrati, or depressors labii inferioris, and the levatores labii inferioris, depress the bone: And below these prints a small rising may be observed, where the depressors commence. On the back-part of the chin, sometimes three, always two, small protuberances appear in the middle. To the uppermost, when it is seen, the frænum of the tongue is connected. From the middle one, the musculi genioglossi rise; and from the lowest, the geniohyoidei have their origin. Below the last, we see two rough sinuosities formed by the digastric muscles.

At the lower and fore part of the external surface of each side of the lower jaw, a small eminence may be observed, where the depressor labiorum communis rises. Near the upper edge of the side a ridge runs lengthwise, to which the under-part of the musculus buccinator is connected.—Internally, towards the upper edge of each side, another ridge appears, from which the mylohyoidei have their origin, and to which the internal membrane of the gums adheres.

In the upper edge of both chin and sides, are a great may deep pits or fockets, for receiving the roots of the teeth. The number and magnitude of these fockets are various, because of the different number, as well

of

Cl

of

pe

the

fre

aft

be

of

ca

jav

m

are

th

pr

de

th

m

th

an

ve

is

pt

ter

fix

fha

in

dy

by

tra

ar

all

an

lo

ar

at

rt I.

ised

the

ice,

are

nin,

re-

the

in-

ond

ne,

ap-

uli

va-

OW

the

in,

ap-

en,

he

he

he

ga-

ot

be

es.

h-

ci-

er

ch

n-

at

ne

ts ll

of

of the teeth themselves, as of their roots, in different These fockets in this lower jaw, as well as in the upper one, are less deep as old age comes on: when freed from the teeth by any means, they are some time after filled up with an offeous net-work, which at last becomes entirely folid, and as fmooth as any other part of the bone; fo that, in a great many old jaws, one cannot observe a vestige of the sockets: but then the jaw becomes less and much narrower (p.) Hence we may know why the chin and nofe of edentulous people are much nearer than before the teeth were loft; while their lips either fall in towards the mouth, or stand prominent forwards.—When new tyeth are protruded, new fockets are formed (q). The lower edge of the chin and sides is smooth and equal, and is commonly called the base of the lower jaw. The ends of the base, where the jaw turns upwards, are called its angles; the external furface of each of which has feveral inequalities upon it, where the maffeter muscle is inferted; as the internal furface also has, where the pterygoideus internus is inferted, and a ligament extended from the styloid process of the temporal bone is fixed.

The processes are two on each side.—The anterior sharp thin coronoid ones have the crotaphite muscles inserted into them. The posterior processes or condyles (r) terminate in an oblong smooth head, supported by a cervix. The heads, whose greatest strength is transverse, and whose convexity is turned forwards, are tipped with a cartilage, as the articulated parts of all other moved bones are. The fore-part of the root and neck of these condyloid processes are a little hollow and rough where the external pterygoid muscles are inserted.

The holes of the lower jaw are two on each fide; one at the root of the processes internally, where a large Vol. I.

I branch

<sup>(</sup>p) Vefal. Anat. lib. 1. cap. 10.

<sup>(9)</sup> Fallop. Observ. Anat. (r) Articulatorii.

t

t

t

1

rum,

branch of the third branch of the fifth pair of nerves enters with an artery, and a vein returns. A fmall sharp process frequently juts out backwards from the edge at the fore-part of this hole, to which a ligament extended from the temporal bone is fixed (s), which faves the nerve and veffels from being too much preffed by the pterygoid muscles.—From the lower side of this hole, either a finall superficial canal or a furrow descends, where a branch of the nerve is lodged, in its way to the mylo-hyoideus muscle, and sublingual gland (t). The other hole is external, at the confines of the chin, where branches of the nerve and veffels come out. The canal betwixt these two holes is formed in the middle of the substance of the bone, and is pierced by a great number of small holes, by which the nerves and blood-veffels of the cancelli and teeth pass. This canal is continued a little further than the external hole at the chin.—On account of the veffels and nerves in the lower jaw, fractures of it may be attended with dangerous symptoms.

The furface of the lower jaw is hard and firm, except at the spongy sockers, where, however, it is stronger than the upper jaw. Its internal fubstance is cellular, without any solid partition between the cancelli in its middle. At the base, especially of the chin, where this bone is most exposed to injuries, the solid sides of it

are thick, compact, and hard.

The lower jaw generally receives the roots of fixteen teeth into its fockets, by gomphosis; and its condyloid processes, covered with cartilage, are articulated with the temporal bones, in a manner that is not commonly described right: for, as was already mentioned in the description of the temporal bones, not only the forepart of the cavity between the zygomatic, auditory, and vaginal processes, but also the adjoining tubercle at the root of the zygomatic process of each os tempo-

<sup>(</sup>s) Weitbrecht. Syndesmolog. sig. 32. 1. (t) Palfya. Anat.-chirur, traité 5. chap. 6.

T.

ves

he

ent

fed

of

ow

inual

els

ed

er-

is.

er-

ind led

ept

ar,

its

it

een oid

ith

nly

the

re-

ry,

cle

00

m,

rum, is covered with a smooth cartilage for this articulation. Here also an intermediate moveable cartilage is placed; which being thin in the middle, and thick at the edges, is concave on both sides; and is connected so firmly by ligaments to each condyle, as to follow the motions of the condyle; and so loosely to the temporal bone, as readily to change its situation from the cavity to the tubercle, and to return again; while the common ligament of the articulation affords space enough for such a change of place backwards and forwards; but, like other ligaments of the joints by ginglimus, is strong and short at the sides, to confine the lateral motions.

When, therefore, the teeth of both jaws coincide, the condyles are lodged fecurely in the temporal cavities; but their motions to either fide must be confined both by the firmness of the ligaments, and the rising brims which are on each fide of the cavities.—When the jaw is brought directly forwards, the condyle and intermediate cartilages defeend and advance forwards upon the tubercles.—In this fituation, the lateral motions are a little more free than in the former one, from the want of rifing brims to stop the condyles.— When the fore-teeth of the lower jaw are moved forwards and to a fide, the condyle of the opposite fide is either advanced from the cavity to the tubercle, while the condyle of the same side remains in the cavity; or if both condyles are on the tubercles, when the jaw is moved obliquely to a fide, the condyle of the fide to which the motion is made flides back from the tubercle to the cavity.——When the mouth is opened by the descent of the lower jaw, the fore-part of it, where the depressing muscles are fixed, is drawn backwards, as well as downwards, while refistance is made to the angles moving backwards by the maffeter and internal pterygoid muscles, and at the same time the external pterygoid draw the condyles and their moveable cartilages forwards; and therefore, when the

p

tl

tl

tl

fi

t

C

ft

il

U

r

e

I

t

t

1

mouth is opened, the condyles are carried forwards upon the tubercles, and the axis of motion of the bone is a little above its angles. But in this fituation there is less resistance, than in any other, to the condyles luxating forwards: a disease which seldom happens, except when people are gaping too wide; and therefore the common practice of nurses, who support the jaw of infants when they are yawning, is reasonable.— In chewing, there is a succession of the motions above described (u).

Here a general remark may be made, That whereever moveable cartilages are found in joints, either the articulated bones are of such a figure, or so joined and fixed by their ligaments, that little motion would be allowed without such cartilages; or else some motions are necessary to the right use of the member, which the form of the articulation would not otherwise admit of. This will more fully appear after the other joints with

fuch cartilages are described.

In a child born to the full time, the lower jaw is composed of two bones, connected by a thin cartilage in the middle of the chin, which gradually offifies, and the two bones intimately unite.—In each of these bones there are five or fix fockets for teeth, as in the

upper jaw.

After I have thus described the incasement of the teeth; the insertion of so many muscles of the tongue, and of the os hyoides; the connection of the membrane of the tongue to the maxillary bone, and the motions of this bone; it is easy to see, that the lower jaw must be a principal instrument in manducation, deglutition, and speech.

### THE TEETH.

THE teeth are the hard white bodies placed in the fockets

(u) For a more full account of this articulation, vid. Edinburgh Medical Essays and Observ. vol. 1. art. 11. and vol. 3. art. 12—Memoires de l'acad. des sciences, 1744.

tI.

rds

ne

ere

les

ns,

re.

he

ve

re-

he

nd

al-

are

he

of.

ith

is

ige

nd

efe

he

he

ie,

rie

ns

ult

n,

he

ets

gh

fockets of both jaws. Their number is generally fixteen above, and as many below; though fome people have more, others have fewer.

The broad thick part of each tooth which appears without the focket, is the base or body (x).—The smaller processes sunk into the maxillæ, are the roots or sangs; which become gradually smaller towards the end tarthest from the base, or are nearly conical; by which the surface of their sides divides the pressure made on the bases, to prevent the soft parts, which are at the small points of the sockets, to be hurt by such pressure. At the place where the base ends and the roots begin, there is generally a small circular depression, which some call the neck or collar.

Without the gums the teeth are covered with no membrane, and they are faid to have no proper periofleum within the fockets; but that is supplied by the resected membrane of the gums, which, after a good injection, may be evidently seen in a young subject, with the vessels from it penetrating into the substance of the teeth; and it may be discovered in any tooth recently pulled, by macerating it in water (y). The adhesion of this membrane to these roots is strengthened by the small surrows observable on them.

Each tooth is composed of its cortex or enamel, and an internal bony substance. The cortex has no cavity or place for marrow; and is so solid and hard, that saws or files can with dissiculty make impression on it. It is thickest upon the base, and gradually, as the roots turn smaller, becomes thinner, but not proportionally to the difference of the size of the base and roots.—

The sibres of this enamel are all perpendicular to the internal substance; and are straight on the base, but at the sides are arched with a convex part towards the roots (z); which makes the teeth resist the compression

I<sub>3</sub> of

<sup>(</sup>x) Corona.

<sup>(1)</sup> Cowper. Anatom. explicat. tab. 92. fig. 7. lit. B.

<sup>(</sup>z) Havers Ofteoleg. Nov. difc. 1.

an

th

fu

ha

th

111

aj

C

a

n

of any hard body between the jaws, with less danger of breaking these sibres than if they had been situated transversely. The spongy sockets in which the teeth are placed, likewise serve better to prevent such an injury than a more solid base would have done.—Notwithstanding the great hardness of this cortex, it is wasted by manducation. Hence the sharp edges of some teeth are blunted and made broad, while the rough surfaces of others are made smooth and slat, as people advance in life.

The bony part of the teeth has its fibres running straight, according to the length of the teeth. When it is exposed to the air, by the breaking or falling off of the hard cortex, it soon corrupts. And thence carious teeth are often all hollow within, when a very small

hole appears only externally.

The teeth have canals formed in their middle, wherein their nerves and blood-veffels are lodged: which
they certainly need, being conflantly wasted by the attrition they are subjected to in manducation; and for
their further growth, not only after they first appear,
but even in adults; as is evident when a tooth is taken
out: for then the opposite one becomes longer, and
those on each side of the empty socket turn boader;
so that when the jaws are brought together, it is scarce
observable where the tooth is wanting (a).

The vessels are easily traced so long as they are in the large canal, but can scarce be observed in their distribution from that to the substance of the teeth of adults. Ruysch (b) however affirms, that after injection he could trace the arteries into the hardest part of the teeth: And Leeuwenhoek (c) suspected the sibres of the cortex to be vessels. In children I have frequently injected the vessels of the teeth as far as their base: and in such as are not entirely ossisted, one can with a lucky injec-

tion)

(b) Thefaur. 10. num. 27.

<sup>(</sup>a) Ingraf. de tumor. cap. 1. p. 24, 25, 26.

<sup>(</sup>c) Arcan Natur. continuat. epift. p. 3.

t I.

of

ted

eth

in-

ot-

va-

me

ur-

ad-

ng

en

of

us

all

c-

ch it-

or

r,

'n

d

;

e

e

3.

d

1

tion fill so many vessels, as to make both the outside and inside of the cortical part appear perfectly red.—
This plentiful supply of vessels must expose the teeth to the same disorders that attack other vascular parts; and such teeth as have the greatest number of vessels must have the most numerous chances of being sezed with these diseases.

Every root of each tooth has such a distinct canal, with vessels and nerves in it. These canals in the teeth with more than one root, come nearer each other as they approach the base of the tooth; and at last are only separted by very thin plates, which being generally incomplete, allow a communication of all the canals; and frequently one common cavity only appears within the base, in which a pulpy substance composed of nerves and vessels is lodged. The condition therefore of the nerves here bears a strong analogy to that of the cutaneous nerves which serve for the sensation of touching.

The entry of the canals for these vessels is a small hole placed a little to a side of the extreme point of each root; sometimes, especially in old people, this hole is entirely closed up, and consequently the nerves and

blood-veffels are destroyed (d).

The teeth are seen for a considerable time in form of mucus contained in a membrane; afterwards a thin cortical plate and some few offcous layers appear within the membrane, with a large cavity filled with mucus in the middle; and gradually this exterior shell turns thicker, the cavity decreases, the quantity of mucus is lessened, and this induration proceeds till all the body is formed; from which the roots are afterwards produced.

In young subjects, different stamina or ruliments of teeth are to be observed. Those next the gums hinder ordinarily the deeper seated ones from making their way out, while these prevent the former from sending

14

out

<sup>(</sup>d) De la Hire, Histoire de l'acad. des sciences, 1699.

Cha

our

bell

baf

the

tha

por

See

cor

tion

oth

fid

mi

th

dy

fta

ke

m

W

or

tu

cc

n

ar

th

e

0

out roots, or from entering deep into the bony fockets of the jaws; by which they come to be less fixed.

Children are feldom born with teeth; but at two years of age they have twenty; and their number does not increase till they are about seven years old, when the teeth that first made their way through the gums are thrust out by others that have been formed deeper in the jaw, and some more of the teeth begin to discover themselves farther back in the mouth. About four. teen years of age, some more of the first crop are shed, and the number is increased. This shedding of the teeth is of good use: for if the first had remained, they would have stood at a great distance one from another; because the teeth are too hard in their outer crust to increase so fast as the jaws do. Whereas, both the second layer and the teeth that come out late, meeting, while they are foft, with a confiderable refiltance to their growth in length from those situated upon them, necessarily come out broad, and fit to make that close guard to the mouth (e) which they now form.

The teeth are joined to the fockets by gomphosis, and the gums contribute to fix them there; as is evident by the teeth falling out when the gums are any way destroyed or made too spongy; as in the scurvy or salivations: whence some (f) class this articulation with the suffarcosis.

The uses of the teeth are to masticate our aliment, and to assist us in the pronunciation of several letters.

Though the teeth fo far agree in their structure, yet, because of some things wherein they differ, they are generally divided into three classes, viz. incifores, canini, and molares.

The Incisores (g) are the four fore-teeth in each jaw, receiving their name from their office of cutting

<sup>(</sup>ε) τραγμος. (f) Drake's Anthropolog. book 4. chap. 3. (g) Γελασινοί, τομικοί, διχαςηρές, κτένες, τομείς, περισκόι, οξείς, Riforii, quaterii, primi, primores, anteriores, acuti.

The

our aliment; for which they are excellently adopted, being each formed into a sharp-cutting edge at their base, by their fore-side turning inwards there, while they are slopped down and hollowed behind (b); so that they have the form of wedges, and therefore their power of acting must be considerably increased.

Seeing, in the action of the incisores, a perpendicular compression is only necessary, without any lateral motion, they are not so firmly fixed in their sockets as the other teeth are, each having only one short root; but that is broader from before backwards, than to either side, to have the greatest strength where it is exposed to the strongest force applied to it (i).

The incifores of the upper jaw, especially the two middle ones (k), are broader and longer generally than

those of the under jaw.

t I.

ets

WO

Oes"

ien

ims

per

cour-

ed,

eth

10-

ter

as,

ite,

ist-

ip-

ke

OW

nd

ent

le-

fa-

ith

nt,

et,

re a-

ch

g

ur

112

In a new-born infant, only the outer shell of the body of these teeth is hardened.—Asterwards, when the stamina of two sets are formed, each has its own socket, those nearest to the edge of the gums being placed more forward, and the others are lodged farther back within the jaw-bones.

Canini (1), from the resemblance to dogs tusks, are one on each side of the incisores in each jaw.—The two in the upper jaw are called eye-teeth, from the communication of nerves which is said to be betwixt them and the eyes.—The two in the lower jaw are named angular or wike-teeth, because they support the angles of the mouth.

The canini are broader, longer, and stronger, than the incisores.—Their bases are formed into a sharp edge, as the incisores are; only that the edge rises into a point in the middle. Each of them has generally but one long root, though sometimes they have two (m).

(b) Or MIGNOS.

(i) Lettre fur l'ofteologie, ascribed to Du Verney.

(k) Duales.

(1) Kuvodovres. Risorii, fractorii, collaterales, collumellares.

(m) Fauchard, Chirurgien dentifte, chap. 1.

Cha

ders

as i

leis

thol

plac

whe

the

dep

equ

qua

the

grin

fpo

tion

me

jaw

tee

are

are

diff

fo

in

are

We

ber

nu

hav

Jav

the

tro

dir

The roots are crooked towards the end. The canini of the upper jaw are larger, longer, and with more crooked roots, than those of the under jaw. The form of their base is sit both for piercing and cutting, and the long crooked root of each makes it secure in the socket.

The canini of a child are in much the same condition as the incifores are.

The Dentes Molares, or Grinders (n), which have got their name because they grind our food, are generally five in each side of each jaw; in all twenty. Their bases are broader, more scabrous, and with a thinner cortical substance, than the other teeth. They have also more roots; and as these roots generally divaricate from each other, the partitions of the sockets between them bear a large share of the great pressure they suffer, and hinder it from acting on their points (o).

The base of the first grinder has an edge pointed in the middle, on its outside, resembling the canini; from which it slopes inwards till it rises again into a point.— It has generally but one root, which sometimes is long and crooked at its point.

The fecond dens molaris has two points on its base, rising near equally on its out and inside ——It has two roots, either separate or run together, but shorter than the root of the first. These two anterior grinders are much smaller than the three that are placed farther back in the mouth.

The third and fourth are very broad in their bases, with four or five points standing out; and they have three or more roots.

The fifth, called commonly dens sapientiæ (p), from its coming through the gums later than the other grinders,

<sup>(</sup>n) Μυλιται, γομφοι, μυλοι, πλατεις, τρας πρες, Maxillares, menfales, clavales, buccarum

<sup>(0)</sup> Lettre fur l'ofteologie.

<sup>(</sup>p) Expersioners \*proverse obsyrver. Senfus, intellectus, ferotini, ztastem complentes, genuini, moderatores.

ders, has four points on its base, which is not so large as the base of the third and sourth, and its roots are less numerous.

The incifores of the upper jaw, being broader than those of the lower jaw, make the superior grinders to be placed so much farther back than the lower ones, that, when they are brought together, by shutting the mouth, the points of the grinders of the one jaw enter into the depressions of the opposite grinders, and they are all equally applied to each other, notwithstanding the ine-

quality of their furface.

at I.

ni of

ore

The

ng,

in

ion

ave

neneir

ner

alfo

om

em

ind

in

om

.-

ng

fe,

wo

an

ire

ck

es,

ve

m

n-

rs,

66,

24

The numerous roots of the dentes molares prevent their loofening by the lateral pressure they suffer in grinding; and as the fockets in the upper jaw are more fpongy, and the teeth are more liable, by their fituation, to fall out (q), the grinders there have more numerous and more separated roots than in the lower jaw (r). The number, however, of the roots of the teeth of both jaws is very uncertain; fometimes they are more, fometimes fewer: frequently feveral roots are joined together; at other times they are all diffinct. The disposition of such as are distinct is alfo various; for in some the roots stand out straight, in others they feparate, and in others again they are crooked inwards. When the roots are united, we can still distinguish them, by remarking the number of small holes at their points, which determine the number of roots each tooth ought to be reckoned to have.

At the time of birth, only two dentes molares in each jaw have begun to offity, and that at little more than the base, which has several sharp points standing out from it. The temporaneous grinders are placed more directly upon the internal set than the other two classes are: sometimes there is a piece of the bone of the jaws

(9) Galen de offib. cap. 5.

<sup>(</sup>r) Fauchard. Chirurg .- dent. chap. 1.

Ch

it l

upo

an

ea

of

ce

wl

ey

de

th

be

fh

bo

W

n

fo

N

between the two fets; in other children, the two fets have no bone interposed between them.

From what has been faid, the answers to the follow.

ing queries may be given.

Why are children subject to salivation, sever, convulfion, vomiting, purging, &c. when their teeth are breeding or cutting the gums?

Why in children do the dentes incifores first cut the

gums, the canini next, and molares last?

Why do children shed their teeth?

Wherefore have these temporaneous teeth generally

no roots, or very fmall ones?

Why have these first teeth sometimes roots, and that more frequently in teeth pulled by art than in those which are shed by nature (3).

Why do these roots frequently come outwards thro'

the gums?

Whence come butter or buck teeth?

How do these teeth sometimes go into the natural row with the others, after pulling a rotten tooth near them?

How have some people got two rows of teeth in one

or both jaws (t)?

Why do the teeth of old people loofen, and then drop ot entire?

Whence arise the new sets of teeth which several old people obtain (u)?

Why are not the gums of toothless old people torn

by the hard fockets in chewing?

Why are the teeth infenfible when flightly filed or

.rafped?

How come they to be fensible of heat or cold, to be set on edge by acids, or to give an uneasy sensation when gritty or sandy substances are rubbed between them?

Why

(1) Fauchard. Chirurgien-dentifte, p. 7.

<sup>(</sup>t) Blaf Comment. ad Vesling. Syntagm. cap. 1, 3.
(u) Hoffman. in Van. Horn. Microcosm. p. 38.

rt I.

fets

ow.

vul.

are

the

ally

that

ofe

iro'

iral

ear

one

rop

old

orn

or

be

on

en

hy

Why does a person who has a pained tooth imagine it longer than any other?

What is the reason of some persons dying convulsed, upon rasping or filing down an overgrown tooth (x).

How do the teeth break and moulder away without any pain in some people, and not in others?

What parts are affected in the tooth-ach?
What are the causes of the tooth-ach?

May worms be reckoned among these causes (y)?

Why are the dentes molares most subject to that difease?

In what different manners ought the several classes of teeth to be extracted when such an operation is necessary?

Whence proceeds the violent obstinate hemorrhagy which sometimes attends the drawing of teeth (z)?

Why is it more difficult and dangerous to draw the eye-teeth than any other?

What makes it impossible frequently to draw grinders without bringing away part of the jaw-bone with them, or breaking the fangs?

Why do teeth foon replaced after being extracted become again fixed in the fockets (a)?

According to the division made of the skeleton, we should now proceed to the description of the trunk of the body. But we must first consider a bone which cannot well be said to belong to either the head or the trunk; nor is it immediately joined to any other, and therefore is very seldom preserved with skeletons. However, it-

(x) Bartholin. Anat. reformat. lib. 4. cap. 12.

(y) Jacob in Act. Hafn. vol. 5. obs. 107. — Pechlin. Observ. Medic. lib. 2. obs. 36. — Bartholin. Hist. Medic. cent. 3. hist 96.

(z) Paré, livre 6. chap. 2.—Rolfinc. lib. 2. cap. 27. & 30.—Moebii Fundam. Medicin. cap 9.—Ephemerid. German. dec. 1. ann. 3. obf. 319.—Fauchard. Chirurg.-dentifte, tom. 1. chap. 23. obferv. 7.

(a) De la Motte Chirurgie, tom. 1. chap. 4. obs. 2.—Fauchard. Chirurgien.-dentiste, tom. 1. chap. 29.

Ch

abo

cer

deu

Ea

ext

bei

wh

lag

WI

dis

mi

mı

for

po

to

ga

ve

R

nı

fo

ve

fti

fe

C

ti

V

tl

tt

il

it is generally described by authors after the bones of the face. In obedience, therefore, to the prevailing me. thod, I shall next examine the structure of the

#### OS HYOIDES.

THE os hyoides (b), is fituated horizontally be, tween the root of the tongue and the larynx. It is properly enough named byoides, from the refemblance it bears to the Greek letter, and may, for a clearer demonstration of its structure, be distinguished into its body, cor.

nua, and appendices.

The body is the middle broad part, convex before and hollow behind.——The convex fore-part is divided into two by a ridge, into the middle of which the mylo-hyoidei, and into the fides the stylo-hyoidei, muscles are inserted.—Above the ridge, the bone is horizontal; but pitted in the middle by the infertion of the two genio-hyoidei muscles, and a little hollowed more laterally by the ballogloffi.—Below the ridge, it is convex; but a little flatted in the middle by the sternohyoidei, and pitted more externally by the coracohyoidei.—The concavity behind faces backwards and downwards to receive the thyroid cartilage, when the larynx and the os hyoides are pulled towards each other by the action of the sterno-hyoidei and hyothyroidei muscles; and to its upper edge, the ligamentous membranes of the epiglottis, tongue, and thyroid cartilage, are fixed.

The cornua of the (c) os hyoides are stretched backwards from each side of its body, where often a small furrow points out the former separation; for in young subjects, the body and cornua are not one continued substance, as they come afterwards to be in adults.—These cornua are not always straight, nor of an equal length; their two plain surfaces stand obliquely sloping from

above

(c) Crura, latera inferiora.

<sup>(</sup>b) Hypsyloides, lambdoides, παρασατη, φαρυγγετερον. Os gutturis, os linguæ, os morsus Adami, assessor, os laude, bicorne.

art I.

's of

me.

be.

pro.

ears Era-

cor.

fore

dinich

dei,

e is

n of wed

e, it

no-

ico.

and the

ach

rol-

ous

car-

ck-

nall

ing

ub-

ele

th;

oill

DVC

ITIS,

above outwards and downwards.—Into the external, the cerato-gloffus is inferted above, and the thyro hyoideus muscle below; and to the one behind, the ligamentous membrane of the tongue and larynx adheres. Each of the cornua becomes gradually smaller as it is extended from the base; but ends in a round tubercle, from which a moveable cartilage stands out, which is connected to the upper process of the cartilago thyroidea.

Where the body of the os hyoides joins on each fide with its cornua, a small styliform process, called appendix (d), rises upwards and backwards, into which the musculi stylo-hyoidei alteri, and part of the hyo-glossis muscles, are fixed. From each of them a ligament is sometimes extended to the styloid processes of the temporal bones, to keep the os hyoides from being drawn too much forwards or downwards. The part of this ligament next to these processes sometimes forms into several cartilages, which afterwards offisy in old people. Ruysch (e) says, that he has seen this offisication continued as far up as the styloid processes, which were therefore joined to the os hyoides by anchyloss.

The fubstance of the os hyoides is cellular; but covered with a firm external plate, which is of sufficient strength to bear the actions of so many muscles as are inferted into it.

It is not articulated with any bone of the body, except by means of the muscles and ligaments mentioned.

The use of the os hyoides, is to serve as a solid lever for the muscles to act with, in raising or depressing the tongue and larynx, or in enlarging and diminishing the capacity of the sauces.

At birth, this bone is in a cartilaginous state; excepting a small point of bone in the middle of its body, and

(d) Crura superioria, latera superiora, ossa graniformia.

(e) Advers. Anat. dec. 3. 9 9.

Ch

pro

vex mo

hol

face

of par

fub

phy

me

the

of

a fi

lag

cal

is c

dec

ter

the

for

the

lig

pai

of mi

the

an

pr

T

fee

in each of the cornua.—The appendices frequently remain cartilaginous many years.

# SECT. II. OF THE TRUNK.

THE TRUNK confifts of the spine, pelvis, and thorax.

### § 1. The SPINE.

The spine (f) is the long pile of bones extended from the condyles of the occiput to the end of the rump. It somewhat resembles two unequal pyramids joined in a common base. It is not, however, straight; for its upper part being drawn backwards by strong muscles, it gradually advances forwards, to support the cesophagus, vessels of the head, &c. Then it turns backwards, to make place enough for the heart and lungs. It is next bended forwards, to support the viscera of the abdomen. It afterwards turns backwards, for the enlargement of the pelvis. And, lastly, it is reslected forwards, for sustaining the lowest great gut.

The spine is commonly divided into true and false vertebræ; the former constituting the long upper pyramid, which has its base below; while the salse vertebræ make the shorter lower pyramid, whose base is

above.

### TRUE VERTEBRÆ.

The true vertebræ (g) are the twenty four upper bones of the spine, on which the several motions of the trunk of our bodies are performed; from which use they have justly got their name.

Each

(f) Ραχις νωτον ακανθα, ιεςα συςιμές σαλεν, Tergum hominis ca-

(g) Στροφεις ετεοφιγγες, Spondyli, ossa orbiculata, ossa vertebrata, verticula.

Each of these vertebræ is composed of its body and

processes.

rt I.

re-

and

ded

the

ids

ht;

ong

the

rns

and

vif-

ds,

is

t.

else

py-

er-

18

er

of

ch

ch

ca-

tas

The body is the thick spongy fore-part, which is convex before, concave backwards, horizontal and plain in most of them above and below --- Numerous small holes, especially on the fore and back part of their furface, give paffage to their veffels, and allow the ligaments to enter their fubstance.—The edges of the body of each vertebra are covered, especially at the forepart, with a ring of bone firmer and more folid than the fubitance of the body any where elfe. These rings feem to be joined to the vertebræ in the form of epiphyses, but are alleged by some (b) to be the liga-They are of great use in preventing ments offified. the fpongy bodies from being broken in the motions of the trunk.

Between the bodies of each two adjoining vertebræ, a fubstance between the nature of ligament and cartilage is interposed; which seems to consist of concentrical curve fibres, when it is cut horizontally: but when it is divided perpendicularly, the fibres appear oblique and decuffating each other (i)——The outer part of the intervertebral ligaments is the most solid and hard, and they gradually become fofter till they are almost in the form of a glairy liquor in the centre; and therefore thefe fubstances were not improperly called mucous ligaments by the ancients (k). The external fibrous part of each is capable of being greatly extended, and of being compressed into a very small space, while the middle fluid part is incompressible, or nearly so; and the parts of this ligament between the circumference and centre approach in their properties to either, in proportion to their more folid or more fluid texture. The middle point is therefore a fulcrum or pivot, on

VOL. I.

(b) Fallop. Observat. Anatom.

(k) Galen. de ufu part. lib. 12. cap. 16.

<sup>(</sup>i) Blancardi Anat. reform. cap. 32.—Weitbrecht, Sydefmolog. fect. 4. 9. 15.

CI

bo

the

fta

ed

pro

its

ter

boo

bo

tog

and

boo

wh

ma

for

ma

tera

eac

rou

thre

mai

plai

terv

by 1

to a

mo whe

enti bacl

recl

of t

which the motion of a ball and focket may be made, with such a gradual yielding of the substance of the ligament, in whichever direction our spines are moved, as saves the body from violent shocks, and their dangerous consequences (1).—This ligamento-cartilaginous substance is firmly fixed to the horizontal surfaces of the bodies of the vertebræ, to connect them; in which it is assisted by a strong membranous ligament, which lines all their concave surface, and by still a stronger ligament that covers all their anterior convex surface.

We may lay it down as a general rule, notwithstanding some exceptions, That the bodies of the vertebra are smaller, and more solid above; but, as they descend, they appear larger and more spongy; and that the cartilages between them are thick, and the surrounding ligaments strong, in proportion to the largeness of the vertebra, and to the quantity of motion they perform: by which disposition, the greater weight is supported on the broadest best-secured base, and the middle of our

body is allowed a large and fecure motion.

From each fide of the body of each vertebra, a bony bridge is produced backwards, and to a fide: from the posterior end of which one slanting process rises, and another descends; the smooth, and what is generally the flattest, side of each of these four processes, which are called the oblique (m), is covered with a smooth cartilage; and the two lower ones of each vertebra are sitted to and articulated with the two upper or ascending oblique processes of the vertebra below, having their articular ligaments sixed into the rough line round their edges.

From between the oblique processes of each side, the vertebra is stretched out laterally into a process that is

named transverse.

From the back-part of the roots of the two oblique and of the transverse process of each side, a broad oblique bony

<sup>(1)</sup> Medical effays and observ. vol. 5. art. 28. (m) Articulatorii, minimi.

I.

de,

li-

ed,

er-

ous

of

nich

r li-

ind-

bræ

end,

car-

g lithe

rm:

our

ony

rom

and

hich

ooth

a are

ding

r ar-

, the

nat is

lique

lique

bony plate is extended backwards; where these meet, the seventh process of the vertebræ takes its rise, and stands out backwards: this being generally sharp-pointed and narrow-edged, has therefore been called spinal process; from which this whole chain of bones has got its name.

Besides the common ligament which lines all the internal surface of the spinal processes, as well as of the bodies, there are particular ligaments that connect the bony bridges and processes of the contiguous vertebræ together.

The substance of the processes is considerably stronger and firmer, and has a thicker external plate, than the bodies of the vertebræ have.

The feven processes form a concavity at their fore-part, which, joined to the one at the back-part of the bodies, makes a great hole; and the holes of all the vertebræ form a long large conduit (n), for containing the spinal marrow.—In the upper and lower edge of each lateral bridge, there is a notch. These are so adapted to each other in the contiguous vertebræ, as to form a round hole in each side between each two vertebræ, through which the nerves that proceed from the spinal marrow and its blood-vessels pass.

The articulations, then, of these true vertebræ are plainly double: for their bodies are joined by the intervening cartilage above described; and their oblique processes, being tipped with cartilages, are so connected by their ligaments as to allow a small degree of motion to all sides. Hence it is evident, that their centre of motion is altered in different positions of the trunk: for, when we bow forwards, the upper moved part bears entirely on the bodies of the vertebræ; if we bend back, the oblique processes support the weight; if we recline to one side, we rest upon the oblique processes of that side and part of the bodies; if we stand erect, all K 2

(n) Ispa eugrys, σωλω, Canalis.

C

**sta** 

nig

br

dy

bu

ni

th

ac

ve

10

fa

CU

ve

it

la

of

fti

th

ca

is

an

th

po

he

ar

of

th

bo

bo

n

1p

g

the bodies and oblique processes have their share in our

support.

Hence it follows, 1. That because the joints of which the spine is composed are so numerous, the spinal marrow, nerves, blood-veffels, &c. are not liable to fuch compression and over-stretching in the motion of the trunk of the body as they would otherwise be, since several vertebræ must be concerned in every motion of the fpine; and therefore a very small curvature is made at the conjunction of any two vertebræ (6). 2. That an erect posture is the furest and firmest, because the furface of contact of the fulcra is largest, and the weight is most perpendicular to them (p). 3. That the muscles which move the fpine act with greater force in bringing the trunk to an erect posture, than in drawing it to any other: For in bending forwards, backwards, or to a fide, the muscles which perform any of these actions are nearer the centre of motion; consequently the lever with which they act is shorter than when the centre of motion is on the part of the vertebra, opposite to that where these muscles are inserted; which is the case in raifing the trunk. This is extremely necessary; fince, in the deflections of the spine from a perpendicular bearing, the weight of the body foon inclines it which way we defign; whereas, in raifing us erect, this great weight must be more than counteracted. calculating the force exerted by the muscles which move the spine, we should always make allowance for the action of the cartilages between the vertebræ, which, in every motion from an erect polture, must be stretched on one fide, and compressed on the other, to both which they refift; whereas, in raifing the trunk, these cartilages affift by their fpringy force (q). 5. We are hence naturally led into the reason of our height of **stature** 

(0) Galen. de usu part. lib. 12. cap. 12.

<sup>(</sup>p) Paaw de offib. part 2. cap. 2. (q) Borelli de motu animal. pars 1. schol. ad propos. 58. Parent. Histoire de l'acad. des sciences, 1702.

t I,

our

nich

nar-

uch

the

fe-

n of

ade

t an

fur-

nt is

cles

ing-

it to

or ,

ions

le-

ntre

e to

cale

ary;

ndi-

es it

this

. In

ove

ac-

, in

hed

ooth

hele

are

t of

ure

Pa-

stature increasing in the morning, and diminishing at night (r): for the intermediate cartilages of the vertebræ being pressed all day long by the weight of our body, become more compact and thin in the evening; but, when they are relieved from this pressure in the night, they again expand themselves to their former thickness: and seeing the bulk of any part must vary according to the different diffention or repletion of the veffels composing it, we may understand how we become taller after a plentiful meal, and decrease after fasting or evacuations (s). 6. From the different articulations of the bodies and oblique processes of the vertebræ, and the different strength of the ligaments, it is plain, that they are formed so as to allow much larger motion forwards than backwards; this last being of much less use, and might be dangerous, by overfiretching the large blood-vessels that are contiguous to the bodies of the vertebræ (t). 7. The intervertebral cartilages shrivelling as they become more folid by age, is the cause why old people generally bow forwards, and cannot raise their bodies to such an erect posture as they had in their youth.

The uses of the true vertebræ are, to give us an erect posture; to allow sufficient and secure motion to the head, neck, and trunk of the body; and to support

and defend the bowels and other foft parts.

At the ordinary time of birth, each vertebra confifts of three bony pieces, connected by cartilages; to wit, the body, which is not fully offified; and a long crooked bone on each fide, on which we fee a finall share of the bony bridge, the oblique processes complete, the beginning transverse processes, and the oblique plate, but no spinal processes: so that the teguments are in no danger of being hurt by the sharp ends of these spinal processes.

(r) Waffe, Philosoph. transact. numb. 383. art. 1.

(t) Galen. de usu part. lib. 1. cap. 16.

<sup>(</sup>s) Abbe Fontenu, Histoire de l'acad. des sciences, 1725.

Cl

pe

ve

Wa

ar

pr

ac

de

by

by

th

T

be

of

In

tic

be

fe

ar

de

bi

m

ar

fc

to

10

b

n

0

P

fa

h

cesses, while a child is in its bended posture in the

womb, nor while it is squeezed in the birth.

From this general mechanism of the spine, an account is eafily deduced of all the different preternatural curvatures which the spine is capable of: For if one or more vertebræ, or their cartilages, are of unequal thickness in opposite sides, the spine must be reclined over to the thinner fide; which now fustaining the greatest share of the weight, must still be more compreffed, confequently hindered from extending itself in proportion to the other fide, which, being too much freed of its burden, has liberty to enjoy a luxuriant The causes on which such an inequality of thickness in different fides of the vertebræ depends may vary. For either it may be owing to an over-diffention of the veffels of one fide, and from thence a preternatural increase of the thickness of that part: or, which more commonly is the case, it may proceed from an obstruction of the vessels, by which the application of proper nourishment to the bony substance is hindered; whether that obstruction depends on the faulty disposition of the veffels or fluids, or if it is produced by an unequal mechanical prefiure occasioned by a paralytic weakness of the muscles and ligaments, or by a spasmodic over-action of the muscles on any side of the spine, or by people continuing long or putting themselves frequently into any posture declining from the erect one: In all these cases one common effect follows, to wit, the vertebræ, or their cartilages, or both, turn thick on that fide where the veffels are free, and remain then on the other fide where the veffels are straitened or obstructed.—Whenever any morbid curvature is thus made, a fecond turn, but in an opposite direction to the former, must be formed; both because the muscles on the convex fide of the spine being stretched, must have a stronger natural contraction to draw the parts to which their ends are fixed, and because the patient makes efforts to keep the centre of gravity of the body perperpendicular to its base, that the muscles may be relieved from a constant violent contractile state, which al-

ways creates uneafiness and pain.

rt I,

the

ac.

ural

e or

qual

ned

the

om-

f in

uch

iant

y of

may

tion

rna-

nich

an n of

ed;

ofi-

y an

no-

ine,

ves

rect

to

urn

ain or

hus

to

uft

sto

ent

ody

er-

When once we understand how these crooked spines are produced, there is little difficulty in forming a just prognosis; and a proper method of cure may be easily contrived, which must vary as to the internal medicines, according to the different causes on which the disease depends: But one general indication must be pursued by surgeons; which is, to counteract the bending force, by increasing the compression on the convex part of the curvature, and diminishing it on the concave side. The manner of executing which in particular cases must be different, and requires a very accurate examination of the circumstances both of the disease and patient. In many such cases I have found some simple directions, as to postures in which the patient's body should be kept, of very great advantage.

Though the true vertebræ agree in the general structure which I have hitherto described; yet, because of several specialities proper to a particular number, they are commonly divided into three classes, viz. cervical,

dorfal, and lumbar.

The CERVICAL (u) are the feven uppermost vertebræ; which are distinguished from the rest by these marks.—Their bodies are smaller and more solid than any others; and slatted on the fore-part, to make way for the cesophagus; or rather this slat sigure is owing to the pressure of that pipe, and to the action of the longi colli and anterior recti muscles. They are also slat behind, where small processes rise, to which the internal ligaments are sixed. The upper surface of the body of each vertebra is made hollow, by a slanting thin process which is raised on each side:—The lower surface is also evacuated, but in a different manner; for here the posterior edge is raised a little, and the one K 4

Cha

hav

COL

fro

fpi

ard

ed

ca tal

ar

co

is

re

fe

th

p

n

fi

a

b

t

before is produced a confiderable way.—Hence we fee how the cartilages between those bones are firmly connected, and their articulations are secure.

The cartilages between these vertebræ are thick, especially at their fore-part; which is one reason why the vertebræ advance forward as they descend, and have

larger motion.

The oblique processes of these bones of the neck more justly deserve that name than those of any other vertebræ. They are situated slanting; the upper ones having their smooth and almost flat surfaces sacing obliquely backwards and upwards, while the inserior oblique processes have these surfaces sacing obliquely forwards and downwards.

The transverse processes of these vertebræ are framed in a different manner from those of any other bones of the spine: For, besides the common transverse process rising from between the oblique processes of each side, there is a second one that comes out from the side of the body of each vertebra; and these two processes, after leaving a circular hole for the passage of the cervical artery and vein, unite, and are considerably bollowed at their upper part; with rising sides, to protect the nerves that pass in the hollow; and at last each side terminates in an obtuse point, for the insertion of muscles.

The *spinal* processes of these cervical bones stand nearly straight backwards, are shorter than those of any other vertebræ, and are forked or double at their ends; and hence allow a more convenient insertion to

muscles.

The thick cartilages between the bodies of these cervical vertebræ, the obliquity of their oblique processes, and the shortness and horizontal situation of their spinal processes, all conspire to allow them large motion.

The holes between the bony cross bridges, for the passage of the nerves from the spinal marrow, have their largest share formed in the lowest of the two vertebræ,

to which they are common.

So

rt I.

fee

on.

ick,

why

ave

ore

rte-

ha-

ob-

obfor-

din

the

ing

ere

ody

ing

ind

eir

hat

in

ind

ny

eir

to

er-

es,

nal

he

eir

æ,

30

li-

So far most of the cervical vertebræ agree; but they have some particular differences, which oblige us to consider them separately.

The first, from its use of supporting the head, has the name of atlas (x); and is also called epistrophea,

from the motion it performs on the second.

The atlas, contrary to all the other vertebræ of the fpine, has no body; but, instead of it, there is a bony arch.——In the convex fore-part of this arch a small rifing appears, where the mufculi longi colli are inferted; and, on each fide of this protuberance, a small cavity may be observed, where the recti interni minores take their rife.—The upper and lower parts of the arch are rough and unequal, where the ligaments that connect this vertebra to the os occipitis, and to the fecond vertebra, are fixed.——The back-part of the arch is concave, fmooth, and covered with a cartilage, in a recent subject, to receive the tooth-like process of the fecond vertebra.—In a first vertebra, from which the fecond has been feparated, this hollow makes the paffage for the fpinal marrow to feem much larger than it really is: On each fide of it a finall rough finuofity may be remarked, where the ligaments going to the fides of the tooth-like process of the following vertebra are fastened; and on each fide a finall rough protuberance and depression is observable, where the transverse ligament, which secures the tooth-like process in the finuolity, is fixed, and hinders that process from injuring the medulla fpinalis in the flections of the head.

The atlas has as little spinal process as body; but, instead thereof, there is a large bony arch, that the muscles which pass over this vertebra at that place might not be hurt in extending the head. On the back, and upper part of this arch there are two depressions, where the recti postici minores take their rise; and at the lower part are two other sinuosities, into which the

Cha

later

part

toot

aut

on

dal

cra

tel

W

18

to

th

pl

la

m

C

n

ligaments which connect this bone to the following one are fixed.

The superior oblique processes of this atlas are large, oblong, hollow, and more horizontal than in any other vertebra.—They rife more in their external than internal brim; by which their articulations with the condyloid processes of the os occipitis are firmer.—Under the external edge of each of these oblique processes is the fossa, or deep open channel, in which the vertebral arteries make the circular turn, as they are about to enter the great foramen of the occipital bone, and where the tenth pair of nerves goes out.——In feveral bodies, I have feen this fossa covered with bone.— The inferior oblique processes, extending from within outwards and downwards, are large, concave, and circular. So that this vertebra, contrary to the other fix, receives the bones with which it is articulated both above and below.

The transverse processes here are not much hollowed or forked; but are longer and larger than those of any other vertebra of the neck, from the origin and infertion of several muscles; of which those that serve to move this vertebra on the second have a considerable lever to act with, because of the distance of their infertion from the axis of revolution.

The hole for the spinal marrow is larger in this than in any other vertebra, not only on account of the marrow being largest here, but also to prevent its being hurt by the motions of this vertebra on the second one.

This large hole, and the long transverse processes, make this the broadest vertebra of the neck.

The condyles of the os occipitis move forwards and backwards in the superior oblique processes of this vertebra; but, from the sigure of the bones forming these joints, it appears, that very little motion can here be allowed to either side; and there must be still less circular motion.

In new-born children this vertebra has only the two

la

lateral pieces offified; the arch, which it-has at its forepart instead of a body, being cartilaginous.

The fecond vertebra colli is called dentata, from the tooth-like process on the upper part of its body. Some authors call it epistrophea, but improperly, since this designation is only applicable to the first, which moves

on this as on an axis.

art I.

one

rge,

ther in-

Con-Un-

effes

rte-

out

and

eral

hin

cir-

fix,

oth

ved

ny

er-

to

ble

in-

an

ar-

ng

ie.

nd

is

ng

re

ſs

0

a-

The body of this vertebra is somewhat of a pyramidal figure, being large, and produced downwards, especially at its fore-fide, to enter into a hollow of the vertebra below; while the upper part has a fquare process, with a small point standing out from it. This it is that is imagined to refemble a tooth (y), and has given name to the vertebra.—The fide of this process, on which the hollow of the anterior arch of the first vertebra plays, is convex, fmooth, and covered with a cartilage; and it is of the fame form behind, for the ligament, which is extended transversely from one rough protuberance of the first vertebra to the other, and is cartilaginous in the middle, to move on it.——A ligament likewise goes out in an oblique transverse direction, from each fide of the processus dentatus, to be fixed at its other end to the first vertebra, and to the occipital bone; and another ligament rifes up from near the point of the process to the os occipitis.

The fuperior oblique processes of this vertebra dentata are large, circular, very nearly in an horizontal position, and slightly convex, to be adapted to the inferior oblique processes of the first vertebra.—A moveable cartilage is said by some authors to be interposed between these oblique processes of the first and second vertebra; but I could never find it.—The inferior oblique processes of this vertebra dentata answer exactly to the description given of those common to all the cer-

vical vertebræ.

The transverse processes of the vertebra dentata are short, very little hollowed at their upper part, and not forked

<sup>(</sup>y) Conoides, pyrenoides, odontoides.

Cha

are ]

was

grea

prel

eith

fee

fub

will

cap

wai

rea

upo

cef

cul

fec

tat

pe

th

ed

fu

OS

u

CI

0

a

f

n

0

forked at their ends; and the canals through which the cervical arteries pass are reflected outwards about the middle substance of each process; so that the course of these vessels may be directed towards the transverse processes of the first vertebra.—Had this curvature of the arteries been made in a part fo moveable as the neck is, while they were not defended by a bone, and fixed to that bone, scarce a motion could have been performed without the utmost hazard of compression, and a stop put to the course of the liquids, with all its train of bad consequences. Hence we observe this fame mechanism several times made use of, when there is any occasion for a sudden curvature of a large artery. This is the third remarkable instance of it we have feen. The first was the passage of the carotide through the temporal bones; and the fecond was that lately described in the vertebral arteries, turning round the oblique processes of the first vertebra, to come at the great hole of the occipital bone.

The spinal process of this vertebra dentata is thick, frong, and short, to give sufficient origin to the musculi recti majores and obliqui inferiores, and to prevent the contusion of these and other muscles in pulling

the head back.

This fecond vertebra consists, at the birth, of sour bony pieces: For, besides the three which I already mentioned as common to all the vertebræ, the tooth-like process of this bone is begun at this time to be offisied in its middle, and is joined as an appendix to the body of the bone.—Lest this appendix be bended or displaced, nurses ought to keep the heads of new-born children from falling too far backwards, by stay-bands, or some such means, till the muscles attain strength sufficient to prevent that dangerous motion.

When we are acquainted with the structure and articulations of the first and second verter ræ, and know exactly the strength and connection of their ligaments, there is no difficulty in understanding the motions that

art I

which

bout

ourse

verle

e of

the

and

been

lion,

lits

this

nere

ery.

lave

ugh

tely

the

the

ck,

uf-

re-

ng

ur

dy

h-

Ti-

he

or

rn

s,

f-

r-

W

t

are performed upon or by the first; though this subject was formerly matter of hor dispute among some of the greatest anatomists (z). It is none of my purpose at present to enter into a detail of the reasons advanced by either party; but to explain the fact, as any one may fee it, who will remove the muscles, which, in a recent subject, hinder the view of these two joints, and then will turn the head into all the different positions it is capable of. The head may then be feen to move forwards and backwards on the first vertebra. as was already faid, while the atlas performs the circumgyratio upon the fecond vertegra; the inferior oblique proceffes of the first vertebra shuffling easily in a circular way on the superior oblique processes of the fecond, and its body or anterior arch having a rotation on the toothlike process, by which the perpendicular ligament that is fent from the point of the tooth-like process to the occipital bone is twisted, while the lateral ligaments that fix the proceffus dentatus to the fides of the first vertebra, and to the os occipitis, are very differently affected; for the one upon the fide towards which the face is turned by the circumgyratio is much shortened and lax, while the opposite one is stretched and made tense, and, yielding at last no more, prevents the head from turning any farther round on the axis. So that these lateral ligaments are the proper moderators of the circumgyratio of the head here; which must be larger or smaller, as these ligaments are weaker or stronger, longer or shorter, and more or less capable of being stretched. Besides the revolution on this axis, the first vertebra can move a small way to either side; but is prevented from moving backwards and forwards by its anterior arch, and by the cross ligament, which are both closely applied to the tooth-like process. Motion forwards here would have been of very bad confequence, as it would have

<sup>(</sup>z) See Euftach. de motu capitis.

have brought the beginning of the spinal marrow upon

the point of the tooth-like process.

The rotatory motion of the head is of great use to us on many accounts, by allowing us to apply quickly our organs of the fenses to objects: And the axis of rotation was altogether proper to be at this place; for, if it had been at a greater distance from the head, the weight of the head, if it had at any time been removed from a perpendicular bearing to the small very moveable joint, and thereby had acquired a long lever, would have broken the ligaments at every turn inconfiderately performed, or these ligaments must have been formed much stronger than could have been connected to such fmall bones. Neither could this circular motion be performed on the first vertebra without danger, because the immoveable part of the medulla oblongata is fo near, as, at each large turn, the beginning of the spinal marrow would have been in danger of being twifted, and of fuffering by the compression this would have made on its tender fibrils.

It is necessary to observe, that the lateral or moderator ligaments confine fo much the motion of the first vertebra upon the fecond, that though this joint may ferve us on feveral occasions, yet we are often obliged to turn our faces farther round than could be done by this joint alone, without the greatest danger of twisting the spinal marrow too much, and also of luxating the oblique processes: therefore, in large turns of this kind, the rotation is affifted by all the vertebræ of the neck and loins; and if this is not fufficient, we employ most of the joints of the lower extremities.—This combination of a great many joints towards the performance of one motion, is also to be observed in several other parts of the body; notwithstanding such motion's being generally faid to be performed by fome fingle joint only.

The third vertebra of the neck is by some called axis; but this name is applied to it with much less rea-

fon

for

be

all

fc

fo

fu

n

0

I.

on

us ur

on

ad

of

a

it,

ve

red

ch

be

fe

fo

i-}-

c

Pt

d

5

fon than to the fecond.—This third, and the three below, have nothing particular in their structure, but all their parts come under the general description formerly given, each of them being larger as they descend.

The feventh (a) vertebra of the neck is near to the form of those of the back, having the upper and lower surfaces of its body less hollow than the others: The oblique processes are more perpendicular; neither spinal nor transverse processes are forked. This seventh and the sixth vertebra of the neck have the hole in each of their transverse processes more frequently divided by a small cross bridge, that goes between the cervical vein and artery, than any of the other vertebræ.

The twelfth DORSAL (b) may be distinguished from the other vertebræ of the spine by the following marks.

Their bodies are of a middle fize betwixt those of the neck and loins:—they are more convex before than either of the other two forts; and are flatted laterally by the pressure of the ribs, which are inserted into small cavities formed in their sides. This flatting on their sides, which makes the sigure of these vertebræ almost an half oval, is of good use; as it affords a firm articulation to the ribs, allows the trachea arteria to divide at a small angle, and the other large vessels to run secure from the action of the vital organs.—These bodies are more concave behind than any of the other two classes.—Their upper and lower surfaces are horizontal.

The cartilages interposed between the bodies of these vertebræ are thinner than in any other of the true vertebræ; and contribute to the concavity of the spine in the thorax, by being thinnest at their fore-part.

The oblique processes are placed almost perpendicular; the upper ones slanting but a little forwards, and the

<sup>(</sup>a) Atlas quibusdam, maxima, magna vertebra, prominens.

<sup>(</sup>b) Θωρακος, μεταρρενν, νωτυ, υποτραχαλιν, antisterni, pectoris, tergi-

C

25

an

an

of

th

ril

pr

VE

fu

m be

m

as

W

its

in

th

th

el

OI

la

pi

fa

the lower ones flanting as much backwards.—They have not as much convexity or concavity as is worth remarking. Between the oblique processes of opposite sides, several sharp processes stand out from the upper and lower parts of the plates which join to form the spinal process: into these sharp processes strong ligaments are fixed for connecting the vertebræ.

The transverse processes of the dorsal vertebræ are long, thicker at their ends than in the middle, and turned obliquely backwards; which may be owing to the pressure of the ribs, the tubercles of which are inferted into a depression near the end of these processes.

The spinal processes are long, small-pointed, and sloping downwards and backwards: from their upper and back part a ridge rises, which is received by a small channel in the fore-part of the spinal process immediately above, which is here connected to it by a ligament.

The conduit of the spinal marrow is here more circular, but, corresponding to the fize of that cord, is smaller than in any of the other vertebræ; and a larger share of the holes in the bony bridges, for the transmission of the nerves, is formed in the vertebra above

The connection of the dorsal vertebræ to the ribs, the thinness of their cartilages, the erect situation of the oblique processes, the length, sloping, and connection of the spinal processes, all contribute to restrain these vertebræ from much motion, which might disturb the actions of the heart and lungs; and in consequence of the little motion allowed here, the intervertebral cartilages sooner shrivel, by becoming more solid: and therefore the first remarkable curvature of the spine observed, as people advance to old age, is in the least stretched vertebræ of the back; or old people first become round-shouldered.

The bodies of the four uppermost dorsal vertebræ deviate from the rule of the vertebræ becoming larger

irt I.

hey

orth

ofite

per

ipi-

ents

are

and

g to

m-

es.

and

per

v a

im-

y a

cir-

ger

nf-

ove

ibs,

of

on-

ain

urb

nce

oral

ob.

east be-

oræ.

ger

as

as they descend: for the first of the sour is the largest, and the other three below gradually become smaller, to allow the trachea and large vessels to divide at smaller angles.

The two uppermost vertebræ of the back, instead of being very prominent forwards, are flatted by the action

of the musculi longi colli and recti majores.

The proportional fize of the two little depressions in the body of each vertebra for receiving the heads of the ribs, seems to vary in the following manner; the depression on the upper edge of each vertebra decreases as far down as the fourth, and after that increases.

The transverse processes are longer in each lower vertebra to the seventh or eighth, with their smooth surfaces, for the tubercles of the ribs, facing gradually more downwards; but afterwards, as they descend, they become shorter, and the smooth surfaces are directed more upwards.

The spinous processes of the vertebræ of the back become gradually longer and more slanting from the first, as far down as the eighth or ninth vertebra; from which they manifestly turn shorter and more erect.

The first (c) vertebra, besides an oblong hollow in its lower edge, that assists in forming the cavity wherein the second rib is received, has the whole cavity for the head of the first rib formed in it.

The fecond has the name of axillary (d), without any

thing particular in its structure.

The eleventh (e) often has the whole cavity for the eleventh rib in its body, and wants the smooth surface

on each transverse process.

The twefth (f) always receives the whole head of the last rib, and has no smooth surface on its transverse processes, which are very short.——I he smooth surfaces of its inserior oblique processes face outwards as

Vol. 1. L the

(/) DIRECTOR. Przeingens.

<sup>(</sup>c) Aogia, Gutturalis. (d) Μασχαλίστης.

<sup>(</sup>e) Aggezne, in neutram partem inclinans.

C

fla

laf

th

in

co

ro

co

8.

eq

tha

ev

the

cef

is I

app

icr

obl

and

ver

ver

pro

pre

bru

lun

wh

mi

tha

is f

wai bra

(

the lumbar do.—And we may fay in general, that the upper vertebræ of the back lose gradually their resemblance to those of the neck, and the lower ones come

nearer to the figure of the lumbar.

The articulation of the vertebræ of the back with the ribs, shall be more particularly considered after the ribs are described. Only it may be proper now to remark, that the ligaments which serve that articulation assist in

connecting the vertebræ.

The lowest order of the true vertebræ is the lumbar (g); which are five bones, that may be distinguished from any others by these marks: 1. Their bodies, tho' of a circular form at their fore-part, are somewhat oblong from one fide to the other; which may be occafioned by the pressure of the large vessels, the aorta and cava, and of the vifcera. The epiphyfes on their edges are larger; and therefore the upper and lower furfaces of their bodies are more concave than in the vertebræ of the back. 2. The cartilages between these vertebræ are much the thickest of any, and render the spine convex within the abdomen, by their greatest thickness being at their fore-part. 3. The oblique processes are strong and deep; those in opposite sides being almost placed in parallel planes; the superior, which are concave, facing inwards, and the convex inferior ones facing outwards: And therefore each of these vertebræ receives the one above it, and is received by the one below; which is not so evident in the other two classes already described. 4. Their transverse processes are finall, long, and almost erect, for allowing large motion to each bone, and fufficient infertion to muscles, and for supporting and defending the internal parts. 5. Betwixt the roots of the superior oblique and transverse processes, a small protuberance may be observed, where some of the muscles that raise the trunk of the body are inferted. 6. Their spinal proceffes are strong, straight, and horizontal, with broad

<sup>(</sup>g) Ocques igues, town, Renum, lumborum.

le

1-

ie

c

80

5,

in

12-

ed

0'

b-

2-

nd

d-

r-

r-

·fc

he

eft

16

es

r,

ex

ot

ei-

he

le

N-

n

r-

1e

e

ond

at

flat fides, and a narrow edge above and below; this last being depressed on each side by muscles. And at the root of these edges, we see rough surfaces for fixing the ligaments. 7. The canal for the numerous cords, called cauda equina, into which the spinal marrow divides, is rather larger in these bones than what contains that marrow in the vertebræ of the back.

8. The holes for the passage of the nerves are more equally formed out of both the contiguous vertebræ than in the other classes; the upper one surnishes, however, the larger share of each hole.

The thick cartilages between these lumbar vertebræ, their deep oblique processes, and their erect spinal processes, are all fit for allowing large motion: though it is not so great as what is performed in the neck; which appears from comparing the arches which the head describes when moving on the neck or the loins only.

The lumbar vertebræ, as they descend, have their oblique processes at a greater distance from each other, and facing more backwards and forwards.

Both transverse and spinal processes of the middlemost vertebræ of the loins are longest and thickest; in the vertebræ above and below, they are less: So that these processes of the first (b) and fifth (i) are the least, to prevent their striking on the ribs or offa ilium, or their bruising the muscles in the motions of the spine.

The epiphyses round the edges of the bodies of the lumbar vertebræ are most raised in the two lowest; which consequently make them appear hollower in the middle than the others are.

The body of the fifth vertebra is rather thinner than that of the fourth.—The spinal process of this fifth is smaller, and the oblique processes face more backwards and forwards than in any other lumbar vertebra.

AFTER confidering the structure of the particular

<sup>(</sup>h) Negerns, Renalis. (i) Anahirns, Fulciens.

Sa

n

0

th

2

a

vertebræ, and their mutual connection, we may obferve a folicitous care taken that they shall not be disjoined but with great difficulty. For, befides being connected by strong ligaments proportioned to the forces which are to be refisted, their bodies either enter fo into each other as to prevent their being displaced any way, as in the vertebræ of the neck; or they are propped on all fides, as those of the back are by the ribs; or their furaces of contact are so broad, as to render the feparation almost impracticable, as in the loins: while the depth and articulation of the oblique processes are exactly proportioned to the quantity of motion which the other parts of the bones allow, or the muscles can perform. Yet as these oblique processes are small, and therefore not capable of so secure a conjunction as the larger bodies, they may fooner yield to a disjoining force; but then their diflocation is not of near fo bad consequence as the separation of the bodies would be: for by the oblique processes being dislocated, the muscles, ligaments, and spinal marrow, are indeed stretched; but this marrow must be compressed, or entirely destroyed, when the body of the vertebra is removed out of its place.

## THE FALSE VERTEBRÆ.

The False Vertebræ compose the under pyramid of the spine. They are distinguished from the bones already described justly enough by this epithet of false; because, though each bone into which they can be divided in young people resembles the true vertebræ in figure, yet none of them contribute to the motion of the trunk of the body; they being intimately united to each other in adults, except at their lower part, where they are moveable; whence they are commonly divided into two bones, os sacrum and coccygis.

Os SACRUM (k), is so called, from being offered in facrifice

<sup>(</sup>k) Ιεζον, σπουδυλος μεγας, Hippocrat. υτοσπουδυλου, Oribaf. Πλατ., Latum, os clunium, clavium.

facrifice by the ancients, or rather because of its largeness in respect of the other vertebræ. This bone is of an irregular triangular shape, broad above, narrow below, convex behind, for the advantageous origin of the muscles that move the spine and thigh backwards; and concave behind, for enlarging the cavity of the pelvis.—Four transverse lines, of a colour different from the rest of the bone which are seen on its fore-part, are the marks of division of the five different bones of

which it confilts in young persons.

tT,

ob. lif-

n-

ces

in-

nv

ped

Or

the ile

are ich

can

ind

the ing

bad

e:

es, ed;

de-

out

nid

nes

le;

di-

in

of

to

ere led

in

ice 210

The fore-part of the os facrum, analogous to the bodies of the true vertebræ, is smooth and flat, to allow a larger space for the contained bowels, without any danger of hurting them; or this flat figure may be owing to the equal preffure of these bowels, particularly of the last gut .- The back-part of it is almost straight, without so large a cavity as the vertebræ have; because the spinal marrow, now separated into the cauda equina, is finall.—The bridges between the bodies and processes of this bone are much thicker, and in proportion shorter, than in the former class of bones.——The strength of these cross bridges is very remarkable in the three upper bones, and is well proportioned to the incumbent weight of the trunk of the body, which thefe bridges fustain in a transverse, consequently an unfavourable, fituation, when the body is erect.

There are only two oblique processes of the os facrum; one standing out on each fide from the upper part of the first bone.—Their plain erect surfaces face backwards, and are articulated with the inferior oblique processes of the last vertebra of the loins, to which each of these processes is connected by a strong ligament, which rifes from a fcabrous cavity round their roots, where mucilaginous glands are also lodged.—Instead of the other oblique processes of this bone, four rough tubercles are to be feen on each fide of its furface behind, from which the mulculus facer has its origin.

The transverse processes here are all grown together

into one large strong oblong process on each side; which, so far as it answers to the first three bones, is very thick, and divided into two irregular cavities by a long perpendicular ridge.—The foremost of the two cavities has commonly a thin cartilaginous skin covering it in the recent subject, and is adapted to the unequal protuberance of the os ilium; and a strong ligament connects the circumference of these surfaces of the two bones.—The cavity behind is divided by a transverse ridge into two, where strong ligamentous strings that go from this bone to the os ilium, with a cellular substance containing mucus, are lodged.

The transverse processes of the two last bones of the os facrum are much smaller than the former.——At their back-part near their edge, a knob and oblong flat surface give rise to two strong ligaments which are extended to the os ischium; and are therefore called

facrosciatic.

The spinal processes of the three uppermost bones of the os facrum appear short, sharp, and almost erect, while the two lower ones are open behind; and sometimes a little knob is to be seen on the fourth, though generally it is bisurcated, without the two legs meeting into a spine; in which condition also the first is often to be seen; and sometimes none of them meet, but leave a sinus, or rather sossa, instead of a canal (1).—The musculus latissimus and longissimus dorsi, facrolumbalis, and glutæus maximus, have part of their origins from these spinal processes.

The canal between the bodies and processes of this bone, for the cauda equina, is triangular; and becomes smaller as it descends, as the cauda also does.—Below the third bone, this passage is no more a complete bony canal, but is open behind; and is only there defended by a strong ligamentous membrane stretched over it, which, with the muscles that cover it, and are

very

<sup>(1)</sup> Verheyen, Anat. tract. 5. cap. 9. - Sue Trad. d'ofteol. p. 127.

rt I.

de;

, is

y a

WO

ing

ual

ent

wo

hat

ıb-

the

At

ng

ed

of

ct,

e-

gh

ng

en

ut

) i -

is

d

very prominent on each fide, is a fufficient defence for the bundle of nerves within.

At the root of each oblique process of this bone, the notch is conspicuous; by which, and such another in the last vertebra of the loins, a passage is left for the twenty-fourth spinal nerve; and in viewing the os facrum, either before or behind, four large holes appear in each fide, in much the fame height as where the marks of the union of its several bones remain. of the largest nerves of the body pass through the anterior holes; and superficial grooves, running outwards from them in different directions, show the course of these nerves.——From the intervals of these grooves, the pyriformis muscle chiefly rises.——The holes in the back-part of the bone are covered by membranes which allow finall nerves to pass through them. The two uppermost of these holes, especially on the fore-side, are the largest; and as the bone descends, the holes turn Sometimes a notch is only formed at the lower part in each fide of this bone; and in other fubjects there is a hole common to it and the os coccygis, through which the twenty-ninth pair of spinal nerves passes; and frequently a bony bridge is formed on the back-part of each fide by a process sent up from the back-part of the os coccygis, and joined to the little knobs which the last bone of the os facrum has instead of a spinal process. Under this bridge or jugum, the twenty-ninth pair of spinal nerves runs in its course to the common holes just now described.

The upper part of the body of the first bone refembles the vertebræ of the loins; but the small fifth bone is oblong transversely, and hollow in the middle of its lower surface.

The fubstance of the os facrum is very fpongy, without any confiderable folid external plates, and is lighter proportionally to its bulk than any other bone in the body; but is fecured from injuries by the thick muscles that cover it behind, and by the strong ligamentous

L 4

mem-

membranes that closely adhere to it.—As this is one of the most remarkable instances of this sort of defence efforded a soft weak bone, we may make the general observation, That wherever we meet with such a bone, one or other, or both these defences, are made use of; the first to ward off injuries, and the second to keep the substance of the bone from yielding too easily.

This bone is articulated above to the last vertebra of the loins, in the manner that the lumbar vertebræ are joined; and therefore the fame motions may be performed here.——The articulation of the lower part of the os facrum to the os coccygis, feems well enough adapted for allowing confiderable motion to this last bone, were it not much confined by ligaments. Laterally, the os facrum is joined to the offa ilium by an immoveable fynchondrosis, or what almost deserves the name of a future: For the cartilaginous crust on the furface of the bones is very thin; and both their furfaces are so scabrous and unequal, as to be indented into each other; which makes fuch a strong connection, that great force is required to separate them, after all the muscles and ligaments are cut.—Frequently the two bones grow together in old subjects.

The uses of the os sacrum are, to serve as the common base and support of the trunk of the body, to guard the nerves proceeding from the end of the spinal marrow, to defend the back-part of the pelvis, and to afford sufficient origin to the muscles which move the

trunk and thigh.

The bones that compose the os facrum of infants, have their bodies separated from each other by a thick cartilage; and, in the same manner as the true vertebræ, each of them consists of a body and two lateral plates, connected together by cartilages; the ends of the plates seldom being contiguous behind.

Os Coccygis (m), or rump-bone, is that triangular chain

<sup>(</sup>m) Oggowuyiov ogos, Caudæ os, spondylium, os cuculi.

chain of bones depending from the os facrum; each bone becoming smaller as they descend, till the last ends almost in a point. The os coccygis is convex behind, and concave before; from which crooked pyramidal figure, which was thought to resemble a cuckow's beak, it has got its name.

This bone consists of four pieces in people of middle age:—In children, very near the whole of it is cartilage: In old subjects, all the bones are united, and become frequently one continued bone with the os sa-

crum.

The highest of the four bones is the largest, with shoulders extended farther to each side than the end of the os facrum; which enlargement should, in my opinion, ferve as a diffinguishing mark to fix the limits of either bone; and therefore should take away all difpute about reckoning the number of bones, of which one or other of these two parts of the false vertebræ is composed; which dispute must still be kept up, so long as the numbering five or fix bones in the os facrum depends upon the uncertain accident of this broad shouldered little bone being united to or separated from it. The upper furface of this bone is a little hollow.— From the back of that bulbous part called its shoulders, a process often rises up on each side, to join with the bifurcated spine of the fourth and fifth bones of the os facrum, to form the bony bridge mentioned in the defcription of the os facrum.—Sometimes these shoulders are joined to the fides of the fifth bone of the os facrum, to form the hole in each fide common to these two bones, for the passage of the twenty-ninth pair of ipinal nerves.—Immediately below the shoulders of the os coccygis, a notch may be remarked in each fide, where the thirtieth pair of the spinal nerves passes.— The lower end of this bone is formed into a finall head, which very often is hollow in the middle.

The three lower bones gradually become smaller, and are spongy; but are strengthened by a strong ligament

which

C

ne

to

W

gr

fh

of

th

in

m

in

W

h

tu

al

al

m

0

0

to

0

a

which covers and connects them.—Their ends, by which they are articulated, are formed in the fame manner as those of the first bone are.

Between each of these four bones of young subjects a cartilage is interposed; therefore their articulation is analogous to that of the bodies of the vertebræ of the neck: For, as has been above remarked, the lower end of the os facrum, and of each of the three superior bones of the os coccygis, has a small depression in the middle; and the upper part of all the bones of the os occygis is a little concave, and confequently the interposed cartilages are thickest in the middle, to fill up both cavities; by which they connect the bones more firmly.—When the cartilages offify, the upper end of each bone is formed into a cavity, exactly adapted to the protuberant lower end of the bone immediately above. From this fort of articulation, it is evident, that, unless when these bones grow together, all of them are capable of motion; of which the first and fecond, especially this last, enjoys the largest share.

The lower end of the fourth bone terminates in a

rough point, to which a cartilage is appended.

To the fides of these bones of the os coccygis, the coccygai muscles (n), and part of the levatores ani,

and of the glutæi maximi, are fixed.

The *fubstance* of these bones is very spongy, and in children cartilaginous; there being only a part of the first bone offissed in a new-born infant.—Since therefore the intestinum rectum of children is not so firmly supported as it is in adults, this may be one reason why they are more subject to a procidentia ani than old people (0).

From the description of this bone, we see how little it resembles the vertebræ; since it seldom has processes,

(n) Douglas, Myograph. chap. 40. — Eustach. tab. 36. n° 45.

(0) Spigel. de humani corp. fabric. lib. 2. cap. 22.—Paaw de offib. par. 2. cap. 3.

t I,

by

me

Cts

18

he

ver

lor

he

OS

er-

up

of

to

ely

nt, of

nd

a

he ni,

in

he

e-

ly hy

0-

it

S,

er

5,

de

never has any cavity for the spinal marrow, nor holes for the passage of nerves.—Its connection hinders it from being moved to either side; and its motion backwards and forwards is much confined: yet, as its ligaments can be stretched by a considerable force, it is a great advantage in the execretion of the secess alvinæ, and much more in child-bearing, that this bone should remain moveable; and the right management of it, in delivering women, may be of great benefit to them (p)—The mobility of the os coccygis diminishing as people advance in age, especially when its ligaments and cartilages have not been kept slexible by being stretched, is probably one reason why the women, who are old maids before they marry, have generally hard labour in child-bed.

The os coccygis ferves to fustain the intestinum rectum; and, in order to perform this office more effectually, it is made to turn with a curve forwards; by which also the bone itself, as well as the muscles and teguments, is preserved from any injury when we sit with our body reclined back.

## \$ 2. Of the PELVIS.

THE fecond part of the trunk of the skeleton, the pelvis, is the cylindrical cavity at the lower part of the abdomen, formed by the os facrum, os coccygis, and ossa innominata; which last therefore fall now in course to be examined.

## OSSA INNOMINATA.

Though the name of offa innominata (q) contributes nothing to the knowledge of their fituation, structure, or office; yet they have been so long universally known by it, that there is no occasion for changing it.—They are two large broad bones, which form the fore-part and sides of the pelvis, and the lower part of the sides

(9) Exerews, προσφυσεις, Sacro conjuncta.

<sup>(</sup>p) Paaw, ibid.—Deventer, Operat. chirurg. cap. 27.

C

b

fi:

V

lo

tl

b

n

C

i

tl

a

t

p

of the abdomen.—In children, each of these bones is evidently divided into three; which are afterwards so intimately united, that scarce the least mark of their former separation remains: This notwithstanding, they are described as consisting each of three bones, to wit, the os ilium, ischium, and pubis; which I shall first describe separately, and then shall consider what is common to any two of them, or to all the three.

Os ILIUM (r), or haunch-bone, is fituated highest of the three, and reaches as far down as one third of the great cavity into which the head of the thigh-bone is received.

The external fide of this bone is unequally convex, and is called its dorfum;—the internal concave furface is by some (but improperly) named its costa. ——The femicircular edge at the highest part of this bone, which is tipped with a cartilage in the recent subject, is named the *(pine, into which the external or descending* oblique muscle of the abdomen is inserted; and from it the internal ascending oblique, and the transverse muscles of the belly, with the glutæus maximus, quadratus lumborum, and latiffimus dorfi, have their origin. Some (s) are of opinion, that it is only the tendinous crust of all these muscles, and not a cartilage, as commonly alleged, that covers this bony edge.—The ends of the spine are more prominent than the surface of the bone below them; therefore are reckoned proceffes.—From the anterior spinal process, the fartorius and facialis muscles have their rife, and the outer end of the doubled tendon of the external oblique muscle of the abdomen, commonly called Falliopus's or Poupart's ligament, is fixed to it. - The infide of the posterior spinal process, and of part of the spine forward from that, is made flat and rough where the facro-lum-

<sup>(</sup>r) Λαγονον, χενεων, Scaphium, lumbare, clunium, clavium, anchas.
(s) Winflow, Exposition anatomique du corps humain, traité des es frais, § 96.

tI.

les

fo

eir

ey

it,

m-

of

he

is

ex,

he

ich

na-

ng

om

rfe

in.

us

m.

he

ice

0.

to-

ter ue

or

00-

ird

m·

lis

as. des balis and longissimus dorsi rise; and to its outside are fixed ligaments, extended to the os facrum and tranfverse processes of the fifth and fourth vertebræ of the loins.(t)—Below the anterior spinal process another protuberance stands out, which by its situation may be diftinguished from the former, by adding the epithet of inferior, where the musculus rectus tibiæ has its origin (u).—Betwixt these two anterior processes the bone is hollowed where the beginning of the fartorius muscle is lodged.—Below the posterior spinal procefs, a fecond protuberance of the edge of this bone is in like manner observable, which is closely applied to the os facrum.—Under this last process a considerable large niche is observable in the os ilium; between the fides of which and the strong ligament that is stretched over from the os facrum to the sharp-pointed process of the os ischium of the recent subject, a large hole is formed, through which the musculus pyriformis, the great sciatic nerve, and the posterior crural vessels, pass, and are protected from compression.

The external broad fide or dorfum of the os ilium is a little hollow towards the fore-part; farther back, it is as much raised; then is considerably concave; and, These inequalities are occasioned lastly, it is convex. by the actions of the muscles that are situated on this furface.—From behind the uppermost of the two anterior spinal processes, in such bones as are strongly marked by the muscles, a semicircular ridge is extended to the hollow passage of the sciatic nerve. Between the spine and this ridge, the glutæus medius takes its rife. Immediately from above the lowest of the anterior spinal processes, a second ridge is stretched to the niche. Between this and the former ridge, the glutæus minimus has its origin.—On the outfide of the posterior ipinal processes, the dorsum of the os ilium is flat and rough, where part of the musculus glutæus maximus

<sup>(</sup>t) Weitbrecht, Sydesmolog. sect. 4. § 39, 40, 46, 47-(u) Baker, Curs. Osteolog. demonstr. 3.

and pyriformis rifes.—The lowest part of this bone is the thickest, and is formed into a large cavity with high brims, to affist in composing the great acetabulum; which shall be considered, after all the three bones that constitute the os innominatum are described.

The internal furface of the os ilium is concave in its broadest fore-part, where the internal iliac muscle has its origin, and some share of the intestinum ilium and colon is lodged.—From this large hollow, a small sinuosity is continued obliquely forwards, at the inside of the anterior inferior spinal process, where part of the ploas and iliacus muscles, with the crural vessels and nerves, pass.—The large concavity is bounded below by a sharp ridge, which runs from behind forwards; and, being continued with such another ridge of the os pubis, forms a line of partition between the abdomen and pelvis.—Into this ridge the broad tendon of the

pfoas parvus is inferted.

All the internal furface of the os ilium, behind this ridge, is very unequal: For the upper part is flat, but fpongy, where the facro-lumbalis and longiffimus dorfi rife.—Lower down, there is a transverse ridge from which ligaments go out to the os facrum.—Immediately below this ridge, the rough unequal cavities and prominences are placed, which are exactly adapted to those described on the fide of the os facrum.—In the fame manner, the upper part of this rough furface is porous, for the firmer adhesion of the ligamentous cellular fubstance; while the lower part is more solid, and covered with a thin cartilaginous skin, for its immoveable articulation with the os facrum.—From all the circumterence of this large unequal furface, ligaments are extended to the os facrum, to fecure more firmly the conjunction of these bones.

The passages of the medullary vessels are very conspicuous, both in the dorsum and costa of many ossa

ilium; but in others they are inconsiderable.

The posterior and lower parts of these bones are thick;

thic con the and cep

Cha

gine of a

fitte figures line for that

leg

fid

pu

bet

profro

Be it po of of

w pl

in

t!

thick; but they are generally exceedingly thin and compact at their middle, where they are exposed to the actions of the musculi glutæi and iliacus internus, and to the pressure of the bowels contained in the belly.

—The substance of the offa ilium is mostly cellular, except a thin external table.

In a ripe child, the spine of the os ilium is cartilaginous; and is afterwards joined to the bone, in form of an epiphyse.—The large lower end of this bone is not

completely offified.

Os Ischium (x), or hip-bone, is of a middle bulk between the two other parts of the os innominatum, is fituated lowest of the three, and is of a very irregular figure.—Its extent might be marked by an horizontal line drawn near through the middle of the acetabulum; for the upper bulbous part of this bone forms some less than the lower half of that great cavity, and the small leg of it rises to much the same height on the other side of the great hole common to this bone and the os

pubis.

tÎ.

is

gh

n;

at

its

as

nd

all

de he

nd

W

3;

os

en

10

is

ut

fi

n

y

)-

0

e

S

d

9

From the upper thick part of the os ischium, a sharp process, called by some spinous, stands out backwards, from which chiefly the musculus coccygæus and superior gemellus, and part of the levator ani, rife; and the anterior or internal facrofciatic ligament is fixed to it.— Between the upper part of this ligament and the bones, it was formerly observed that the pyriform muscle, the polterior crural vessels, and the sciatic nerve, pals out of the pelvis.—Immediately below this process, a finuolity is formed for the tendon of the musculus obturator internus.——In a recent subject, this part of the bone, which ferves as a pulley on which the obturator mulcle plays, is covered with a ligamentous cartilage, that, by two or three small ridges, points out the interstices of the fibres in the tendon of this muscle.——The outer surface of the bone at the root of this spinous process

bı

re

er

th

bo

to

la

ei

fo

th

tl

0

tl

t

is made hollow by the pyriformis or iliacus externus muscle.

Below the finuofity for the obturator muscle, is the great knob or tuberofity, covered with cartilage or tendon (y).—The upper part of the tuberofity gives rife to the inferior gemellus muscle.—To a ridge at the infide of this, the external or posterior facrosciatic ligament is fo fixed, that between it, the internal ligament, and the finuofity of the os ischium, a passage is left for the internal obturator muscle — The upper thick fmooth part of the tuber, called by some its dorfum, has two oblique impressions on it. The inner one gives origin to the long head of the biceps, flexor tibiæ, and feminervofus muscles; and the semimembranosus rises from the exterior one, which reaches higher and nearer the acetabulum than the other.—The lower, thinner, more scabrous part of the knob which bends forwards, is also marked with two flat surfaces; whereof the internal is what we lean upon in fitting, and the external gives rife to the largest head of the triceps adductor femoris.—Between the external margin of the tuberofity and the great hole of the os innominatum, there is frequently an obtuse ridge extended down from the acetabulum, which gives origin to the quadratus femoris. -As the tuber advances forwards, it becomes smaller, and is rough, for the origin of the musculus transversalis and erector penis.—The small leg of it, which mounts upwards to join the os pubis, is rough and prominent at its edge, where the two lower heads of the triceps or quadriceps adductor femoris take their rife.

The upper and back part of the os ischium is broad and thick; but its lower and fore part is narrower and thinner.—Its substance is of the structure common to

broad bones.

The os ilium and pubis of the same side are the only bones which are contiguous to the os ischium.

The part of the os ischium, which forms the acetabulum,

<sup>(</sup>y) Winflow, Exposit. Anat. de os frais, \$ 96.

İ.

lus

he

ile

in-

ra-

nt, for

ck

las

res nd

fes

er

er,

is,

n-

nal

e-

ty e-

e-

IS.

r,

a-

ts

nt

or

id

d

0

\*

2-

bulum, the spinous process, the great tuber, and the recurved leg, are all cartilaginous at birth.—The tuber, with part of the leg or process above it, becomes an epiphyse before this bone is fully formed.

The Os Pubis (z), or share-bone, is the least of the three parts of the os innominatum, and is placed at the upper fore-part of it.—The thick largest part of this bone is employed in forming the acetabulum; from which becoming much finaller, it is stretched inwards to its fellow of the other fide, where again it grows larger, and fends a finall branch downwards to join the end of the small leg of the os ischium.—The upper fore-part of each os pubis is tuberous and rough where the musculus rectus and pyramidalis are inserted.— From this a ridge is extended along the upper edge of the bone, in a continued line with such another of the os ilium, which divides the abdomen and pelvis. The ligament of Fallopius is fixed to the internal end of this ridge, and the fmooth hollow below it is made by the ploas and iliacus internus muscles passing with the anterior crural vessels and nerves behind the ligament. -Some way below the former ridge, another is extended from the tuberous part of the os pubis downwards and outwards towards the acetabulum; between thele two ridges the bone is hollow and smooth, for lodging the head of the pectineus muscle.—Immediately below, where the lower ridge is to take the turn downwards, a winding niche is made, which is comprehended in the great foramen of a skeleton; but is formed into a hole by a fubtended ligament in the recent subject, for the passage of the posterior crural nerve, an artery, and a vein.—The internal end of the os pubis is rough and unequal, for the firmer adhesion of the thick ligamentous cartilage that connects it to its fellow of the other fide: - The process which goes down from that to the os ischium is broad and rough

<sup>(</sup>z) nons, Pectinis, penis, pudibundum, fenestratum.

b

0

1

d

i

ti

tl

la

tl

tl

e

W

p

10

tl

it

tl

liga-

before, where the gracilis and upper heads of the triceps, or rather quadriceps, adductor femoris have their origin.

The fubstance of the os pubis is the same as of other

broad bones.

Only a part of the large end of this bone is offified, and the whole leg is cartilaginous, in a child born at the full time.

Betwixt the os ischium and pubis a very large irregular hole is left, which, from its refemblance to a door or shield, has been called thyroides. This hole is all, except the nitch for the posterior crural nerve, filled up, in a recent subject, with a strong ligamentous membrane, that adheres very firmly to its circumference. From this membrane chiefly the two obturator muscles, external and internal, take their rife.—The great defign of this hole, befides rendering the bone lighter, is to allow a strong enough origin to the obturator muscles, and sufficient space for lodging their bellies, that there may be no danger of disturbing the functions of the contained vifcera of the pelvis by the actions of the internal, nor of the external being bruifed by the thigh-bone, especially by its lesser trochanter, in the motions of the thigh inwards: Both which inconveniences must have happened, had the offa innominata been complete here, and of fufficient thickness and strength to ferve as the fixed point of these muscles.-The bowels fometimes make their way through the nitch for the veffels, at the upper part of this thyroid hole; and this causes a hernia in this place (a).

In the external furface of the offa innominata, near the outfide of the great hole, a large deep cavity is formed by all the three bones conjunctly: For the os pubis constitutes about one fifth, the os ilium makes fomething less than two fifths, and the os ischium as much more than two fifths. The brims of this cavity are very high, and are still much more enlarged by the

<sup>(</sup>a) Memoires de l'acad. de chirurgie, tom. 1. p. 709, &c.

1-

ir

er

d,

at

e.

a

is

ed

m-

e.

es,

le-

13

10

es,

ns

ot

he

he

re-

ita

nd

ch

e;

ear

TS

OS

es

as.

ity

he ga-

ligamentous cartilage, with which they are tipped in a recent subject. From this form of the cavity it has been called acetabulum; and for a diltinguishing character, the name of the bone that constitutes the largest share of it is added; therefore acetabulum offis ischii (b) is the name this cavity commonly bears.—Round the base of the supercilia the bone is rough and unequal, where the capfular ligament of the articulation is fixed. —The brims at the upper and back part of the acetabulum are much larger and higher than any where elfe; which is very necessary to prevent the head of the femur from flipping out of its cavity at this place, where the whole weight of the body bears upon it, and confequently would otherwise be constantly in danger of thrusting it out.—As these brims are extended downwards and forwards, they become lefs; and at their internal lower part a breach is made in them; from the one fide of which to the other, a ligament is placed in the recent subject; under which a large hole is left, which contains a fatty cellular fubstance and vessels. The reason of which appearance has afforded matter of To me it feems evidently contrived for allowing a larger motion to the thigh inwards: For if the bony brims had been here continued, the neck of the thigh-bone must have struck upon them when the thighs were brought across each other; which, in a large itrong motion this way, would have endangered the neck of the one bone, or brim of the other. Then the veffels which are distributed to the joint may fafely enter at the finuofity in the bottom of the breach; which being, however, larger than is necessary for that purpose, allows the larger mucilaginous gland of the joint to escape below the ligament, when the head of the thigh-bone is in hazard of prefling too much upon it in the motions of the thigh outwards (c).—Belides this difference in the height of the brims, the acetabulum M 2

<sup>(</sup>b) Coxæ, coxendicis.

<sup>(</sup>c) Petit. Memoires de l'acad. des sciences, 1722.

1

n

1

1

lum is otherwise unequal: for the lower internal part of it is depressed below the cartilaginous surface of the upper part, and is not covered with cartilage; into the upper part of this particular depression, where it is deepest and of a semilunar form, the ligament of the thigh-bone, commonly, though improperly, called the round one, is inserted; while, in its more superficial lower part, the large mucilaginous gland of this joint is lodged. The largest share of this separate depression is formed in the os ischium.

From what has been faid of the condition of the three bones composing this acetabulum in new-born children, it must be evident that a considerable part of this cavity is cartilaginous in them.

THE offa innominata are joined at their back-part to each fide of the os facrum by a fort of future, with a very thin intervening cartilage, which ferves as fo much glue to cement those bones together; and strong ligaments go from the circumference of this unequal furface, to connect them more firmly. The offa innominata are connected together at their fore-part by the ligamentous cartilage interposed between the two offa pubis.—These bones can therefore have no motion in a natural state, except what is common to the trunk of the body, or to the os facrum. But it has been difputed, whether or not they loofen so much from each other, and from the os facrum, in child-birth, by the flow of mucus to the pelvis, and by the throes of labour, as that the offa pubis recede from each other, and thereby allow the paffage between the bones to be enlarged.—Several observations (d) show that this relaxation fometimes happens: But those who had trequently opportunities of diffecting the bodies of women who died immediately after being delivered of chil-

<sup>(</sup>d) Bauhin. Theat. anat. lib. 1. cap. 49.—Spigel. Anat. lib. 2. cap. 24.—Riolan. Anthropog. lib. 6. cap. 12.—Diemer. broek, Anat. lib. 9. cap. 16.

T.

of ne

10

is

ne

ne

al

18

13

ne

rn

ut

to

a

ar-

1-

ne

la

in

of

uh

ic

a.

r,

is

e-

n |-

1,

1.

dren, teach us to beware of regarding this as the common effect of child-birth; for they found such a relaxation in very few of the bodies which they examined (e).

Considering what great weight is supported in our erect posture, by the articulation of the ossa innominata with the ossacrum, there is great reason to think, that if the conglutinated surfaces of these bones were once separated, (without which the ossa pubis cannot shuffle on each other), the ligaments would be violently stretched, if not torn; from whence many disorders would arise (f).

Each os innominatum affords a focket (the acetabulum) for the thigh-bones to move in; and the trunk of the body rolls here so much on the heads of the thighbones, as to allow the most conspicuous motions of the trunk, which are commonly thought to be performed by the bones of the spine.—This articulation is to be more fully described after the ossa femoris are examined.

The pelvis, then, has a large open above where it is continued with the abdomen; is strongly senced by bones on the sides, back and fore part; and oppears with a wide opening below, in the skeleton; but, in recent subject, a considerable part of the opening is silled by the sacrosciatic ligaments, pyriform, internal obturator, levatores ani, gemini, and coccygæi muscles, which support and protect the contained parts better than bones could have done; so that space is only left at the lowest part of it, for the large excretories, the vesica urinaria, intestinum rectum, and, in semales, the uterus, to discharge themselves.

M 3

§ 3. Bones

(f) Ludov. in Ephem. German. dec. 1. ann. 3. obf. 255.

<sup>(</sup>e) Hildan. epist. cent. obs. 46.—Dionis Sixieme demonst. des.—Morgagn. Advers. 3. animad. 15.

W

n

ri

te

2

W

0

t

n

## § 3. Bones of the THORAX.

The Thorax (g), or chest, which is the only part of the trunk of the body which we have not yet described, reaches from below the neck to the belly; and, by means of the bones that guard it, is formed into a large cavity: The figure of which is somewhat conoidal: but its upper smaller end is not finished, being left open for the passage of the wind-pipe, gullet, and large blood-vesses; and its lower part, or base, has no bones, and is shorter before than behind; so that, to carry on our comparison, it appears like an oblique section of the conoid. Besides which, we ought also to remark, that the lower part of this cavity is narrower than some way above (b); and that the middle of its back-part is considerably diminished by the bones standing forwards into it.

The bones which form the thorax are the twelve dorfal vertebræ behind, the ribs on the fides, and the

sternum before.

The vertebræ have already been described as part of the spine; and therefore are now to be passed.

### THE RIBS.

THE ribs, or costa (i), (as if they were custodes, or guards, to those principal organs of the animal machine, the heart and lungs), are the long crooked bones placed at the side of the chest, in an oblique direction downwards in respect of the back-bone.—Their number is generally twelve on each side; though frequently eleven or thirteen have been found (k).——Sometimes the ribs are found preternaturally conjoined or divided (1).

The

(1) Sue trad. d'osteolog. p. 141.

<sup>(</sup>g) Pectus, cassum. (b) Albin. de ossib. § 169.

<sup>(</sup>i) Πλευραι, περιστερία, σπαθαι.
(k) Riolan. Comment. de offibus, cap. 19.—Marchetti, cap. 9.
Cowper Explicat. tab. 93. and 94.—Morgagn. Advers. anat.

I.

art

le-

y ;

ed

at

e-

et,

e,

fo an

ht

r-

lle

es

ve

ne

of

or

a-

es

n

ir

d

C

The ribs are all concave internally; where they are also made smooth by the action of the contained parts, which, on this account, are in no danger of being hurt by them; and they are convex externally, that they might refift that part of the preffure of the atmosphere which is not balanced by the air within the lungs during in/piration.—— The ends of the ribs next the vertebræ are rounder than they are after these bones have advanced forwards, when they become flatter and broader, and have an upper and lower edge, each of which is made rough by the action of the intercostal muscles inserted into them. These muscles, being all of nearly equal force, and equally stretched in the interstices of the ribs, prevent the broken ends of these bones, in a fracture, from being removed far out of their natural place, to interrupt the motion of the vital organs.—The upper edge of the ribs is more obtule and rounder than the lower, which is depressed on its internal fide by a long fossa, for lodging the intercostal veffels and nerves; on each fide of which there is a ridge, to which the intercostal muscle's are fixed. The fossa is not observable, however, at either end of the ribs; for, at the posterior or root, the vessels have not yet reached the ribs; and, at the fore-end, they are split away into branches, to ferve the parts between the ribs: Which plainly teaches furgeons one reason of the greater fafety of performing the operation of the empyema towards the fides of the thorax, than either near the back or the breaft.

At the posterior end (m) of each rib, a little head is formed, which is divided by a middle ridge into two plain or hollow surfaces; the lowest of which is the broadest and deepest in most of them. The two plains are joined to the bodies of two different veriebræ, and the ridge forces itself into the intervening cartilage.—

A little way from this head, we find, on the external surface, a small cavity, where mucilaginous glands are lodged;

CC

th

CC

el

28

p

fu

r

fi

lodged; and round the head, the bone appears fpongy, where the capfular ligament of the articulation is fixed.—Immediately beyond this a flatted tubercle rifes, with a finall cavity at and roughness about its root, for the articulation of the rib with the transverse process of the lowest of the two vertebræ, with the bodies of which the head of the rib is joined.—Advancing farther on this external furface, we observe in most of the ribs another smaller tubercle, into which ligaments which connect the ribs to each other, and to the transverse processes of the vertebræ and portions of the longissimus dorsi, are inserted.—Beyond this the ribs are made flat by the facro-lumbalis muscle, which is inferted into the part of this flat furface farthest from the spine, where each rib makes a considerable curve, called by some its angle.—Then the rib begins to turn broad, and continues fo to its anterior end (n), which is hollow and fpongy, for the reception of, and firm coalition with, the cartilage that runs thence to be inferted into the sternum, or to be joined with some other cartilage.—In adults, generally the cavity at this end of the ribs is smooth and polished on its surface; by which the articulation of the cartilage with it has the appearance of being defigned for motion, but it has none.

The fubstance of the ribs is spongy, cellular, and only covered with a very thin external lamellated surface, which increases in thickness and strength as it approaches the vertebræ.

To the fore-end of each rib a long broad and strong cartilage is fixed, and reaches thence to the sternum, or is joined to the cartilage of the next rib. This course, however, is not in a straight line with the rib; for generally the cartilages make a considerable curve, the concave part of which is upwards; therefore, at their insertion into the sternum, they make an obtuse angle above, and an acute one below.—These cartila-

I.

n.

is

its

fe

0-

nin

li-

to of

ne

ch

m

e,

rn

m

ner

nd Dy

1e

as

y

25

is

t

C

-

ges are of such a length as never to allow the ribs to come to a right angle with the spine; but they keep them situated so obliquely as to make them angle very considerably obtuse above, till a force exceeding the elasticity of the cartilage is applied.—These cartilages, as all others, are firmer and harder internally than they are on their external surface; and sometimes, in old people, all their middle substance becomes bony, while a thin cartilaginous lamella appears externally (o). The offisication, however, begins frequently at the external surface.—The greatest alternate motions of the cartilages being made at their great curvature, that part remains frequently cartilaginous, after all the rest is of-sisted (p).

The ribs then are articulated at each end, of which the one behind is doubly joined to the vertebræ; for the head is received into the cavities of two bodies of the vertebræ, and the larger tubercle is received into the depression in the transverse process of the lower vertebra.—When one examines the double articulation, he must immediately see, that no other motion can here be allowed than upwards and downwards; fince the transverse process hinders the rib to be thrust back; the relistance on the other fide of the sternum prevents the ribs coming forward; and each of the two joints, with the other parts attached, oppose its turning round. But then it is likewise as evident, that even the motion upwards and downwards can be but small in any one rib at the articulation itself. But as the ribs advance forwards, the distance from their centre of motion increafing, the motion must be larger; and it would be very confpicuous at their anterior ends, were not they relisted there by the cartilages, which yield so little, that the principal motion is performed by the middle part of the ribs, which turns outwards and upwards, and occasions the twist remarkable in the long ribs

<sup>(0)</sup> Vefal. lib. 2. cap. 19.

p) Havers, Ofteolog. Nov. dife. 5. p. 289.

creases

g

n

h

S

at the place near their fore-end where they are most re-

fifted (q).

Hitherto I have laid down the structure and connection which most of the ribs enjoy, as belonging to all of them; but must now consider the specialties wherein any of them differ from the general description gi-

ven, or from each other.

In viewing the ribs from above downwards, their figure is still straighter; the uppermost being the most crooked of any.—Their obliquity in respect of the fpine increases as they descend; so that though their distances from each other is very little different at their back-part, yet at their fore-ends the diltances between the lower ones must increase.—In consequence, too, of this increased obliquity of the lower ribs, each of their cartilages makes a greater curve in its progress from the rib towards the sternum; and the tubercles, that are articulated to the transverse processes of the vertebræ, have their smooth surfaces gradually facing more upwards.—The ribs becoming thus more oblique, while the sternum advances forwards in its descent, makes the distance between the sternum and the anterior end of the lower ribs greater than between the sternum and the ribs above; consequently the cartilages of those ribs that are joined to the breast-bone are longer in the lower than in the higher ones. - These cartilages are placed nearer to each other as the ribs descend, which occasions the curvature of the cartilages to be greater.

The length of the ribs increases from the first and uppermost rib, as far down as the seventh; and from that to the twelfth, as gradually diminishes.—The superior of the two plain, or rather hollow surfaces, by which the ribs are articulated to the bodies of the vertebræ, gradually increases from the first to the sourth rib, and is diminished after that in each lower rib.—The distance of their angles from the heads always in-

<sup>(9)</sup> Winslow Memoires de l'acad. des sciences, 1720.

creases as they descend to the ninth, because of the greater breadth of the facrolumbalis muscle (r).

The ribs are commonly divided into true and false.

The true (s) costæ are the seven upper ones of each side, whose cartilages are all gradually longer as the ribs descend, and are joined to the breast-bone: so that, being pressed constantly between two bones, they are slatted at both ends; and are thicker, harder, and more liable to ossify, than the other cartilages that are not subject to so much pressure. These ribs include the heart and lungs; and therefore are the proper or true

custodes of life.

tI.

re-

ec-

all

re-

gi-

eir oft

he

di-

eir

en of

eir

m

te-

re

ile

nd

nd

bs

he

re

ch

nd

m

e-

Dy

r-

th

nes

The five inferior ribs of each fide are the false or baflard (t), whose cartilages do not reach to the sternum; and therefore, wanting the refiltance at their fore-part, they are there pointed; and on this account, having less pressure, their substance is softer.—The cartilages of these false ribs are shorter as the ribs descend.—To all thele five ribs the circular edge of the diaphragm is connected; and its fibres, instead of being stretched immediately transversely, and so running perpendicular to the ribs, are pressed so as to be often, especially in exspiration, parallel to the plane in which the ribs lie: Nay, one may judge by the attachments which these fibres have fo frequently to the fides of the thorax a confiderable way above where their extremities are inferted into the ribs, and by the fituation of the viscera always to be observed in a dead subject laid supine, that there is constantly a large concavity formed on each fide by the diaphragm within these bastard-ribs, in which the stomach, liver, spleen, &c. are contained; which being only reckoned among the vifcera naturalia, have occasioned the name of bastard custodes to these bones.

Hence

(s) rmoat Germanæ, legitimæ.

<sup>(</sup>r) Winflow, Exposition anatomique des os secs, § 643.

<sup>(</sup>t) Μαλθοκαι, χουδροδεις, ακανθαι, κίενες, ροαι, Adulterinæ, spuriæ, illegitimæ.

ti

th

of

as

wis

h

tl 'I

b

t

Hence, in simple fractures of the false ribs, without fever, the stomach ought to be kept moderately filled with food, lest the pendulous ribs falling inwards, should thereby increase the pain, cough, &c. (u).—
Hence likewise we may learn how to judge better of the seat of several diseases, and to do the operation of the empyema, and some others, with more safety than we can do if we follow the common directions.

The eight upper ribs were formerly (x) classed into pairs, with particular names to each two, to wit, the crooked, the folid, the pectoral, the twisted: But these names are of so little use, that they are now generally

neglected.

The first rib of each side is so situated, that the flat fides are above and below, while one edge is placed inwards and the other outwards, or nearly fo; therefore sufficient space is left above it for the subclavian veffels and muscle; and the broad concave surface is opposed to the lungs: But then, in consequence of this fituation, the channel for the intercostal vessels is not to be found; and the edges are differently formed from all the other, except the second; the lower one being rounded, and the other sharp.——The head of this rib is not divided into two plain furfaces by a middle ridge, because it is only articulated with the first vertebra of the thorax.——Its cartilage is offified in adults, and is united to the sternum at right angles.—Frequently this first rib has a ridge rising near the middle of its posterior edge, where one of the heads of the scalenus muscle rifes.-Farther forward it is flatted, or sometimes depressed by the clavicle.

The fifth, fixth, and seventh, or rather the fixth, seventh, eighth, and sometimes the fifth, fixth, seventh, eighth, ninth ribs, have their cartilages at least contiguous; and frequently they are joined to each

other

<sup>(</sup>u) Hippocrat. de articulo, § 51.—Paré, lib. 15. cap. 11. (x) Laurent. Hift. Anat. lib. 2. cap. 29.—Paaw, de offibus, pars 3. cap. 2.

tI.

ut

ed ls,

of

of

an

to he

le

at

ed e-

is

is

ot

m

g

of

is

y

S

S

.

1

r

other by cross cartilages; and most commonly the cartilages of the eighth, ninth, tenth, are connected to the former and to each other by firm ligaments.

The eleventh, and fometimes the tenth rib, has no tubercle for its articulation with the transverse process of the vertebra, to which it is only loosely fixed by ligaments.—The fossa in its lower edge is not so deep as in the upper ribs, because the vessels run more towards the interstice between the ribs.—Its fore-end is smaller than its body, and its short small cartilage is but loosely connected to the cartilage of the rib above.

The twelfth rib is the shortest and straightest.——Its head is only articulated with the last vertebra of the thorax; therefore is not divided into two surfaces.—
This rib is not joined to the transverse process of the vertebra; and therefore has no tubercle, being often pulled necessarily inwards by the diaphragm, which an articulation with the transverse process would not have allowed.——The fossa is not found at its under edge, because the vessels run below it.——The fore-part of this rib is smaller than its middle, and has only a very small pointed cartilage fixed to it.——To its whole internal side the diaphragm is connected.

The motions and uses of the ribs shall be more particularly treated of after the description of the sternum.

The heads and tubercles of the ribs of a new-born child have cartilages on them; part of which becomes afterwards thin epiphyses.—The bodies of the ribs encroach gradually after birth upon the cartilages; so that the latter are proportionally shorter, when compared to the ribs, in adults than in children.

Here I cannot help remarking the wife providence of our Creator, in preserving us from perishing as soon as we come into the world. The end of the bones of the limbs remain in a cartilaginous state after birth, and are many years before they are entirely united to the main body of their several bones; whereas the condyles

an

F

to

ar

al

fa

CI

tl

t

dyles of the occipital bone, and of the lower jaw, are true original processes, and ossisted before birth; and the heads and tubercles of the ribs are nearly in the same condition: and therefore the weight of the large head is sirmly supported; the actions of sucking, swallowing, respiration, &c. which are indispensably necessary for us as soon as we come into the world, are performed without danger of separating the parts of the bones that are most pressed on in these motions. Whereas, had these processes of the head, jaw, and ribs, been epiphyses at birth, children must have been exposed to danger of dying by such a separation; the immediate consequences of which would be the compression of the beginning of the spinal marrow, or want of food, or a stop put to respiration.

### THE STERNUM.

THE sternum (y), or breast-bone, is the broad slat bone, or pile of bones, at the fore-part of the thorax. The number of bones which this should be divided into, has occasioned debates among anatomists, who have considered it in subjects of different ages.—In adults of a middle age, it is composed of three bones, which easily separate after the cartilages connecting them are destroyed. Frequently the two lower bones are found intimately united; and very often, in old people, the sternum is a continued bony substance from one end to the other; though we still observe two, sometimes three, transverse lines on its surface; which are marks of the former divisions.

When we consider the sternum as one bone, we find it broadest and thickest above, and becoming smaller as it descends. The internal surface of this bone is somewhat hollowed for enlarging the thorax: but the convexity on the external surface is not so conspicuous, because the sides are pressed outwards by the true ribs; the round heads of whose cartilages are received into seven

<sup>(</sup>y) Erneos, Os pectoris, ensiforme, scutum cordis.

I.

re

he

ge

al-

ef-

re

ne

3%

id

en

ne

nnt

at

K.

1-

10

n

s,

g

d

n

d

r

feven smooth pits formed in each side of the sternum, and are kept firm there by strong ligaments, which on the external surface have a particular radiated texture (z). Frequently the cartilaginous sibres thrust themselves into the bony substance of the sternum, and are joined by a fort of suture.—The pits at the upper part of the sternum are at the greatest distance one from another, and, as they descend, are nearer; so that the two lowest are contiguous

The fubstance of the breast-bone is cellular, with a very thin external plate, especially on its internal surface, where we may frequently observe a cartilaginous crust spread over it. (a)—On both surfaces, however, a strong ligamentous membrane is closely braced; and the cells of this bone are so small, that a considerable quantity of osseous sibres must be employed in the composition of it. Whence, with the defence which the muscles give it, and the moveable support it has from the cartilages, it is sufficiently secured from being broken: for it is strong by its quantity of bone; its parts are kept together by ligaments; and it yields enough to elude considerably the violence offered (b).

So far may be faid of this bone in general; but the three bones, of which, according to the common account, it is composed in adults, are each to be examined.

The first, all agree, is somewhat of the figure of a heart, as it is commonly painted; only it does not terminate in a sharp point.—This is the uppermost thickest part of the sternum.

The upper middle part of this first bone, where it is thickest, is hollowed, to make place for the trachea arteria; though this cavity (c) is principally formed by the bone being raised on each side of it, partly by the clavicles

<sup>(</sup>z) Ruysch. Catalog. Rar. fig. 9.

<sup>(</sup>a) Jac. Sylv. in Galen de offibus, cap. 12.

<sup>(</sup>b) Senec. in Memoires de l'acad des sciences, 1724.

<sup>(</sup>c) Zeayn, Jugulum, furcula superior.

clavicles thrusting it inwards, and partly by the sternomastoidei muscles pulling it upwards.—On the outside of each tubercle there is an oblong cavity, that,
in viewing it transversely from before backwards, appears a little convex. Into these glenæ the ends of the
clavicles are received.—Immediately below these, the
sides of this bone begin to turn thinner; and in each
a superficial cavity or a rough surface is to be seen,
where the first ribs are received or joined to the sternum—In the side of the under-end of this sirst bone,
the half of the pit for the second rib on each side is
formed.—The upper part of the surface behind is covered with a strong ligament, which secures the clavicles; and is afterwards to be more particularly taken
notice of.

The fecond or middle division of this bone is much longer, narrower, and thinner, than the first; but excepting that it is a little narrower above than below, it is nearly equal all over in its dimensions of breadth or thickness.—In the sides of it are complete pits for the. third, fourth, fifth, and fixth ribs, and an half of the pits for the second and seventh; the lines, which are marks of the former division of this bone, being extended from the middle of the pits of one fide to the middle of the corresponding pits of the other fide. Near its middle an unoffified part of the bone is sometimes found; which, freed of the ligamentous membrane or cartilage that fills it, is described as a hole: and in this place, for the most part, we may observe a transverse line, which has made authors divide this bone into two.—When the cartilage between this and the first bone is not offified, a manifest motion of this upon the first may be observed in respiration, or in raising the sternum, by pulling the ribs upwards, or distending the lungs with air in a recent subject.

The third bone is much less than the other two, and has only one half of the pit for the seventh rib formed in it; wherefore it might be reckoned only an appen-

X

Ch

dix

tila

lag

the

COL

app

is 1

in

lar

cu

co

po

ba

br

for

fer

gr

a

no

th

fu

C

fo

tl

0

d

f

t.

10-

ut-

at,

p-

he

he

ch

en,

er-

ie,

is

:0-

la-

en

ch

X-

it

or

ne.

ne

re

.

le

-

1-

a

e

n

dix of the sternum.—In young subjects it is always cartilaginous, and is better known by the name of cartilago xiphoides or ensiformis (d) than any other: though the ancients often called the whole sternum ensiforme; comparing the two first bones to the handle, and this appendix to the blade, of a fword.——This third bone is feldom of the fame figure, magnitude, or fituation, in any two subjects; for sometimes it is a plain triangular bone, with one of the angles below, and perpendicular to the middle of the upper fide, by which it is connected to the fecond bone.—In other people the point is turned to one fide, or obliquely forwards or backwards. Frequently it is all nearly of an equal breadth, and in feveral subjects it is bifurcated; whence fome writers give it the name of furcella or furula inferior; or elle it is unoffified in the middle.——In the greatest number of adults, it is offished, and tipped with a cartilage; in some, one half of it is cartilaginous; and in others, it is all in a cartilaginous state. Generally several oblique ligaments, fixed at one end to the cartilages of the ribs, and by the other to the outer furface of the xiphoid bone, connect it firmly to those cartilages (e).

So many different ways this small bone may be formed without any inconvenience: But then some of these positions may be so directed, as to bring on a great train of ill consequences; particularly when the lower end is offissed, and is too much turned outwards or inwards (f), or when the conjunction of this appen-

dix with the second bone is too weak (g).

Vol. I. N The

(d) Clypealis, gladialis, mucronata, malum granatum. scutum stomachi, epiglottalis, cultralis, medium furculæ inferiores, scutiformis, ensiculata.

(e) Weitbrecht, Syndefmolog. p. 121.

(f) Rolfinc. Differt. Anat lib. 2. cap. 41.—Paaw de offib. pars 1. cap. 3. & pars 3. cap. 3.—Codronchi de prolapfu cartilagin. mucronat.

(g) Paaw, de offib. pars 1. cap. 3. & part 3. cap. 3.—Borrich. Act. Hafn. vol. 5. ob. 79 —Bonet. Sepulchret. Anat. tom. 2. lib. 3.

§ 5. Append. ad obf. 8. et ibid. § 7. obf. 19.

(

tl

fi

C

r

10

0

a

i

f

r

b

r

d

The sternum is joined by cartilages to the seven upper ribs, unless when the first coalesces with it in an intimate union of substance; and its unequal cavity on each side of its upper end is sitted for the ends of the clavicles.

The sternum most frequently has four round small bones, surrounded with cartilage, in children born at the full time; the uppermost of these, which is the sirst bone, being the largest.—Two or three other very small bony points are likewise to be seen in several children.—The number of bones increases for some years, and then diminishes, but uncertainly, till they are at last united into those above described of an adult.

The uses of this bone are, to afford origin and infertion to several muscles; to sustain the mediastinum; to defend the vital organs, the heart and lungs, at the fore-part; and, lastly, by serving as a moveable sustain of the ribs, to affist considerably in respiration: which action, so far as it depends on the motion of the

bones, we are now at liberty to explain.

When the ribs that are connected by their cartilages to the sternum, or to the cartilages of the true ribs, are acted upon by the intercostal muscles, they must all be pulled from the oblique position which their cartilages kept them in, nearer to right angles with the vertebræ and sternum, because the first or uppermost rib is by much the most fixed of any; and the cartilages making a great refistance to raising the anterior ends of the ribs, their large arched middle parts turn outwards as well as upwards. The sternum, pressed strongly on both fides by the cartilages of the ribs, is pushed forwards, and that at its several parts, in proportion to the length and motion of its supporters the ribs; that is, most at its lower end. The sternum and the cartilages, thus raifed forwards, must draw the diaphragm connected to them; confequently fo far stretch it, and bring it nearer to a plane. The power that raises this bone and the cartilages, fixes them sufficiently up.

in-

on the

iall

at

ery

hil-

irs,

at

er-

to

the

ul-

n:

the

ges

bs,

ila-

er-

rib

ges

nds

ut-

fed

is

ro-

he

ım

he

far

ver

uf-

tly

ficiently to make them refift the action of the diaphragm, whose fibres contract at the same time, and thrust the viscera of the abdomen downwards. ——The arched part of the ribs being thus moved outwards, their anterior ends and the sternum being advanced forwards, and the diaphragm being brought nearer to a plain furface, instead of being greatly convex on each fide within each cavity of the thorax, it is evident how considerably the cavity, of which the nine or ten upper ribs are the fides, must be widened, and made deeper and longer. While this is doing in the upper ribs, the lower ones, whose cartilages are not joined to the sternum or to other cartilages, move very differently, tho they conspire to the same intention, the enlargement of the thorax: for having no fixed point to which their anterior ends are fastened, and the diaphragm being inferted into them at the place where it runs pretty straight upwards from its origin at the vertebræ, these ribs are drawn downwards by this strong muscle, and by the muscles of the abdomen, which at this time are refifting the stretching force of the bowels; while the intercostal muscles are pulling them in the contrary direction, to wit, upwards. The effect, therefore, of either of these powers, which are antagonists to each other, is very little, as to moving the ribs either up or down; but the muscles of the abdomen, pushed at this time outwards by the vifcera, carry these ribs along with them.—Thus the thorax is not only not allowed to be shortened, but is really widened at its lower part, to affift in making fufficient space for the due distention of the lungs.

As foon as the action of these several muscles ceases, the elastic cartilages, extending themselves to their natural situation, depress the upper ribs, and the sternum subsides; the diaphragm is thrust up by the viscera abdominalia, and the oblique and transverse muscles of the belly serve to draw the inserior ribs inwards at the

N 2

fame

ta

W

W

of

ne

is

w vi is

er

th

at

CC

re

ar

ar

in

te

fi

de

tub

ft lo

e

al

a

fame time.—By these causes, the cavity of the breast is diminished in all its dimensions.

Though the motions above described of the ribs and sternum, especially of the latter bone, are so small in the mild respiration of a healthy person, that we can scarce observe them; yet they are manifest whenever we designedly increase our respiration, or are obliged to do it after exercise, and in several diseases.

# SECT. III. OF THE SUPERIOR EXTREMITIES.

A Uthors are much divided in their opinions about the number of bones which each superior extremity (b) should be said to consist of; some describing the clavicle and scapula as part of it, others classing these two bones with those of the thorax: But since most quadrupeds have no clavicles, and the human thorax can perform its functions right when the scapula is taken away (i), whereas it is impossible for us to have the right use of our arms without these bones, I must think that they belong to the superior extremities; and therefore shall divide each of them into the shoulder, arm, fore-arm, and hand.

# § 1. Bones of the SHOULDER.

THE SHOULDER consists of the clavicle and fea-

### CLAVICULA.

CLAVICULA, or collar-bone (k), is the long crooked bone, in figure like an Italic f, placed almost horizon-tally

<sup>(</sup>h) κωλα, γυα, εκφυαδες, Enata, adnata, explantata membra, artus.

<sup>(</sup>i) Philosoph. Transact. numb. 449. § 5.
(k) Os jugulare, jugulum, furcula, ligula, clavis, humerus quibusdam.

tI.

aft

nd

in

can

ver

ged

ES.

out

tre-

ing

ling

nce

ho-

a is

ave

nust

and

der,

sca-

ked

zon-

tally

, ar-

qui-

tally between the upper lateral part of the sternum, and what is commonly called the top of the shoulder: which, as a clavis or beam, it bears off from the trunk of the body.

The clavicle, as well as other long round bones, is larger at its two ends than in the middle. next to the sternum (1) is triangular: The angle behind is confiderably protruded, to form a sharp ridge, to which the transverse ligament, extended from one clavicle to the other, is fixed (m). The fide opposite to this is somewhat rounded. The middle of this protuberant end is as irregularly hollowed as the cavity in the sternum for receiving it is raised: but, in a recent subject, the irregular concavities of both are supplied by a moveable cartilage; which is not only much more closely connected every where by ligaments to the circumference of the articulation than those of the lower jaw are, but it grows to the two bones at both its internal and external end; its substance at the external end being foft, but very strong, and resembling the intervertebral cartilages (n).

From this internal end, the clavicle, for about twofifths of its length, is bended obliquely forwards and downwards. On the upper and fore part of this curvature a fmall ridge is feen, with a plain rough furface before it; whence the musculus sterno-hyoideus and sterno-mastoideus have in part their origin. Near the lower angle a small plain furface is often to be remarked, where the first rib and this bone are contiguous (0), and are connected by a firm ligament (p). From this a rough plain furface is extended outwards, where the pectoral muscle has part of its origin. Behind, the

bone

<sup>(1)</sup> Hagarpayis.

<sup>(</sup>m) Riolan. Encheirid. anat. lib. 6. cap. 13. - Winflow, Expol. Anat. des os frais, § 248 .- Weitbrecht. Act. Petropolit. tom. 4. p 255. et Syndesmolog sect. 2. I. § 3.

<sup>(</sup>n) Weitbrecht. Syndefmolog. fect. 2. I. § 6.

<sup>(0)</sup> Dionis, Sixieme demonst. des os.

<sup>(</sup>p) Weitbrecht. Syndesmolog. sect. 2. I. § 7.

ti

th

tr

ot

b

th

21

m

c

tl

a

e

th

tl

0

ra

it

bone is made flat and rough by the infertion of the larger share of the subclavian muscle. After the clavicle begins to be bended backwards, it is round: but it foon after becomes broad and thin; which shape it retains to its external end. Along the external concavity, a rough finuofity runs, from which some part of the deltoid muscle takes its rise: Opposite to this, on the convex edge, a scabrous ridge give infertion to a share of the cucullaris muscle. The upper furface of the clavicle here is flat; but the lower is hollow, for lodging the beginning of the musculus subclavius; and towards its back-part a tubercle rifes; to which, and a roughness near it, the strong short thick ligament connecting this bone to the coracoid process of the scapula is fixed.

The external end (q) of this bone is horizontally oblong, smooth, sloping at the posterior side, and tipped in a recent subject with a cartilage, for its articulation with the acromion scapulæ. Round this the bone is spongy, for the sirmer connection of the ligaments.

The medullary arteries having their direction obliquely outwards, enter the clavicles by one or more

fmall passages in the middle of their back-part.

The fubstance of this bone is the same as of the other

round long bones.

The triangular unequal interior end of each clavicle has the cartilage above described interposed betwixt it and the irregular cavity of the sternum — The ligaments which surround this articulation to secure it, are so short and strong, that little motion can be allowed any way; and the strong ligament that is stretched across the upper furcula of the sternum, from the posterior prominent angle of the one clavicle to the same place of the other clavicle, serves to keep each of these bones more firmly in their place.—By the assistance, however, of the moveable intervening cartilage, the clavicle can, at this joint, be raised or depressed, and moved

1

a

.

.

g

8

)•

d

n

18

e

er

le

it

a-

re

d

d

0.

e

se!

e,

ne id id moved backwards and forwards so much, as that the external end, which is at a great distance from that axis, enjoys very conspicuous motions. The articulation of the exterior end of the clavicle shall be considered after the description of the scapula.

The clavicles of infants are not deficient in any of their parts; nor have they any epiphyles at their extremules joined afterwards to their bodies, as most other such long bones have, which preserves them from being bended too much, and from the danger of any unoslified parts being separated by the force which pulls the arms forwards.

The uses of the clavicles are, to keep the scapulæ, and consequently all the superior extremities, from falling in and sorward upon the thorax; by which, as in most quadrupeds, the motions of the arms would be much confined, and the breast made too narrow. The clavicles likewise afford origin to several muscles, and a defence to large vessels.

From the fituation, figure, and use of the clavicles, it is evident that they are much exposed to fractures; that their broken parts must generally go by each other; and that they are difficultly kept in their place afterwards.

#### SCAPULA.

Scapula, or shoulder-blade (q), is the triangular bone situated on the outside of the ribs; with its long-est side, called its base, towards the spinal processes of the vertebræ; and with the angle at the upper part of this side about three inches, and the lower angle at a greater distance, from these processes. The back-part of the scapula has nothing but the thin ends of the serratus anticus major and subscapularis muscles between it and the ribs: But as this bone advances forwards, its distance from the ribs increases. The upper or shortest

<sup>(</sup>r) Ωμοπλατος, επινωτιον, Latitudo humeri, sceptulum vel scutulum opertum, spatula, ala, humerus, clypeus, scutum thoracis.

fide, called the superior costa of the scapula, is nearly horizontal, and parallel with the fecond rib. The lower fide, which is named the inferior costa, is extended obliquely from the third to the eighth rib. The fituation of this bone, here described, is when people are fitting or standing in a state of inactivity, and allowing the members to remain in the most natural easy po-The inferior angle of the scapula is very acute; the upper one is near to a right angle; and what is called the anterior does not deferve the name, for the two fides do not meet to form an angle. The body of this bone is concave towards the ribs, and convex behind, where it has the name of dorfum (s) Three proceffes are generally reckoned to proceed from the scapula. The first is the large spine that rises from its convex furface behind, and divides it unequally. The fecond process stands out from the fore-part of the upper fide; and, from its imaginary refemblance to a crow's beak, is named coracoides (t). The third process is the whole thick bulbous fore-part of the bone.

After thus naming the feveral constituent parts of the scapula, the particular description will be more easily

understood.

The base, which is tipped with cartilage in a young subject, is not all straight: for above the spine it runs obliquely forwards to the superior angle, that here it might not be too protuberant backwards, and so bruise the muscles and teguments. Into the oblique space the musculus patientiæ is inserted. At the root of the spine, on the back-part of the base, a triangular plain surface is formed by the pressure of the lower sibres of the trapezius. Below this the edge of the scapula is scabrous and rough, for the insertion of the serratus major anticus and rhomboid muscles.

The back-part of the inferior angle is made smooth by the latissimus dorsi passing over it. This muscle also

alters

<sup>(</sup>s) XEXIVIOV.

<sup>(</sup>t) Anchoroides, figmoides, digitalis, ancistroides.

Ī.

ly

N-

ed

1-

re

g

0-

:;

18 ie

of

e.

0-

a-

ts

IC ).

).

e y

e

e

n f alters the direction of the inferior costa some way forwards from this angle: and fo far it is flatted behind by the origin of the teres major. As the inferior costa advances forward, it is of considerable thickness, is flightly hollowed and made fmooth behind by the teres minor, while it has a fossa formed into it below by part of the subscapularis; and between the two a ridge with a fmall depretion appears, where the longus extensor cubiti has its origin.

The superior costa is very thin: and near its forepart there is a femilunar niche, from one end of which to the other a ligament is stretched; and sometimes the bone is continued to form one, or fometimes two, holes for the passage of the scapular blood-vessels and Immediately behind this femilunar cavity the coracohyoid muscle has its rife. From the niche to the termination of the fossa for the teres minor, the scapula is narrower than any where elfe, and supports the third

This part has the name of cervix.

The whole dorfum of the scapula is always faid to be convex; but, by reason of the raised edges that surround it, it is divided into two cavities by the spine, which is stretched from behind forwards, much nearer to the superior than to the inferior costa.—The cavity above the spine is really concave where the supra-spinatus muscle is lodged; while the surface of this bone below the fpine, on which the infra-spinatus muscle is placed, is convex, except a fosta that runs at the fide of the inferior costa.

The internal or anterior furface of this bone is hollow, except in the part above the spine, which is convex.——The subscapularis muscle is extended over this furface, where it forms feveral ridges and intermediate depressions, commonly mistaken for prints of the ribs; they point out the interffices of the bundles of fibres of which the fubscapularis muscle is composed (u).

The

<sup>(4)</sup> Winflow, in Memoires de l'acad. des sciences, 1722.

The spine (x) rises small at the base of the scapula. and becomes higher and broader as it advances forwards.—On the fides it is unequally hollowed and crooked, by the actions of the adjacent muscles.-Its ridge (y) is divided into two rough flat furfaces: Into the upper one, the trapezius muscle is inserted; and the lower one has part of the deltoid fixed to it. - The end of the spine, called acromion (z), or top of the shoulder, is broad and flat, and is fometimes only joined to the spine by a cartilage (a).—The anterior edge of the acromion is flat, smooth, and covered with a cartilage, for its articulation with the external end of the clavicle; and it is hollowed below, to allow a paffage to the infra and fupra spinati muscles, and free motion to the os humeri.

The coracoid (b) process is crooked, with its point inclining forwards; fo that a hollow is left at the lower fide of its root, for the passage of the infra-scapularis muscle.—The end of this process is marked with three plain furfaces. Into the internal, the ferratus minor anticus is inferted: from the external, one head of the biceps flexor cubiti rifes; and from the lower one, the coraco-brachialis has its ori-At the upper part of the root of this process, immediately before the femilunar cavity, a fmooth tubercle appears, where a ligament from the clavicle is fixed. From all the external fide of this coracoid apophyse, a broad ligament goes out, which becomes narrower where it is fixed to the acromion. The fharp pain, violent inflammation, and tedious cure of contusions in this part, are probably owing to these tendons and ligaments being hurt.

From the cervix fcapulæ the third process is produ-

ced.

(x) Ραχις, υπιζοχη ωμοπλατών, Eminentia scapularum.

(y) Pterigium, crista.

<sup>(</sup>z) Επωμις αγχυροειδης, κορακοειδης, κατακλεις, Acromii os, fummus armus, rostrum porcinum, processus digitalis.

<sup>(</sup>a) Sue Trad. d'osteol. p. 160. (b) 'Αγχυροειδης, σιγμοειδης, Rostriformis.

ced. The fore-part of this is formed into a glenoid cavity (c), which is of the shape of the longitudinal section of an egg, being broad below and narrow above. Between the brims of this hollow and the fore-part of the root of the spine, a large sinuosity is lest for the transmission of the supra and infra spinati muscles; and on the upper part of these brims we may remark a sinooth surface, where the second head of the biceps slexor cubiti has its origin. The root of the supercilia is rough all round, for the sirmer adhesion of the capsular ligament of the articulation, and of the cartilage which is placed on these brims, where it is thick, but becomes very thin as it is continued towards the middle of the cavity, which it lines all over.

The medullary veffels enter the fcapula near the bafe

of the spine.

The fubstance of the scapula, as in all other broad flat bones, is cellular, but of an unequal thickness: for the neck and third process are thick and strong; the inferior costa, spine, and coracoid process, are of a middle thickness; and the body is so pressed by the muscles, as

to become thin and diaphanous.

The scapula and clavicle are joined by plain surfaces, tipped with cartilage (d); by which neither bone is allowed any considerable motion, being tightly tied down by the common capsular ligament, and by a very strong one which proceeds from the coracoid process; but divides into two before it is fixed into the clavicle, with such a direction, as can either allow this bone to have a small rotation, in which its posterior edge turns more backwards, while the anterior one rises farther forwards; or it can yield to the fore-part of the scapula moving downwards, while the back-part of it is drawn upwards; in both which cases, the oblong smooth articulated surfaces of the clavicle and scapula are not in the same plane, but stand a little transversely, or across

<sup>(</sup>c) AMONOTUNIS.

<sup>(</sup>d) Acromion, xaraxxiis, Claufuræ.

each other, and thereby preserve this joint from luxations, to which it would be subject if either of the bones was to move on the other perpendicularly up and down, without any rotation. Sometimes a moveable ligamentous cartilage is found in this joint; otherwhiles fuch a cartilage is only interposed at the anterior half of it; and in some old subjects I have found a fesamoid bone here (e). The scapula is connected to the head, os hyoides, vertebræ, ribs, and arm-bone, by muscles, that have one end fastened to these bones, and the other to the scapula, which can move it upwards, downwards, backwards, or forwards; by the quick succession of these motions, its whole body is carried in a circle. But being also often moved as upon an axis perpendicular to its plane, its circumference turns in a circle whose centre this axis is (f). Whichever of these motions it performs, it always carries the outer end of the clavicle and the arm along with it. The glenoid cavity of this bone receives the os humeri, which plays in it as a ball in a focket, as will be explained more hereafter.

The use of the scapula is, to serve as a fulcrum to the arm; and by altering its position on different occasions, to allow always the head of the os humeri a right-situated socket to move in; and thereby to assist and to enlarge greatly the motions of the superior extremity, and to afford the muscles which rise from it more advantageous actions, by altering their directions to the bone which they are to move.—This bone also serves to defend the back-part of the thorax, and is often employed to sultain weights, or to resist forces,

too great for the arm to bear.

The base, acromion, coracoid process, and head of the scapula, are all in a cartilaginous state at birth; and the three first are joined as epiphyses; while the head, with the glenoid cavity, is not formed into a distinct

<sup>(</sup>e) Jac. Sylv. Isagog. Anat. lib. 1. cap. 2.
(f) See Winslow, Memoires de l'acad. des sciences, 1726,

distinct separate bone, but is gradually produced by the offification of the body of this bone being continued forwards.

# § 2. The ARM.

THE Arm has only one bone, best known by the Latin name of os humeri (g); which is long, round, and

nearly straight.

The upper end of this bone (b) is formed into a large round smooth head, whose middle point is not in a straight line with the axis of the bone, but stands obliquely backwards from it. The extent of the head is distinguished by a circular fossa surrounding its base, where the head is united to the bone, and the capfular ligament of the joint is fixed. Below the fore-part of its base two tubercles stand out: The smallest one. which is fituated most to the inside, has the tendon of the fubscapularis muscle inserted into it. ——The larger more external protuberance is divided, at its upper part, into three fmooth plain furfaces; into the anterior of which the musculus supra-spinatus; into the middle or largest, the infra-spinatus; into the one behind, the teres minor, is inferted.—Between these two tubercles, exactly in the fore-part of the bone, a deep long fossa is formed, for lodging the tendinous head of the biceps flexor cubiti; which, after passing, in a manner peculiar to itself, through the cavity of the articulation, is tied down by a tendinous sheath extended across the fossa; in which, and in the neighbouring tubercles, are feveral remarkable holes, which are penetrated by the tendinous and ligamentous fibres, and by veffels.—On each fide of this fossa, as it descends in the os humeri, a rough ridge, gently flatted in the middle, runs from the roots of the tubercles. The tendon of the

(b) Acrocolium.

<sup>(</sup>g) Ακρολία, αλένη. Os brachii, armi, adjutorium, parvum, bra-

the pectoral muscle is fixed into the anterior of these ridges, and the latissimus dorsi and teres major are inferted into the internal one. A little behind the lower end of this last, another rough ridge may be observed, where the coraco-brachialis is inferted. From the back. part of the root of the largest tubercle a ridge also is continued, from which the brevis extensor cubiti rifes. This bone is flatted on the infide, about its middle, by the belly of the biceps flexor cubiti. In the middle of this plain furface, the entry of the medullary artery is feen flanting obliquely downwards. At the fore-fide of this plane the bone rifes in a fort of ridge, which is rough, and often has a great many small holes in it, where the tendon of the strong deltoid muscle is inferted; on each fide of which the bone is smooth and flat, where the brachiæus internus rifes. The exterior of these two flat surfaces is the largest; behind it a superficial spiral channel, formed by the muscular nerve and the veffels that accompany it, runs from behind forwards and downwards. The body of the os humeri is flatted behind by the extensors of the fore-arm.

Near the lower end of this bone, a large sharp ridge is extended on its outside, from which the musculus spinator radii longus, and the longest head of the extensor carpi radialis, rise. Opposite to this, there is another small ridge to which the aponeurotic tendon, that gives origin to the sibres of the internal and external brachiæi muscles, is fixed; and from a little depression on the fore-side of it, the pronator radii teres rises.

The body of the os humeri becomes gradually broader towards the lower end, where it has feveral process; at the roots of which there is a cavity before and another behind (i). The anterior is divided by a ridge into two; the external, which is the least, receives the end of the radius; and the internal receives the coronoid

noid process of the ulna in the flexions of the fore-arm, while the posterior deep triangular cavity lodges the olecranon in the extensions of that member. The bone betwixt these two cavities is pressed so thin by the proceffes of the ulna, as to appear diaphanous in feveral subjects. The fides of the posterior cavity are stretched out into two processes, one on each side: These are called condyles; from each of which a strong ligament goes out to the bones of the fore-arm. The external condyle, which has an oblique direction also forwards in respect of the internal, when the arm is in the most natural posture (k), is equally broad, and has an obtuse fmooth head rifing from it forwards. From the rough part of the condyle, the inferior head of the bicornis, the extensor digitorum communis, extensor carpi ulnaris, anconæus, and fome part of the spinator radii brevis, take their rife; and on the fmooth head the upper end of the radius plays. Immediately on the outfide of this, there is a finuofity made by the shorter head of the bicornis muscle, upon which the muscular nerve is placed. The internal condyle is more pointed and protuberant than the external, to give origin to some part of the flexor carpi radialis, pronator radii teres, palmaris longus, flexor digitorum fublimis, and flexor carpi ulnaris. Between the two condyles, is the trochlea or pulley; which confifts of two lateral protuberances, and a middle cavity, that are fmooth and covered with cartilage. When the fore-arm is extended, the tendon of the internal brachiæus muscle is lodged in the fore-part of the cavity of this pulley. The external protuberance, which is less than the other, has a sharp edge behind; but forwards, this ridge is obtufe, and only separated from the little head, already described, by a fmall fossa, in which the joined edges of the ulna and radius move. The internal protuberance of the pulley is largest and highest; and therefore, in the motions of the ulna upon it, that bone would be inclined outwards

<sup>(4)</sup> Winflow, Memoires de l'acad. des feiences, 1722.

(

t

t

b

fide.—Between this internal protuberance and condyle, a finuofity may be remarked, where the ulnar nerve passes.

The fubstance and the internal structure of the os humeri is the same, and disposed in the same way, as in

other long bones.

The round head at the upper end of this bone is articulated with the glenoid cavity of the scapula; which being fuperficial, and having long ligaments, allows the arm a free and extensive motion.—These ligaments are, however, confiderably strong. For, befides the common capfular one, the tendons of the muscles perform the office, and have been described under the name of ligaments.—Then the acromion and coracoid process, with the strong broad ligaments stretched betwixt them, secure the articulation above, where the greatest and most frequent force is applied to thrust the head of the bone out of its place. It is true, that there is not near fo strong a defence at the lower part of the articulation; but, in the ordinary postures of the arm, that is, so long as it is at an acute angle with the trunk of the body, there cannot be any force applied at this place to occasion a luxation, fince the joint is protected fo well above.

The motions which the arm enjoys by this articulation are to every fide; and by the fuccession of these different motions, a circle may be described. Besides which, the bone performs a small rotation round its own axis. But though this can be performed with the round head in all positions; yet as these vary, the essects upon the body of the bone are very different: For if the middle of the head is the centre of rotation, as it is when the arm hangs down by the side, the body of the bone is only moved forwards and backwards; because the axis of motion of the head is nearly at right angles with the length of the bone (1); whereas, when the arm is raised to right angles with the trunk of the body, the centre of motion, and the axis of the bone, come to be of the same straight line; and therefore the body of the os humeri performs the same motion with its head. Though the motions of the arm seem to be very extensive, yet the larger share of them depends on the motions of the scapula. The lower end of the os humeri is articulated with the bones of the fore-arm, and carries them with it in all its motions, but serves as a base on which they perform the motions peculiar to themselves; as shall be described afterwards.

Both the ends of this bone are cartilaginous in a new-born infant, and the large head with the two tubercles, and the trochlea with the two condyles, become epiphyses before they are united to the body of the bone.

## § 3. The Fore-ARM.

THE fore-arm (m) confifts of two long bones, the ulna and radius; whose situation, in respect of each other, is oblique in the least straining or most natural posture; that is, the ulna is not directly behind, nor on the outfide of the radius, but in a middle fituation between these two, and the radius crosses it. The situation, however, of these bones, and of all the other bones of the fuperior extremity that are not yet defcribed, is frequently altered: and therefore, to fhun repetitions, I defire it may be now remarked, that, in the remaining account of the superior extremity, I understand by the term of posterior, that part which is in the fame direction with the back of the hand; by anterior, that answering to the palm; by internal, that on the same side with the thumb; by external, the side nearest to the little finger; supposing the hand always to be in a middle position between the pronation and fupination.

Vol. I. O UL.

<sup>(</sup>m) Cubitus, taxus, axon, suyer, Ulna, lacertus.

#### ULNA.

ULNA (n), so named from its being used as a meafure, is the longest of the two bones of the fore-arm, and situated on the outside of the radius.

At the upper end of the ulna are two processes.-The posterior is the largest, and formed like a hook, whose concave surface moves upon the pulley of the os · humeri, and is called olecranon (o), or top of the cubit. The convex back-part of it is rough and scabrous, where the longus, brevis, and brachiæus externus, are inserted. The olecranon makes it unnecessary that the tendons of the extensor muscles should pass over the end of the os humeri; which would have been of ill consequence in the great flexions of this joint, or when any confiderable force is applied to this part (p). The anterior process is not so large, nor does it reach so high, as the one behind; but is sharper at its end, and therefore is named coronoid.—Between these two processes, a large semicircular or sigmoid concavity is left; the furface of which, on each fide of a middle rifing, is flanting, and exactly adapted to the pulley of the bone of the arm.—Across the middle of it, there is a small finuofity for lodging mucilaginous glands; where, as well as in a small hollow on the internal side of it, the cartilage that lines the rest of its surface is wanting.— Round the brims of this concavity the bone is rough, where the capfular ligament of the joint is implanted. -Immediately below the olecranon, on the backpart of the ulna, a flat triangular spongy surface appears, on which we commonly lean.—At the internal fide of this, there is a larger hollow furface, where the musculus anconæus is lodged; and the ridge at the

<sup>(</sup>n) Cubitus, \*nxuc, \*eo\*nxvov, Focile majus, canna vel arundo major, et inferior brachii.

<sup>(</sup>p) Winflow, Exposition anatomique du corps humain, traité des os secs, § 979.

infide of this gives rife to the musculus supinator radii brevis.—Between the top of the ridge and the coronoid process is the semilunated smooth cavity, lined with cartilage; in which, and a ligament extended from the one to the other end of this cavity, the round head of the radius plays.—Immediately below it, a rough hollow gives lodging to mucilaginous glands.—Below the root of the coronoid process, this bone is scabrous and unequal, where the brachiæus internus is inserted.—On the outside of that, we observe a smooth concavity, where the beginning of the slexor digitorum profundus sprouts out.

The body of the ulna is triangular.—The internal angle is very sharp where the ligament that connects the two bones is fixed: The sides which make this angle are flat and rough, by the action and adhesion of the many muscles which are situated here. At the distance of one-third of the length of the ulna from the top, in its fore-part, the passage of the medullary vessels is to be remarked slanting upwards. The external side of this bone is smooth, somewhat convex, and the angles at each edge of it are blunted by the pressure of

the muscles equally disposed about them.

As this bone descends, it becomes gradually smaller; so that its lower end terminates in a little head, standing on a small neck. Towards the fore but outer part of which last, an oblique ridge runs, that gives rise to the pronator radii quadratus. The head is round, smooth, and covered with a cartilage on its internal side, to be received into the semilunar cavity of the radius; while a styloid process (q) rises from its outside, to which is fixed a strong ligament that is extended to the os cuneiforme and pisisforme of the wrist. Between the back-part of that internal smooth side and this process, a sinuosity is lest for the tendon of the extensor carpi ulnaris. On the fore-part of the root of the process, such another depression may be remarked for the

<sup>(</sup>q) reaposion, Malleolus externus.

fage of the ulnar artery and nerve. The end of the bone is smooth, and covered with a cartilage. Between it and the bones of the wrist, a doubly concave moveable cartilage is interposed; which is a continuation of the cartilage that covers the lower end of the radius, and is connected loosely to the root of the styloid process, and to the rough cavity there; in which mucila-

ginous glands are lodged.

The ulna is articulated above with the lower end of the os humeri, where these bones have depressions and protuberances corresponding to each other, so as to allow an easy and secure extension of the fore-arm to almost a straight line with the arm, and slexion to a very acute angle; but, by the slanting position of the pulley, the lower part of the fore-arm is turned outwards in the extension, and inwards in the flexion (r); and a very small kind of rotation is likewise allowed in all positions, especially when the ligaments are most relaxed by the fore-arm being in a middle degree of slexion. The ulna is also articulated with the radius and carpus, in a manner to be related afterwards.

#### RADIUS.

Radius (s), so called from its imagined resemblance to a spoke of a wheel or to a weaver's beam, is the bone placed at the inside of the fore-arm. Its upper end is formed into a circular little head, which is hollowed for an articulation with the tubercle at the side of the pulley of the os humeri; and the half of the round circumference of the head next to the ulna is smooth, and covered with a cartilage, in order to be received into the semilunated cavity of that bone.—

Below the head, the radius is much smaller; therefore this part is named its cervix, which is made round by the action of the supinator radii brevis.—At the external root of this neck, a tuberous process rises; into

(r) Winflow, Memoires de l'acad. des sciences, 1722.

<sup>(1)</sup> Kigxis, παραπηχίον, Focile minus, canna minor, arundo minor-

n

f

f

d

d

11

f

r

e

the outer part of which the biceps flexor cubiti is inferted.—From this a ridge runs downwards and inwards, where the fupinator radii brevis is inferted; and a little below, and behind this ridge, there is a rough scabrous surface, where the pronator radii teres is fixed.

The body of the radius is not straight, but convex on its internal and posterior surfaces; where it is also made round by the equal pressure of the circumjacent muscles, particularly of the extensors of the thumb; but the surfaces next to the ulna are flatted and rough, for the origin of the muscles of the hand; and both terminate in a common sharp spine, to which the strong ligament extended betwixt the two bones of the forearm is fixed. A little below the beginning of the plain surface, on its fore-part, where the slexor muscle of the last joint of the thumb takes its origin, the passage of the medullary vessels is seen slanting upwards. The radius becomes broader and flatter towards the lower end, especially on its fore-part, where its pronator quadratus muscle is situated.

The lower end of the radius is larger than the superior; though not in fuch a disproportion as the upper end of the ulna is larger than its lower end. -- Its backpart has a flat strong ridge in the middle, and fossæ on each fide ——In a fmall groove, immediately on the outside of the ridge, the tendon of the extensor tertii internodii pollicis plays. In a large one beyond this, the tendons of the indicator and of the common extenfor muscles of the fingers pass. Contiguous to the ulna there is a fmall depression made by the extensor minimi digiti.—On the infide of the ridge there is a broad depression, which seems again subdivided, where the two tendens of the bicornis, or extensor carpi radialis, are lodged. The internal fide of this end of the radius is also hollowed by the extensors of the first and tecond joint of the thumb; immediately above which, a little rough furface shows where the supinator radii lon-

0 3

gus is inserted.—The ridges at the sides of the grooves, in which the tendons play, have an annular ligament fixed to them, by which the feveral sheaths for the tendons are formed. The fore-part of this end of the radius is also depressed, where the flexors of the fingers and flexor carpi radialis pass.——The external fide is formed into a femilunated fmooth cavity, lined with a cartilage, for receiving the lower end of the ulna.-The lowest part of the radius is formed into an oblong cavity; in the middle of which is a small transverse rifing, gently hollowed, for lodging muciláginous glands; while the rifing itself is infinuated into the conjunction of the two bones of the wrift that are received into the cavity.—The internal fide of this articulation is fenced by a remarkable process (t) of the radius, from which a ligament goes out to the wrift, as the styloid process of the ulna with its ligament guards it on the outside.

The ends of both the bones of the fore-arm being thicker than the middle, there is a confiderable distance between the bodies of these bones; in the larger part of which a strong tendinous, but thin ligament, is extended, to give a large enough surface for the origin of the numerous sibres of the muscles situated here, that are so much sunk between the bones as to be protected from injuries, which they would otherwise be exposed to. But this ligament is wanting near the upper end of the fore-arm, where the supinator radii brevis, and slexor digitorum profundus, are imme-

diately connected (u).

Both ends of the bones of the fore-arm are first car-

tilages, and then epiphyses, in children.

As the head of the radius receives the tubercle of the os humeri, it is not only bended and extended along with the ulna, but may be moved round its axis in any position; and that this motion round its axis may be sufficiently large, the ligament of the articulation is extended

(t) Malleollus internus, processus styloides.
(u) Weitbrecht. Syndesmolog. sig. 10, 11.

tended farther down than ordinary on the neck of this bone, before it is connected to it; and it is very thin at its upper and lower part, but makes a firm ring in the middle. This bone is also joined to the ulna by a double articulation: for above, a tubercle of the radius plays in a focket of the ulna; whilst below, the radius gives the socket, and the ulna the tubercle. But then the motion performed in these two is very differrent: for, at the upper end, the radius does no more than turn round its axis; while, at the lower end, it moves in a fort of cycloid upon the round part of the ulna; and as the hand is articulated and firmly connected here with the radius, they must move together. When the palm is turned uppermost, the radius is faid to perform the fupination: when the back of the hand is above, it is said to be prone. But then the quickness and large extent of these two motions are affished by the ulna, which, as was before observed, can move with a kind of fmall rotation on the floping fides of the pulley. This lateral motion, though very inconfiderable in the joint itself, is conspicuous at the lower end of fuch a long bone; and the strong ligament connecting this lower end to the carpus, makes the hand more readily to obey these motions. When we design a large circular turn of our hand, we increase it by the rotation of the os humeri, and fometimes employ the fpine and inferior extremities to make these motions of pronation or supination of the hand large enough.

# 6 4. The HAND.

THE hand (x) comprehends all from the joint of the wrist to the points of the fingers. Its back-part is convex, for greater firmness and strength; and it is concave before, for containing more surely and conveniently such bodies as we take hold of. One half of the hand has an obscure motion in comparison of what the other O<sub>4</sub> has,

<sup>(</sup>x) 'Axeoxie, Summa manus.

tr

21

cij

OI

na m

la fo

ar

h

to

te

b

0

lo

0

v fi

ti

V

a

0

b

has, and ferves as a base to the moveable half; which can be extended back very little farther than to a straight line with the fore-arm, but can be considerably bended forwards.

As the bones that compose the hand are of different shapes and uses, while several of them that are contiguous agree in some general characters; the hand is, on this account, commonly divided into carpus, metacarpus, and singers; among which last the thumb is reckoned.

#### CARPUS.

THE carpus (y) is composed of eight small spongy bones, fituated at the upper part of the hand. I shall describe each of these bones, under a proper name taken from their figure (z); because the method of ranging them by numbers leaves anatomists too much at liberty to debate, very idly, which ought to be preferred to the first number; or, which is worse, several, without explaining the order they observe, differently apply the same numbers, and so confound their readers. But, that the description of these bones may be in the fame order as they are found in the generality of anatomical books, I shall begin with the range of bones that are concerned in the moveable joint of the wrift, or are connected to the fore-arm, and shall afterwards confider the four that support the thumb and offa metacarpi of the fingers.

The eight bones of the CARPUS are, Os scaphoides, lunare, cuneiforme, pisiforme, trapezium, trapezoides,

magnum, unciforme.

The feaphoides is fituated most internally of those that are articulated with the fore-arm. The lunare is immediately on the outside of the former.—The cuneiforme is placed still more externally, but does not reach so high up as the other two.—The pistforme stands forwards

(z) Lyfer. Cult. Anat. lib 5. cap. 2.

<sup>(</sup>y) Kruc. Brachiale, prima palmæ pars, rasetta.

wards into the palm from the cuneiforme.—The trapezium is the first of the second row, and is situated betwixt the scaphoides and first joint of the thumb.—The trapezoides is immediately on the outside of the trapezium.—The os magnum is still more external.—The un-

ciforme is farther to the fide of the little finger.

Os scaphoides (a) is the largest of the eight, excepting one. It is convex above, concave and oblong below; from which small resemblance of a boat it has got its name. Its smooth convex surface is divided by a rough middle fossa, which runs obliquely cross it. The upper largest division is articulated with the radius. Into the sossa the common ligament of the joint of the wrist is sixed; and the lower division is joined to the trapezium and trapezoides. The concavity receives more than an half of the round head of the os magnum. The external side of this hollow is formed into a semilunar plane, to be articulated with the following bone.—The internal, posterior, and anterior edges are rough, for fixing the ligaments that connect it to the surrounding bones.

Os lunare (b) has a smooth convex upper surface, by which it is articulated with the radius. The internal side, which gives the name to the bone, is in the form of a crescent, and is joined with the scaphoid;—the lower surface is hollow, for receiving part of the head of the os magnum. On the outside of this cavity is another smooth, but narrow, oblong sinuosity, for receiving the upper end of the os unciforme:—on the outside of which a small convexity is found, for its connection with the os cuneiforme. Between the great convexity above, and the first deep inferior cavity, there is a rough sossa, in which the circular ligament of the joint of the wrist is fixed.

Os cuneiforme (c) is broader above, and towards the back of the hand, than it is below and forwards; which gives

<sup>(</sup>a) Kotunosides, Naviculare.

<sup>(</sup>b) Lunatum.

<sup>(</sup>c) Triquetrum.

d

gives it the resemblance of a wedge. The superior slightly convex surface is included in the joint of the wrist, being opposed to the lower end of the ulna.— Below this the cuneiform bone has a rough sossa, wherein the ligament of the articulation of the wrist is fixed. On the internal side of this bone, where it is contiguous to the os lunare, it is smooth and slightly concave. Its lower surface, where it is contiguous to the os unciforme is oblong, somewhat spiral, and concave. Near the middle of its anterior surface a circular plane appears, where the os pisiforme is sustained.

Os pisiforme (d) is almost spherical, except one circular plane, or slightly hollow surface, which is covered with cartilage for its motion on the cuneiform bone, from which its whole rough body is prominent forwards into the palm; having the tendon of the slexor carpiulnaris, and a ligament from the styloid process of the ulna, fixed to its upper part; the transverse ligament of the wrist is connected to its internal side; ligaments extended to the unciform bone, and to the os metacarpi of the little singer, are attached to its lower part; the abductor minimi digiti has its origin from its forepart; and, at the internal side of it, a small depression is formed, for the passage of the ulnar nerve.

Trapezium (e) has four unequal fides and angles in its back-part, from which it has got its name.—Above, its surface is smooth, slightly hollowed, and semicircular, for its conjunction with the os scaphoides.—Its external side is an oblong concave square, for receiving the following bone. The inferior surface is formed into a pulley; the two protuberant sides of which are external and internal. On this pulley the first bone of the thumb is moved.—At the external side of the external protuberance, a small oblong smooth surface is formed by the os metacarpi indicis. The fore-part of the trapezium is prominent in the palm, and near

(d) Cartilaginosum, subrotundum, rectum.

<sup>(</sup>e) Os cubiforme, trapezoides, multangulum, majus.

to the external fide has a finuofity in it, where the tendon of the flexor carpi radialis is lodged; on the ligamentous fheath of which the tendon of the flexor tertii internodii pollicis plays: And still more externally the bone is fcabrous, where the transverse ligament of the wrist is connected, the abductor and flexor primi internodii pollicis have their origin, and ligaments go out to the first bone of the thumb.

Os trapezoides (f), so called from the irregular quadrangular figure of its back-part, is the smallest bone of the wrist except the pisiforme. The figure of it is an irregular cube. It has a small hollow surface above, by which it joins the scaphoides; a long convex one internally, where it is contiguous to the trapezium; a small external one, for its conjunction with the os magnum; and an inferior convex surface, the edges of which are, however, so raised before and behind, that a sort of pulley is formed, where it sustains the os

metacarpi indicis.

Os magnum (g) fo called because it is the largest bone of the carpus, is oblong, having four quadrangular fides, with a round upper end, and a triangular plain one below. The round head is divided by a fmall rifing, opposite to the connection of the os scaphoides and lunare, which together form the cavity for receiving it. On the infide a short plain surface joins the os magnum to the trapezoides. On the outfide is a long narrow concave furface, where it is contiguous to the os unciforme. The lower end, which fuftains the metacarpal bone of the middle finger, is triangular, flightly hollowed, and farther advanced on the internal fide than on the external, having a confiderable oblong depression made on the advanced inside by the metacarpal bone of the fore-finger; and generally there is a small mark of the os metacarpi digiti annularis on its external fide.

Os

<sup>(</sup>f) Trapezium, multangulum minus. (g) Maximum, capitatum.

C

fi

fo

d

tl

n

tl

t

a

te

b

h

Os unciforme (b) has got its name from a thin broad process that stands out from it forwards into the palm, and is hollow on its infide, for affording paffage to the tendons of the flexors of the fingers. To this process also the transverse ligament is fixed that binds down and defends these tendons; and the flexor and abductor muscles of the little finger have part of their origin from The upper plain furface is fmall, convex, and joined with the os lunare: The internal fide is long, and flightly convex, adapted to the contiguous os mag-The external furface is oblique, and irregularly convex, to be articulated with the cuneiform bone. The lower end is divided into two concave furfaces; the external is joined with the metacarpal bone of the little finger; and the internal one is fitted to the metacarpal bone of the ring-finger.

In the description of the preceding eight bones, I have only mentioned those plain surfaces covered with cartilage, by which they are articulated to each other, or to some other bones, except in some few cases, where fomething extraordinary was to be observed; and I have defignedly omitted the other rough furfaces, left, by crowding too many words in the defcription of fuch small bones, the whole should be unintelligible. But these scabrous parts of the bones may eafily be understood after mentioning their figure, if it is observed, that they are generally found only towards the back or palm of the hand; that they are all plain, larger behind than before; and that they receive the different ligaments, by which they are either connected to neighbouring bones or to one another; for these ligaments cover all the bones, and are so accurately applied to them, that at first view the whole carpus of a recent subject appears one smooth bone (i).

As the furfaces of these bones are largest behind, the figure

(b) Cuneiforme.

<sup>(</sup>i) Galen. de usu part. lib. 2. cap. 8. For a particular description of these ligaments, see Weitbrecht. Syndesmolog. p. 5.-68.

figure of the whole conjoined must be convex there, and concave before; which concavity is still more increased by the os pisiforme, and process of the os unciforme, standing forwards on one side, as the trapezium does on the other: And the bones are securely kept in this form by the broad strong transverse ligament connected to those parts of them that stand prominent into the palm of the hand. The convexity behind renders the whole sabric stronger, where it is most exposed to injuries; and the large anterior hollow is necessary for a safe passage to the numerous vessels, nerves, and tendons of the singers.

The fubstance of these bones is spongy and cellular,

but strong in respect of their bulk.

The three first bones of the carpus make an oblong head, by which they are articulated with the cavity at the lower ends of the bones of the fore-arm, fo as to allow motion to all fides; and by a quick succession of these motions, they may be moved in a circle. But as the joint is oblong, and therefore the two dimensions are unequal, no motion is allowed to the carpus round its axis, except what it has in the pronation and fupination along with the radius.—The articulation of the first three bones of the superior row, with the bones of the inferior, is fuch as allows of motion, especially backwards and forwards; to the fecurity and eafiness of which, the reception of the os magnum into the cavity formed by the scaphoides and lunare contributes considerably: And the greatest number of the muscles that ferve for the motion of the wrist on the radius, being inferted beyond the conjunction of the first row of bones with the fecond, act equally on this articulation as they do on the former; but the joint formed with the radius being the most easily moved, the first effect of these muscles is on it; and the second row of the carpus is only moved afterwards. By this means a larger motion of the wrift is allowed than otherwise it could have had fafely: for if as large motion had been

Ł

b

tl

tl

n

It

fi

given to one joint, the angle of flexion would have been very acute, and the ligaments must have been longer than was consistent with the sirmness and security of the joint. The other articulations of the bones here being by nearly plain surfaces, scarce allow of any more motion, because of the strong connecting ligaments, than to yield a little, and so elude the force of any external power; and to render the back of the wrist a little more flat, or the palm more hollow, on proper occasions. The articulations of the thumb and metacarpal bones shall be examined afterwards.

The uses of the carpus are to serve as a base to the hand, to protect its tendons, and to afford it a free

large motion.

All the bones of the carpus are in a cartilaginous

flate at the time of birth.

On account of the many tendons that pass upon the lower end of the fore-arm and the carpus, and of the numerous ligaments of these tendons and of the bones, which have lubricating liquors supplied to them, the pain of sprains here is acute, the parts take long time to recover their tone, and their swellings are very obstinate.

# METACARPUS.

METACARPUS (k) consists of sour bones which suffain the singers. Each bone is long and round, with its ends larger than its body.—The upper end, which some call the base, is slat and oblong, without any considerable head or cavity; but it is however somewhat hollowed, for the articulation with the carpus: It is made slat and smooth on the sides where these bones are contiguous to each other. Their bodies are slatted on their back-part by the tendons of the extensors of the singers. The anterior surface of these bodies is a little concave, especially in their middle; along which

<sup>(</sup>k) Kreis, προκαρπίου, στηθος, ανδηρού, κτινίου, Postbrachiale, pectus, palma, pecten.

a sharp ridge stands out, which separates the musculi interossei placed on each side of these bones, which are

there made flat and plain by these muscles.

Their lower ends are raised into large oblong smooth heads, whose greatest extent is forwards from the axis of the bone. At the fore-part of each side of the root of each of these heads, one or two tubercles stand out, for fixing the ligaments that go from one metacarpal bone to another, to preserve them from being drawn as funder: Round the heads a rough ring may be remarked, for the capsular ligaments of the first joints of the singers to be fixed to; and both sides of these heads are slat, by pressing on each other.

The substance of the metacarpal bones is the same

with that of all long bones.

At the time of birth, these bones are cartilaginous at

both ends, which afterwards become epiphyses.

The metacarpal bones are joined above to the offar carpi and to each other by nearly plain surfaces. These connections are not fit for large motions. The articulation of their round heads at the lower ends with the cavities of the first bones of the singers, is to be taken notice of hereafter.

The concavity on the fore-part of these metacarpal bones, and the placing their bases on the arched carpus, cause them to form a hollow in the palm of the hand, which is often useful to us. The spaces between them lodge muscles, and their small motion makes them sit supporters for the singers to play on.

Though the offa metacarpi fo far agree, yet they may be distinguished from each other by the following

marks.

The os metacarpi indicis is generally the longest.——
Its base, which is articulated with the os trapezoides, is hollow in the middle. The small ridge on the internal side of this oblong cavity is smaller than the one opposite to it, and is made slat on the side by the trapezium. The exterior ridge is also smooth, and slat on its out-

fide,

fide, for its conjunction with the os magnum; immediately below which a femicircular smooth flat surface shows the articulation of this to the second metacarpal bone. The back-part of this base is flatted where the long head of the extensor carpi radialis is inserted, and its fore-part is prominent where the tendon of the flexor carpi radialis is fixed. The external side of the body of this bone is more hollowed by the action of muscles, than the internal. The tubercle at the internal root of its head is larger than the external. Its base is so simply fixed to the bone it is connected with, that it has no motion.

Os metacarpi medii digiti is generally the second in length: but often it is as long as the former; fometimes it is longer; and frequently it appears only to equal the first by the os magnum being farther advanced downwards than any other bone of the wrift. Its base is a broad superficial cavity, flanting outwards; the internal posterior angle of which is so prominent, as to have the appearance of a process. The internal fide of this base is made plain in the same way as the external fide of the former bone, while its external fide has two hollow circular furfaces, for joining the third metacarpal bone; and between these surfaces there is a rough fossa, for the adhesion of a ligament, and lodging mucilaginous glands. The shorter head of the bicornis is inserted into the back-part of this base. The two sides of this bone are almost equally flatted; only the ridge on the fore-part of the body inclines outwards. The tubercles at the fore-part of the root of the head are equal. The motion of this bone is very little more than the first metacarpal one has; and therefore these two firmly result bodies pressed against them by the thumb or fingers, or both.

Os metacarpi digiti annularis is shorter than the second metacarpal bone. Its base is semicircular and convex, for its conjunction with the os unciforme. On its internal side are too smooth convexities, and a middle

fossa,

fossa, adapted to the second metacarpal bone. The external side has a triangular smooth concave surface to join it with the fourth one. The anterior ridge of its body is situated more to the out than to the inside.—— The tubercles near the head are equal.— The motion of this third metacarpal bone is greater than the motion of the second.

Os metacarpi minimi digiti is the smallest and sharpest. Its base is irregularly convex, and rises slanting outwards.—Its internal side is exactly adapted to the third metacarpal bone.—The external has no smooth surface, because it is not contiguous to any other bone; but it is prominent where the extensor carpi ulnaris is inserted.—As this metacarpal bone is furnished with a proper moving muscle, has the plainest articulation, is most loosely connected and least confined, it not only enjoys a much larger motion than any of the rest, but draws the third bone with it, when the palm of the hand is to be made hollow by its advancement forwards, and by the prominence of the thumb opposite to it.

## THUMB AND FINGERS.

THE thumb and four fingers are each composed of three long bones.

The Thumb (1) is fituated obliquely in respect of the fingers, neither opposite directly to them, nor in the same plane with them.—All its bones are much thicker and stronger in proportion to their length, than the bones of the fingers are: Which are extremely necessary, fince the thumb counteracts all the fingers.

The first bone of the thumb has its base adapted to the double pulley of the trapezium: For, in viewing it from one side to the other, it appears convex in the middle; but when considered from behind forwards, it is concave there.—The edge at the fore-part of this Vol. I.

P base

<sup>(1)</sup> Avrixiip, dixbisuhos, Magnus digitus, promanus.

base is produced farther than any other part; and round the back-part of the base a rough sossa may be seen, for the connection of the ligaments of this joint. The body and head of this bone are of the same shape as the ossa metacarpi; only that the body is shorter, and the head slatter, with the tubercles at the fore-part

of its root larger.

The articulation of the upper end of this bone is uncommon: For, though it has protuberances and depressions adapted to the double pulley of the trapezium; yet it enjoys a circular motion, as the joints do where a round head of the one plays in the orbicular socket of another; only it is somewhat more confined, and less expeditious, but stronger and more secure than such joints generally are.

This bone of children is in the same state with the

metacarpal bones.

The fecond bone of the thumb has a large base formed into an oblong cavity, whose greatest length is from one side to the other.—Round it several tubercles may be remarked, for the insertion of ligaments.—Its body is convex, or a half-round behind; but flat before, for lodging the tendon of the long slexor of the thumb, which is tied down by ligamentous sheaths that are fixed on each side to the angle at the edge of this flat surface.—The lower end of this second bone has two lateral round protuberances, and a middle cavity, whose greatest extent of smooth surface is forwards.

The articulation and motion of the upper end of this fecond bone is as singular as that of the former.—For its cavity being joined to the round head of the sirst bone, it would seem at first view to enjoy motion in all directions; yet, because of the strength of its lateral ligaments, oblong sigure of the joint itself, and mobility of the sirst joint, it only allows slexion and extension;

and these are generally much confined.

The third-bone of the thumb is the smallest, with a large base, whose greatest extent is from one side to the other.

other.—This base is formed into two cavities and a middle protuberance, to be adapted to the pulley of the former bone.—Its body is rounded behind; but is slatter than in the former bone, for sustaining the nail.—It is slat and rough before, by the insertion of the slexor tertii internodii.—This bone becomes gradually smaller, till near the lower end, where it is a little enlarged, and has an oval scabrous edge.

The motion of this third bone is confined to flexion

and extension.

The orderly disposition of the bones of the fingers into three rows, has made them generally obtain the name of three phalanges (m)—All of them have half-bound convex surfaces, covered them with an aponeurosis, formed by the tendons of the extensors, lumbricales, and interossei, and placed directly backwards, for their greater strength; and their slat concave part is forwards, for taking hold more surely, and for lodging the tendons of the flexor muscles.—The ligaments for keeping down these tendons are fixed to the angles that are between the convex and concave sides.

The bones of the first phalanx (n) of the fingers answer to the description of the second bone of the thumb; only that the cavity in their base is not so oblong; nor is their motion on the metacarpal bones so much confined: For they can be moved laterally or circularly; but have no rotation, or a very small degree of it, round their axis.

Both the ends of this first phalanx are in a cartilaginous state at the birth; and the upper one is afterwards affixed in form of an epiphyse.

The fecond bone (0) of the fingers has its base formed into two lateral cavities, and a middle protuberance;

<sup>(</sup>m) Scytalidæ, internodia, scuticula, agmina, acies, condyli are

<sup>(</sup>n) Προχονδυλοι. (o

<sup>(</sup>o) KovSunos.

while the lower end has two lateral protuberances and a middle cavity; therefore it is joined at both ends in the same manner, which none of the bones of the thumb are.

This bone is in the same condition with the former in children.

The third bone (p) differs nothing from the defcription of the third bone of the thumb, excepting in the general distinguishing marks; and therefore the fecond and third phalanx of the singers enjoy only flexion and extension.

The upper end of this third phalanx is a cartilage in a ripe child; and is only an epiphyse after, till the full growth of the body.

All the difference of the phalanges of the several fingers consists in their magnitude.—The bones of the middle finger (q) being the longest and largest,—those of the fore-finger (r) come next to that in thickness, but not in length, for those of the ring-finger (s) are a little longer. The little finger (t) has the smallest bones. Which disposition is the best contrivance for holding the largest bodies; because the longest singers are applied to the middle largest periphery of such substances as are of a spherical figure (u).

The uses of all the parts of our superior extremities are so evident in the common actions of life, that it is needless to enumerate them here; and therefore I shall proceed to the last part of the skeleton. Only, lest I should seem to have forgot the small bones at the joints of the hand, I desire now to refer to the description of them,

(ρ) Μετακονδύλοι, ριζωνεχια.

(r) Δεκτικός, Indicator, λιχανός, demonstrativus, falutaris.

<sup>(</sup>q) Karatuyev, opaxxelos, Infamis, impudicus, verpus, famolus, oblecenus.

<sup>(3)</sup> Ίατρικος, παραμισος, δακτυλιωτής, ιπιβατής, Annularis, medicus, cordis digitus.

<sup>(</sup>t) Μυωψ, ωτιτης, Auricularis, minimus.
(z) Galen. de ufu part. lib. 1. cap. 24-

them, under the common title of fesamoid bones, which I have placed after the bones of the feet.

SECT. IV. OF THE INFERIOR EXTREMITIES.

THE INFERIOR EXTREMITIES depend from the acetabula of the offa innominata; and are commonly divided into three parts, viz. the thigh, leg, and foot.

§ 1. The THIGH.

THE thigh (x) has only one bone; which is the longest of one body, and the largest and strongest of any of the cylindrical bones. The situation of it is not perpendicular; for the lower end is inclined confiderably inwards; fo that the knees are almost contiguous, while there is a confiderable diffance between the thighbones above; which is of good use to us, fince sufficient space is thereby left for the external parts of generation, the two great cloacæ of urine and fæces, and for the large thick muscles that move the thigh inwards. At the fame time this fituation of the thighbones renders our progression quicker, surer, straighter, and in less room: for, had the knees been at a greater distance from each other, we must have been obliged to describe some part of a circle with the trunk of our body in making a long step; and when one leg was raifed from the ground, our centre of gravity would have been too far from the base of the other, and we should consequently have been in danger of falling; so that our steps would neither have been straight nor firm, nor would it have been possible to walk in sa narrow path, had our thigh-bones been otherwise placed. In confequence, however, of the weight of the body bearing fo obliquely on the joint of the knee by P 3 this

<sup>(</sup>x) Mnpov, Femen, coxa, agis, anchæ os, crus, femur.

this situation of the thigh-bones, weak ricketty children become in-knee'd.

The upper end of the thigh-bone is not continued in a straight line with the body of it, but is set off oblique. ly inwards and upwards, whereby the distance here between these two bones, at their upper part is considerably increased.—This end is formed into a large fmooth round head (y), which is the greater portion of a fphere unequally divided .- Towards its lower internal part a round rough spongy pit is observable, where the strong ligament, commonly, but unjustly, called the round one, is fixed, to be extended from thence to the lower internal part of the receiving cavity, where it is confiderably broader than near to the head of the thigh bone. The fmall part below the head, called the cervix, of the os femoris, has a great many large holes, into which the fibres of the strong ligament, continued from the capfular, enter, and are thereby furely united to it; and round the root of the neck, where it rifes from the bone, a rough ridge is found, where the capfular ligament of the articulation itself is connected.—Below the back-part of this root, the large unequal protuberance, called trochanter major (z), stands out; the external convex part of which is distinguished into three different surfaces; whereof the one on the fore-part is fcabrous and rough, for the infertion of the glutæus minimus; the superior one is smooth, and has the glutæus medius inserted into it; and the one behind is made flat and fmooth, by the tendon of the gluteeus maximus paffing over it.—The upper edge of this process is sharp and pointed at its back-part, where the glutæus medius is fixed; but forwards it is more obtuse, and has two superficial pits formed in it: Into the superior of these the pyriformis is implanted; and the obturator internus and gemini are fixed into the lower one. - From the backmost pro-

(y) Vertebrum.

<sup>(</sup>z) TANTOS, Rotator natis, malum granatum testiculorum.

minent part of this great trochanter, a rough ridge runs backwards and downwards, into which the quadratus is inferted.—In the deep hollow, at the internal upper fide of this ridge, the obturator externus is implanted.

—More internally, a conoid process, called trochanter minor (a), rises for the infertion of the musculus psoas and iliacus internus; and the pectineus is implanted into a rough hollow below its internal root. The muscles inferted into these two processes being the principal instruments of the rotatory motion of the thigh, have occasioned the name of trochanters to the processes.—The tendons that are fixed into or pass over the great trochanter, cause bruises by falls on this part to be attended with great pain and weakness of the limb, which

generally remain long.

The body of the os femoris is convex on the forepart, and made hollow behind, by the action of the muscles that move it and the leg, and for the conveniency of fitting, without bearing too much on these muscles; and probably the weight of the legs depending from the thighs in that posture contributes to this curvature. The fore-part of the thigh-bone is a little flatted above by the beginning of the cruræus muscle, as it is also below by the same muscle and the rectus. -Its exernal furface is likewise made flat below by the vastus externus, where it is separated from the former by an obtuse ridge. The valtus internus depresses a little the lower part of the internal furface.—The posterior concave surface has a ridge rising in its middle, commonly called linea afpera, into which the triceps is inferted, and the fhort head of the biceps flexor tibiæ rifes from it .- At the upper part of it the medullary veffels enter by a fmall hole that runs obliquely upwards. A little above which there is a rough fossa or two, where the tendon of the glutæus maximus is fixed .-The lower end of the linea aspera divides into two, which descend towards each fide. The two vasti muscles

muscles have part of their origin from these ridges; and the long tendon of the triceps is fixed to the internal, by means of part of the fascia aponeurotica of the thigh.—Near the beginning of the internal ridge, there is a discontinuation of the ridge, where the crural artery passes through the aponeurosis.—Between these two rough lines, the bone is made slat by the large blood-vessels and nerves which pass upon it; and near the end of each of these ridges a small smooth protuberance may often be remarked, where the two heads of the external gastrocnemius muscle take their rise, and where sesamoid bones are sometimes sound (b); and from the fore-part of the internal tubercle a strong

ligament is extended to the infide of the tibia.

The lower end of the os femoris is larger than any other part of it, and is formed into a great protuberance on each fide, called its condyles; between which a confiderable cavity is found, especially at the backpart, in which the crural veffels and nerves lie immerfed in fat.—The internal condyle is longer than the external, which must happen from the oblique position of this bone, to give less obliquity to the leg. Each of these processes seems to be divided in its plain smooth furface. The mark of division on the external is a notch, and on the internal a small protuberance. fore-part of this division, on which the rotula moves, is formed like a pulley, the external fide of which is highest. Behind, there are two oblong large heads, whose greatest extent is backwards, for the motion of the tibia: and from the rough cavity between them, but near to the base of the internal condyle, the strong ligament, commonly called the cross one, has its rife .-A little above which a protuberance gives infertion to the tendon of the triceps. The condyles, both on the outer and inner fide of the knee are made flat by the muscles passing along them. On the back-part of the internal, a flight depression is made by the tendons of the

<sup>(</sup>b) Vefal. lib. 1. cap. 28, & 30.

the gracilis and fartorius; and on the external such another is formed by the biceps slexor cruris; behind which a deep soffa is to be observed, where the poplitæus muscle has its origin. From the tubercle immediately before this cavity, a strong ligament goes out to the upper part of the sibula.—Round this lower end of the thigh-bone, large holes are found, into which the ligaments for the security of the joint are fixed, and blood-vessels pass to the internal substance of the bone.

All the processes of the femur are cartilaginous in new-born children; and afterwards become small apo-

physes, with large epiphyses.

f

0

C

e

of

C

The thigh-bone being articulated above with the acetabulum of the offa innominata, which affords its round head a fecure and extensive play, can be moved to every fide; but is restrained in its motion outwards by the high brims of the cavity, and by the round ligament; for otherwise the head of the bone would have been frequently thrust out at the breach of the brims on the infide, which allows the thigh to move confiderably inwards.—The body of this bone enjoys little or no rotatory motion, though the head most commonly moves round its own axis; because the oblique progress of the neck and head from the bone is fuch, that the rotatory motion of the head can only bring the body of the bone forwards and backwards. Nor is this head, as in the arm, ever capable of being brought to a straight direction with its body: so far, however, as the head can move within the cavity backwards and forwards, the rest of the bone may have a partial rotation.—When the thigh-bone resists the actions of its muscles more than the trunk of the body can then do, at in standing, these muscles have their effect on the trunk, causing it to bend forward, raising it up, inclining it to the one or the other fide, twilting it obliquely, &c. which the rolling of the acetabula of the offa innominata on the round heads of the thighbones,

bones is well fitted for.—The os femoris is articulated below to the tibia and rotula in the manner afterwards to be described.

The nearness of the small neck to the round head of the thigh-bone, and its upper end being covered with very thick muscles, make greater difficulty in distinguishing between a luxation and fracture here, than in any other part of the body.

# § 2. The LEG.

THE leg (c) is composed, according to the common account, of two bones, tibia and fibula, though it seems to have a very good title to a third, the rotula; which bears a strong analogy to the olecranon of the ulna, and moves always with the other two.

# TIBIA.

Tibia (d), so called from its resemblance to an old musical pipe or flute, is the long thick triangular bone, situated at the internal part of the leg, and continued

in almost a straight line from the thigh-bone.

The upper end of the tibia is large, bulbous, and spongy, and is divided into two cavities by a rough irregular protuberance (e), which is hollow at its most prominent part, as well as before and behind. The amerior of the two ligaments that compose the great cross one is inserted into the middle cavity, and the depression behind receives the posterior ligament.—The two broad cavities at the sides of this protuberance are not equal: for the internal is oblong and deep, to receive the internal condyle of the thigh-bone; while the external is more superficial and rounder, for the external condyle.—In each of these two cavities of a

(c) Kunun, Crus, tibia.

 <sup>(</sup>d) Προκνημιον, οντικνημιον, Focile majus, arundo major, canna major, canna domestica cruris.
 (e) Διανρυσις, εξοχη νευροχονκρωδης, Tuber, tuberculum.

recent subject, a semilunar cartilage is placed, which is thick at its convex edge, and becomes gradually thinner towards the concave or interior edge. The middle of each of these cartilages is broad, and the ends of them turn narrower and thinner as they approach the middle protuberance of the tibia. The thick convex edge of each cartilage is connected to the capfular and other ligaments of the articulation, but fo near to their rife from the tibia, that the cartilages are not allowed to change place far; while the narrow ends of the cartilages becoming almost ligaments, are fixed at the infertion of the strong cross ligament into the tibia, and feem to have their substance united with it; therefore a circular hole is left between each cartilage and the ligament, in which the most prominent convex part of each condyle of the thigh-bone moves. The circumference of these cavities is rough and unequal, for the firm connection of the ligaments of the joint. Immediately below the edge at its back-part, two rough flatted protuberances stand out: into the internal, the tendon of the semimembranosus muscle is inserted; and a part of the cross ligament is fixed to the external.— On the outfide of this last tubercle, a smooth slightlyhollowed furface is formed by the action of the poplitæus muscle.

Below the fore-part of the upper end of the tibia, a considerable rough protuberance (f) rises, to which the strong tendinous ligament of the rotula is fixed.—
On the internal side of this, there is a broad scabrous slightly-hollowed surface, to which the internal long ligament of the joint, the aponeurosis of the vastus internus, and the tendons of the seminervosus, gracilis, and sartorius, are fixed. The lowest part of this surface is therefore the place where the tibia ought to be sawed through in an amputation, so as not to have too long and troublesome a stump, and at the same time to preserve its motions, by saving the proper muscles.

muscles. Below the external edge of the upper end of the tibia, there is a flat circular furface, covered in a recent fubject with cartilage, for the articulation of the fibula; between which and the anterior knob there is a rough hollow, from which the tibialis anticus, and extensor digitorum longus, take their origin. From the fmooth flat furface, a ridge runs obliquely downwards and inwards, to give rife to part of the folæus, tibialis posticus, and flexor digitorum longus, and infertion to the aponeurofis of the femimembranofus which covers the poplitæus, and to some of the external fibres of this last-named muscle. At the inside of this ridge an oblique plain furface is left, where the greatest part of the musculus poplitæus is inserted. The remaining body of the tibia is triangular. The anterior angle is very sharp, and is commonly called the spine or shin(g). This ridge is not straight; but turns first inwards, then outwards, and lastly inwards again. The plain internal fide is smooth and equal, being little subjected to the actions of muscles; but the external side is hollowed above by the tibialis anticus, and below by the extenfor digitorum longus and extenfor pollicis longus. The two angles behind these sides are rounded by the action of the muscles; the posterior side comprehended between them is not fo broad as those already mentioned, but is more oblique and flatted by the action of the tibialis posticus and flexor digitorum longus.-Some way above the middle of the bone, the internal angle terminates, and the bone is made round by the pressure of the musculus solæus. Near to this the pasfage of the medullary veffels is feen flanting obliquely downwards.

The lower end of the tibia is made hollow, but so as a small protuberance rises in the middle. The internal side of this cavity, which is smooth, and, in a recent subject, is covered with cartilage, is produced into a considerable process, commonly named malleolus inter-

nus

<sup>(</sup>g) Axavaa, Spina, crea, linea prima tibiæ, angulus acutus.

nus (b); the point of which is divided by a notch, and from it ligaments are sent out to the foot. We ought to observe here, that this internal malleolus is situated more forwards than the internal condyle of the upper end of this bone; which is necessary to be remembered in reducing a fracture of the leg (i). The external side of this end of the tibia has a rough irregular semi-lunar cavity formed in it for receiving the lower end of the sibula. The posterior side has two lateral grooves, and a small middle protuberance. In the internal depression, the tendons of the musculus tibialis posticus and slexor digitorum longus are lodged; and in the external, the tendon of the slexor longus pollicis plays.—

From the middle protuberance, ligamentous sheaths go out, for tying down these tendons.

The articulations and motions of the tibia shall be explained, after all the three bones of the leg are described.

Both the ends of the tibia are cartilages at birth, and become afterwards epiphyses.

#### FIBULA.

FIBULA (k) is the small long bone, placed on the outside of the leg, opposite to the external angle of the tibia; the shape of it is irregularly triangular.

The head of the fibula has a superficial circular cavity formed on its inside, which, in a recent subject, is covered with a cartilage, but so closely connected to the tibia by ligaments, as to allow only a small motion backwards and forwards. This head is protuberant and rough on its outside, where a strong round ligament and the musculus biceps are inserted; and, below the back-part of its internal side, a tubercle may

<sup>- (</sup>h) Σφυρον, πεσον, Talus, clavicula, clavilla interior, clavilla do-mestica.

<sup>(</sup>i) Winflow, Exposition anatomique des os secs, § 865.

<sup>(</sup>k) Hagaxynatov, Berone, focile minus, atundo minor, canna minor cruris, sura, radius.

be remarked, that gives rife to the strong tendinous

part of the folæus muscle.

The body of this bone is a little crooked inwards and backwards: which figure is owing to the actions of the muscles; but is still further increased by nurses, who often hold children carelessly by the legs. sharpest angle of the fibula is forwards; on each fide of which the bone is confiderably, but unequally, depreffed by the billies of the feveral muscles that rise from or act upon it; and in old people, these muscles make distinct sinuosities for themselves- The external surface of the fibula is depressed obliquely from above downwards and backwards by the two peronæi. internal furface is unequally divided into two narrow longitudinal planes, by an oblique ridge extended from the upper part of the anterior angle, to join with the lower end of the internal angle. To this ridge the ligament stretched between the two bones of the leg is connected. The anterior of the two planes is very narrow above, where the extensor longus digitorum and extensor longus pollicis arise from it; but is broader below, where it has the print of the nonus Vefalii. The posterior plane is broad and hollow, giving origin to the larger share of the tibialis posticus. The internal angle of this bone has a tendinous membrane fixed to it, from which fibres of the flexor digitorum longus take their rife. The posterior surface of the fibula is the plainest and smoothest, but is made flat above by the folæus, and is hollowed below by the flexor pollicis longus. In the middle of this furface, the canal for the medullary veffels may be feen flanting downwards.

I have taken particular notice of the entry and direction of the medullary vessels of the large bones of the extremities (1); because, in several chirurgical cases, a surgeon who is ignorant of this may do mischief to his patient. Thus, for example, if these vessels are opened very near to their entry into the bone, or while

they

<sup>(1)</sup> Havers, Ofteolog. Nov. difc. 1. p. 59-

they are in the oblique passage through it, an obstinate hæmorrhagy may enfue: for the arteries being connected to the bony passage, styptics, and other like corrugators, are vainly applied; compressing instruments can do no fervice, and ligatures cannot be employed. There feems to be a particular defign in the contrivance of these canals; those in the os humeri, tibia, and fibula, running obliquely downwards from their external entry; whereas in the radius, ulna, and os femoris. they flant upwards, whereby the arteries and nerves which are fent into these three last bones, must fuffer a confiderable reflexion before they come at the cancelli. The reason of this diversity may perhaps be, that the arteries, which are so small within the bones as to have no strong contractile propelling force in their coats, and where they are not affifted by the action of any moving neighbouring organ, should have, at least in their pasfage through the bone, a favourable descent for their liquids: which, it is evident, they have in the defcending oblique passages formed for them in the first class of bones, to wit, the os humeri, tibia, and fibula, which are generally depending; and they also most frequently acquire the like advantage in the radius. ulna, and os femoris; because the hand, in the most natural posture, is higher than the elbow: and when we fit or lie, the lower end of the thigh-bone comes to be at least as high raised as the upper. In standing and walking, or when the arms are moved, the blood must indeed afcend as it passes through the bones of the forearm and thigh; but the pressure of the muscles, then in action, on the vessels, before they enter the bones, is fufficient to compensate the disadvantage of their courfe. This reasoning seems to be still enforced, by observing, that this passage is always nearer the upper than the lower ends of these bones.

The lower end of the fibula is extended into a spongy oblong head; on the inside of which is a convex, irregular, and frequently a scabrous, surface, that is recei-

ved by the external hollow of the tibia, and fo firmly joined to it by a very thin intermediate cartilage and strong ligaments, that it scarce can move.—Below this the fibula is stretched out into a coronoid process, that is smooth, covered with cartilage on its internal side. and is there contiguous to the outfide of the first bone of the foot, the astragalus, to secure the articulation. This process, named malleolus externus, being fituated farther back than the internal malleolus, and in an oblique direction, obliges us naturally to turn the forepart of the foot outwards (m). At the lower internal part of this process, a spongy cavity for mucilaginous glands may be remarked; from its point ligaments are extended to the astragalus, os calcis, and os naviculare, bones of the foot; and from its infide short strong ones go out to the astralagus. On the back-part of it a finuofity is made by the tendons of the peronæi When the ligament extended over these tenmuscles. dons from the one fide of the depression to the other is broke, stretched too much, or made weak by a sprain, the tendons frequently start forwards to the outside of the fibula.

The conjunction of the upper end of the fibula with the tibia is by plain surfaces tipped with cartilage; and at its lower end the cartilage seems to glue the two bones together; not, however, so firmly in young people, but that the motion at the other end of such a long radius is very observable.——In old subjects I often see the two bones of the leg grown together at their lower ends.

The principal use of this bone is to afford origin and insertion to muscles; the direction of which may be a little altered on proper occasions, by its upper part shuffling backwards and forwards.—It likewise helps to make the articulation of the foot more secure and firm.—The ends of the tibia and sibula being larger than their middle, a space is here left, which is filled

<sup>(</sup>m) Winflow, Memoires de l'acad. des sciences, 1722.

up with fuch another ligament as I described extended between the bones of the fore-arm; and which is also discontinued at its upper part, where the tibialis anticus immediately adheres to the solæus and tibialis positicus; but every where else it gives origin to muscular fibres (n).

Both the ends of this bone are cartilaginous in a ripe child, and assume the form of appendices before they

are united to its body.

### ROTULA.

ROTULA (6) is the small flat bone situated at the. fore-part of the joint of the knee.—Its shape resembles the common figure of the heart with its point downwards.—The anterior convex furface of the rotula is pierced by a great number of holes, into which fibres of the strong ligament that is spread over it enter .-Behind, its furface is smooth, covered with cartilage, and divided by a middle convex ridge into two cavities, of which the external is largest; and both are exactly adapted to the pulley of the os femoris, on which they are placed in the most ordinary unstraining postures of the leg: but when the leg is much bended, the rotula descends far down on the condyles; and when the leg is fully extended, the rotula rifes higher in its upperpart than the pulley of the thigh bone. The plain smooth surface is surrounded by a rough prominent edge, to which the capfular ligament adheres: Below, the point of the bone is scabrous, where the strong tendinous ligament from the tubercle of the tibia is fixed. The upper horizontal part of this bone is flatted and unequal, where the tendons of the extensors of the leg are inserted.

The substance of the rotula is cellular, with very Vol. I. Q thin

(n) Weitbrecht, Syndefmolog. p. 156.

<sup>(</sup>ο) 'Επιμυλις, μυλακεις, κογχος, επιγουατις, πλανιστεδεον, Patella, monla, genu, scutiforme os, cartilaginosum, disciforme, oculus genu,

thin external firm plates: but then these cells are fo small, and such a quantity of bone is employed in their formation, that scarce any bone of its bulk is so strong. Besides, it is covered all over with a thick ligament (as it was observed that this fort of bones generally is). to connect its fubstance, and is moveable to one fide or other: therefore it is fufficiently ftrong to refift the ordinary actions of the large muscles that are inserted into it, or any common external force applied to it; while a fixed process, such as the olecranon, would not have been sufficient to bear the whole weight of our bodies, which frequently falls on it, and would have hindered the rotatory motion of the leg. Notwithstanding these precautions to preferve this bone from fuch injuries. yet I have feen a transverse fracture in it, when, by the report of the patient, and of the people about him, and by the want of fwelling, discolouring, or other mark of bruise or contusion, it was plain the bone was broken by the violent straining effort of the muscles (p). Tho' my patient recovered the use of the joint of the knee, vet I think it reasonable to believe, that this fort of fracture is commonly attended with difficulty of motion after the broken parts of the rotula are reunited; because the callous matter probably extends itself into the cavity of the joint, where it either grows to some of the parts, or makes fuch an inequality on the furface of this bone, as does not allow it to perform the necessary motions on the condyles of the femur (q).

At the ordinary time of birth, the rotula is entirely cartilaginous, and scarcely assumes a bony nature so

foon as most epiphyses do.

(q) Paré, liv. 15. cap. 22.

The parts which conflitute the joint of the knee being now described, let us examine what are its motions, and how performed.—The two principal motions are flexion and extension. In the former of these, the leg may be brought to a very acute angle with the thigh,

<sup>(</sup>p) See Ruysch. Observ. anat.-chirurg. obs. 3.

by the condyles of the thigh-bones being round and made smooth far backwards. In performing this, the rotula is pulled down by the tibia. When the leg is to be extended, the rotula is drawn upwards, confequently the tibia forwards, by the extensor muscles; which, by means of the protuberant joint, and of this thick bone with its ligament, have in effect the chord with which they act fixed to the tibia at a confiderable angle, therefore act with advantage; but are restrained from pulling the leg farther than to a straight line with the thigh, by the posterior part of the cross ligament, that the body might be supported by a firm perpendicular column: For at this time the thigh and leg are as little moveable in a rotatory way, or to either fide, as if they were one continued bone -But when the joint is a little bended, the rotula is not tightly braced, and the posterior ligament is relaxed; therefore this bone may be moved a little to either fide, or with a final! rotation in the superficial cavities of the tibia; which is done by the motion of the external cavity backwards and forwards, the internal ferving as a fore of axis (r). Seeing, then, one part of the cross ligament is situated perpendicularly, and the posterior part is stretched obliquely from the internal condyle of the thigh outwards, that posterior part of the cross ligament prevents the leg's being turned at all inwards; but it could not hinder it from turning outwards almost round, was not that motion confined by the lateral ligaments of this joint; which can yield little. This rotation of the leg " outwards is of good advantage to us in croffing our legs, and turning our feet outwards, on feveral neceffary occasions; though it is altogether fit this motion should not be very large, to prevent frequent luxations here — While all these motions are performing, the part of the tibia that moves immediately on the condyles is only fo much as is within the cartilaginous

U

re

1,

by-

<sup>(</sup>r) Winslow, Exposition anatomique du corps humain, traité des

rings, which by the thickness on their outsides make the cavities of the tibia more horizontal, by raising their external side where the surface of the tibia slants downwards. By this means the motions of this joint are more equal and steady than otherwise they would have been. The cartilages being capable of changing a little their situation, are sit for doing this good office in the different motions and postures of the member, and likewise contribute to make the motions larger and quicker.

On account of the very large furface of the bones forming the joint of the knee, and the many strong ligaments connecting them, luxations seldom happen here. But these very ligaments, the aponeurosis passing over this joint, the quantity of fat and mucilaginous glands necessary for lubricating it, make it more subject to white swellings, dropsies, and such other disorders, than any other joint of the body.

# § 3. The FOOT.

THE foot is divided, as well as the hand, into three parts, viz tarfus, metatarfus, and toes: In the description of which the several surfaces shall be named, according to their natural situation, viz. the broad of the foot shall be called superior; the sole, inferior; the side on which the great toe is, internal; that where the little toe is, external.

### TARSUS.

THE tarfus (t) confilts of feven spongy bones; to wit, the altragalus, os calcis, naviculare, cuboides, curelforme externum, cuneiforme medium, and cuneiforme internum

The aftragalus is the uppermost of these bones.—
The os galeis is below the astragalus, and is considerably prominent backwards beyond the other bones, to form

form the heel.—The os naviculare is in the middle of the internal fide of the tarfus.—The os cuboides is the most external of the row of four bones at its fore part.
—The os cuneiforme externam is placed at the inside of the cuboid.—The cuneiforme medium is between the external and internal cuneiform bones; and the internal cuneiform is put at the internal side of the foot.

That the description of these bones may not be immoderately swelled with repetition, I desire, once for all, to observe, That wherever a ridge is mentioned, without a particular use assigned, a ligament is understood to be fixed to it; or where a spongy rough cavity, depression, or sossa, is remarked, without naming its use, a ligament is inserted, and mucilaginous glands are lodged: for such will occur in the detail of each of these bones.

Astragalus. The upper part of the astragalus (t) is formed into a large smooth head (u), which is slightly hollowed in the middle; and therefore refembles a superficial pulley, by which it is sitted to the lower end of the tibia. The internal side of this head is slat and smooth, to play on the internal malleolus.—The external side has also such a surface, but larger, for its articulation with the external malleolus. Round the base of this head there is a rough fossa; and immediately before the head, as also below its internal smooth surface, we find a considerable rough cavity.

The lower furface of the astragalus is divided by an irregular deep rough fossa; which at its internal end is narrow, but gradually widens as it stretches obliquely outwards and forwards. The smooth surface, covered with cartilage, behind this fossa, is large, oblong, extended in the same oblique situation with the sossa, and concave for its conjunction with the os calcis. The back-part of the edge of this cavity is produced into

(u) Tireveni.

<sup>(</sup>t) 'Acrese, Talus, ballistæ os, malleolus, chaib, quatrio, os teffaræ, claviculæ, nuciforme,

two sharp-pointed rough processes; between which is a depression made by the tenden of the flexor pollicis longus.——The lower furface before the fossa is convex, and composed of three distinct smooth planes.— The long one behind, and the exterior or shortest, are articulated with the heel-bone; while the internal, which is the most convex of the three, rests and moves upon a cartilaginous ligament, that is continued from the calcaneum to the os fcaphoides. Without which ligament the astragalus could not be sustained; but would be preffed out of its place by the great weight it supports, and the other bones of the tarfus would be separated. Nor would a bone be fit here, because it must have been thicker than could conveniently be allowed; otherwife it would break, and would not prove fuch an easy bending base, to lessen the shock which is given to the body in leaping, running, &c.

The fore-part of this bone is formed into a convex oblong smooth head, called by some its praces, which is received by the os naviculare. Round the root of this head, especially on the upper surface, a rough softa

may be remarked.

The astragalus is articulated above to the tibia and fibula, which together form one cavity. Though in this articulation the bones have prominences and cavities, fo small as might-allow motions in all directions, yet the flexion and extension are the most considerable, the other motions being confined by the malleoli, and by the strong ligaments which go out from the points of these processes to the altragalus and os calcis.— When the foot is bended, fo far as it is commonly when we stand, no lateral or rotatory motion is allowed in this joint; for then the head of the astragalus is sunk deep between the mallcoli, and the ligaments are tenfe: but when the foot is extended, the astragalus can move a little to either fide, and with a finall rotation. By this contrivance the foot is firm, when the weight of the body is to be supported on it; and when a foot

is raised, we are at liberty to direct it more exactly to the place we intend next to step upon. The astragalus is joined below to the os calcis; and before to the os naviculare, in the manner to be explained when these bones are described.

A confiderable share of this bone is offified in a newborn infant.

Behind, it is formed into a large knob, commonly called the beel: the furface of which is rough behind, where the tendo Achillis is inferted into it; and above it, it is hollow and spongy. Farther forwards, on the upper surface of the calcaneum, there is an irregular oblong smooth convexity, adapted to the concavity at the back-part of the astragalus: and beyond this a narrow soffa is seen, which divides it from two small concave smooth surfaces, that are joined to the fore-part of the astragalus.—Behind the posterior of these smooth surfaces, which is the largest, a small sinuosity is made by the tendon of the slexor digitorum longus; at the fore-part of which a small rough protuberance appears, that gives rise to the musculus extensor digitorum brevis.

The external fide of this bone is flat, with a superficial fossa running horizontally, in which the tendon of the musculus peronæus longus is lodged. The internal fide of the heel-bone is hollowed, for lodging the origin of the massa cornea Jac. Sylvii, and for the safe passage of tendons, nerves, and arteries. Under the side of the internal smooth concavity, a particular groove is made by the tendon of the slexor pollicis longus; and from the thin protuberance on this internal side the cartilaginous ligament that supports the astragalus, goes out to the os naviculare; on which ligament, and on the edge of this bone to which it is fixed, the groove is formed for the tendon of the slexor digitorum profundus.

Q 4 The

<sup>(</sup>v) Os calcis \*\*\* calcar pedis.

The lower furface of this bone is pressed slat at the back-part, by the weight of our bodies; and immediately before this plane, there are two tubercles, from the internal of which the musculus abductor pollicis, slexor digitorum sublimis, as also part of the aponeurosis plantaris, and of the abductor minimi digiti, have their origin; and the other part of the abductor minimi digiti and aponeurosis plantaris rises from the external. Before these protuberances this bone is concave, for lodging the slexor muscles; and at its fore-part we may observe a rough depression, from which, and a tubercle behind it, the signment goes out that prevents this bone from being separated from the os cuboides.

The fore-part of the os calcis is formed into an oblong pulley-like fmooth furface, which is circular at its upper external end, but is pointed below. The fmooth

furface is fitted to the os cuboides.

Though the furfaces by which the astragalus and os calcis are articulated, seem sit enough for motion; yet the very strong ligaments by which these bones are connected, prevent it, and render this principal part of our base, which rests on the ground, to wit, the os calcis, sirm.

A large share of the heel-bone is offisied at the ordinary time of birth, and the large knob appears after-

wards in form of an epiphyfe.

I.

ne

e-

m

S,

0-

re

ni l.

70

y

le

le

)-

h

S

t

r

bones of the middle toe, and of the toe next to the little one.—On the outfide of the os naviculare there is a semicircular smooth surface, where it is joined to the os cuboides.—The fore-part of this bone is all covered with cartilage, and is divided into three smooth planes, fitted to the three offa cuneiformia.

The os naviculare and astragalus are joined as a ball and socket; and the naviculare moves in all directions in turning the toes inwards, or in raising or depressing either side of the foot, though the motions are greatly restrained by the ligaments which connect this to the other bones of the tarfus. A weakness of these ligaments causes sometimes an unnatural turn of the forepart of the foot inwards.

The os naviculare is wholly cartilaginous in a newborn infant.

Os cuboides (y) is a very irregular cube.—Behind, it is formed into an oblong unequal concavity, adapted to the fore part of the os calcis. - On its internal fide, there is a small semicircular smooth cavity, to join the os naviculare.—Immediately before which, an oblong fmooth plane is made by the os cuneiforme externum. Below this the bone is hollow and rough.—On the internal fide of the lower furface, a round protuberance and fossa are found, where the musculus adductor pollicis has its origin. On the external fide of this fame furface, there is a round knob, covered with cartilage; immediately before which a fmooth foffa may be obferved, in which the tendon of the peronæus primus runs obliquely cross the foot; and on the knob the thin flat cartilage proper to this muscle plays; in place of which fometimes a bone is found:—More externally than the knob, a rough hollow is made, for the frong ligaments stretched betwixt this bone and the os calcis.—Before, the furface of the os cuboides is flat, smooth, and flightly divided into two planes, for fu**ftaining** 

<sup>(</sup>y) Πολυμορτον, Cubiforme, quadratum, grandinosum, varium, tessaræ, multiforme.

staining the os metatarsi of the little toe, and of the toe next to it.

The form of the back-part of the os cuboides, and the ligaments connecting the joint there with the os calcis, both concur in allowing little motion in this part.

The offification of this bone is scarcely begun at the

birth.

Os cuneiforme externum (z), if we regard its fituation or medium by its bulk, is much of the shape of a wedge, being broad and flat above, with long fides running obliquely downwards, and terminating in a sharp edge.—Ihe upper surface of this bone is an oblong fquare.—The one behind is nearly a triangle, but not complete at the inferior angle, and is joined to the os naviculare. The external fide is an oblong fquare divided as it were by a diagonal; the upper half of it is smooth, for its conjunction with the os cuboides: The other is a scabrous hollow, and in its superior anterior angle a small smooth impression is made by the os metatarfi of the toe next to the little one. The internal fide of this bone is also quadrangular, with the fore-part of its edge made flat and smooth by the os metatarsi of the toe next to the great one, and the back-part is also flat and fmooth where the os cuneiforme medium is contiguous to it. The fore-part of this bone is an oblong triangle, for fultaining the os metatarfi of the middle toe.

Os cuneiforme medium, or minimum, is still more exactly the shape of a wedge than the former.—Its upper part is square;—its internal side has a slat smooth surface above and behind, for its conjunction with the following bone; with a small rough solla below; and a considerable share of it is rough and hollow.—The external side is smooth and a little hollowed, where it is contiguous to the last described bone.—Behind, this bone is triangular, where it is articulated with the os navi-

<sup>(</sup>z) Chalcoideum externum.

naviculare; and it is also triangular at its fore-part, where it is contiguous to the os metatarsi of the toe

next to the great one.

Os cuneiforme maximum, or internum, differs from the two former in its fituation, which is more oblique than theirs.—Befides, its broad thick part is placed below, and the fmall thin point is above and outwards; while its under broad furface is concave, for allowing a fafe paffages to the flexor of the great toe. The furface of this os cuneiforme behind, where it is joined to the os naviculare, is hollow, fmooth, and of a circular figure below, but pointed above. The external fide confifts of two fmooth and flat furfaces, whose direction is nearly at right angles with each other. With the posterior, that runs obliquely from below forwards and upwards, the os cuneiforme minimum is joined; and with the anterior, whose direction is longitudinal, the os metatarfi of the toe next to the great one is connected.—The fore-part of this bone is fumilunar, but flat and fmooth, for fustaining the os metatarfi of the great toe.— The internal fide is fcabrous, with two remarkable tubercles below, from which the musculus abductor pollicis rifes, and the tibialis anticus is inferted into its upper part.

The three cunciform bones are all so secured by ligaments, that very little motion is allowed in any of them, and they are cartilaginous in a sectus of nine months.

These seven bones of the tarsus, when joined, are vex above, and leave a concavity below, for lodging safely the several muscles, tendons, vessels, and nerves, that lie in the sole of the soot. In the recent subject, their upper and lower surfaces are covered with strong ligaments which adhere firmly to them; and all the bones are so tightly connected by these and the other ligaments, which are fixed to the rough ridges and sosse mentioned in the preceding description of the particular bones, that, notwithstanding the many surfaces covered with cartilage, some of which are of the form of the very moveable articulations, no more motion is

here allowed, than only to prevent too great a shock of the sabric of the body in walking, leaping, &c. by falling on too solid a base; which, if it was one continued bone, would likewise be much more liable to be broken; and, in order to make our foot accommodate itself to the surfaces we tread on, by becoming more or less hollow, or by raising or depressing either side of it, as might be judged by what was said of the particular bones.

Sprains here occasion, as in the wrist, great pain and obstinate tumours, which too often cause carious bones.

## METATARSUS.

METATARSUS (z) is composed of five bones, which, in their general characters, agree with the metacarpal bones; but may be distinguished from them by the following marks: 1. They are longer, thicker, and stronger. 2. Their anterior round ends are not so broad, and are less in proportion to their bases.

3. Their bodies are sharper above and slatter on their sides, with their inferior ridge inclined more to the outside. 4. The tubercles at the lower parts of the round head are larger.

The first or internal metatarfal bone is easily distinguished from the rest by its thickness. The one next to it is the longest, and with its sharp edge almost perpendicular. The others are shorter and more oblique, as their situation is more external. Which general remarks, with the description I am now to give of each, may teach us to distinguish them from each other.

Os metatarsi policis is by far the thickest and strongest, as having much the greatest weight to sustain. Its base is oblong, irregularly concave, and of a semilunar sigure, to be adapted to the os cuneiforme maximum. The inferior edge of this base is a little prominent and rough,

<sup>(</sup>z) Erndor, mediov, Planta, planum, vestigium, folium, pectus, prz-cordium, pectusculum.

rough, where the tendon of the peronœus primus muscle is inserted. On its outside an oblique circular depression is made by the second metatarsal bone. Its round head has generally on its fore-part a middle ridge, and two oblong cavities, for the ossa sesand on the external side a depression is made by the

following bone.

With a triangular base supported by the os cuneiforme medium and the external side produced into a process; the end of which is an oblique smooth plane, joined to the os cuneiforme externum.—Near the internal edge of the base, this bone has two small depressions, made by the os cuneiforme maximum, between which is a rough cavity.—Farther forwards we may observe a smooth protuberance, which is joined to the foregoing bone.—On the outside of the base are two oblong smooth surfaces, for its articulation with the following bone; the superior smooth surface being extended longitudinally, and the inferior perpendicularly; between which there is a rough sossa.

Os metatarsi of the middle toe, is the second in length.

—Its base, supported by the os cuneiforme externum, is triangular, but slanting outwards, where it ends in a sharp-pointed little process; and the angle below is

not completed.

The internal fide of this base is adapted to the preceding bone; and the external fide has also two smooth surfaces covered with cartilage, but of a different sigure; for the upper one is concave, and, being round behind, turns smaller as it advances forwards; and the lower surface is little, smooth, convex, and very near the edge of the base.

Os metatarsi of the fourth toe, is near as long as the former, with a triangular flanting base joined to the os cuboides, and made round at its external angle; having one hollow smooth surface on the outside, where it is pressed upon by the following bone; and two on the

internal

internal fide, corresponding to the former bone; behind which is a long narrow furface impressed by the os cuneiforme externum.

Os metatarsi of the little toe, is the shortest, situated with its two slat sides above and below, and with the ridges laterally.—The base of it, part of which rests on the os cuboides, is very large, tuberous, and produced into a long-pointed process externally, where part of the abductor minimi digiti is sixed; and into its upper part the peronæus secundus is inserted. Its inside has a slat conoidal surface, where it is contiguous to the preceding bone.

When we stand, the fore-ends of these metatarsal bones, and the os calcis, are our only supporters; and therefore it is necessary they should be strong, and

should have a confined motion.

#### TOES.

THE bones of the toes are much akin to those of the thumb and fingers; particularly the two of the great toe are precisely formed as the two last of the thumb; only their position, in respect of the other toes, is not oblique; and they are proportionally much stronger, because they are subjected to a greater force; for they sustain the force with which our bodies are pushed forwards by the foot behind at every step we make; and on them principally the weight of the body is supported, when we are raised on our tip-toes.

The three bones in each of the other four toes, compared to those of the singers, differ from them in these particulars. They are less, and smaller in proportion to their lengths:—Their bases are much larger than their anterior ends:—Their bodies are more narrow above and below, and flatter on the sides. The first phalanx is proportionably much longer than the bones of the

fecond and third, which are very short.

Of the four, the toe next to the great one has the largest bones in all dimensions, and more externally the

next to it, have the fecond and third bones intimately united into one; which may be owing to their little motion, and the great pressure they are subject to.

The toes are of good use to us in walking; for, when the sole is raised, they bring our body, with its centre

of gravity, perpendicular to the advanced foot.

The bones of the METATARSUS and TOES, are in the fame condition in children as those of the metacarpus and fingers.

THE only bones now remaining to complete the defcription of the skeleton, are the small ones which are found at the joints of the singers and toes and in some other parts, called

#### OSSA SESAMOIDEA.

THESE are of very different figures and fizes, though they are generally faid to refemble the feed of the fefamum. They feem to me nothing elfe than the ligaments of the articulations, or the firm tendons of strong muscles, or both, become bony by the compression which they suffer. Thus the sesamoid bones at the beginning of the gastrocnemii muscles, are evidently composed of the tendinous fibres only. These, at the first joint of the great toe, are as plainly the same continued substance with the ligaments and the tendons of the adductor, flexor, brevis, and abductor.-That which is fometimes double at the fecond joint of that toe is part of the capfular ligament; and if we enumerate the other fefamoid bones that are at any time found, we may observe all of them formed in this man-Their number, figure, fituation, and magnitude, are so uncertain, that it were in vain to infift on the differences of each; and therefore I shall only in general remark.

1. That wherever the tendons and ligaments are firmest, the actions of the muscles strongest, and the compression

compression greatest, there such bones are most com-

2. That, cæteris paribus, the older the subject is in which they are sought, their number is greater, and

their fize larger.

3. The more labour any person is inured to, he has, caterus paribus, the most numerous and largest offa se-famoidea.

However, as the two at the first joint of the great toe are much larger than any other, are early formed, and are seldom wanting in an adult, we may judge, that, besides the more forcible cause of their formation, there should also be some particular advantage necessary at this place, rather than elsewhere; which may possibly be, to allow the slexor muscles to send their tendons along this joint, secure from compression in the hollow between the two oblong sesamoid bones; while, by removing these tendons from the centre of motion, and giving them the advantage of an angle at their insertion, the force of the muscle is increased, and therefore the great superincumbent weight of our body in progression is more easily raised.

# APPENDIX.

### MARKS of a FEMALE SKELETON.

TO finish the description of the bones, is generally to conclude the ofteology: but, that no part of the fubject may be left untouched, I think it necessary to subjoin the distinguishing marks of the Male and Female skeletons; and have chosen to illustrate them principally in the latter; because women having a more delicate constitution, and affording lodging and nourishment to their tender fætuses till they have sufficient ftrength and firmness to bear the injuries of the atmosphere, and contact of other more solid substances, their bones are frequently incomplete, and always of a make in some parts of the body different from those of the robust male; which agree to the description already given, unless where the proper specialities of the semale were particulary remarked; which could not be done in all places where they occur, without perplexing the order of this treatife: Therefore I choose rather to sum them up here by way of appendix.

The causes of the following specialities of the semale bones may be reduced to these three: 1. A weak lax constitution. 2. A sedentary unactive life, increasing that constitution. 3. A proper frame for being mothers.

The bones of women are smaller in proportion to their length than those of men; because the force of their muscles is not so great, nor is such strong external force applied to them to prevent their stretching out in length.

The depressions, ridges, scabrous surfaces, and other inequalities made by the muscles, are not so conspicuous Vol. I.

in them; because their muscles are neither so thick nor strong, nor so much employed, to make so strong prints on their bones.

The os frontis is more frequently divided by a continuation of the fagittal future; which depends on the first and second general causes assigned above for the specialities in their bones, as will appear after reflecting on the account given formerly of the middle internal spine of this bone.

Their clavicles are less crooked; because their arms have been less foreibly pulled forwards; which, in our European women, especially those of distinction, is more

hindered by their garb.

Their sternum is more raised by long cartilages below, that the thorax might be there widened in some proportion to what it is shortened by the pressure upon

the diaphragm when they are with child.

The defect of bone, or the hole, in the middle of the sternum, is oftenest found in them; to allow the passage of the mammary vessels, say some. But, in my opinion, this is owing to a lax constitution, by which the offisication is not so soon completed as in men, where the action of the solids is vigorous, and the circulation of the sluids is brisk: for a much smaller hole might have served this purpose; and the branches of the internal mammary vessels which are sent to the external parts of the thorax, do not pass here, but between the cartilages of the ribs, before these are joined to the sternum.

The cartilago xiphoides, is oftener bifcurated in women than men, for the reason assigned in the preceding paragraph, viz. a less forcible power of ossistation.

The superior cartilages of the ribs sooner offify, to

support the weight of the mammæ.

The middle cartilages are more flat and broad by the

weight of the breafts.

The inferior cartilages are longer, for enlarging the cheft.

When

or

ts

i-

le.

ne

ð-

r-

13-

ur

re

e-

le

n

10

Te

1-

ne

re

n

ht

n-

al

10

ne

0.

0

le

11

Weak women, who have born many children when young, often have the vertebræ of their back bended forwards, and their sternum depressed, or become roundshouldered and flat-breakted (a), by the pressure and weight of the impregnated uterus, and by the itrong action of the abdominal muscles.

The os facrum is broader, and turned much more

backwards, for enlarging the pelvis.

The os coccygis is more moveable, and much less

bended forwards, to facilitate the birth.

The offa ilium are more hollow, and more reflected outwards, and confequently further removed from each other, in order to widen the lower part of their abdomen, and for the better support of the impregnated uterus.

The ridge on the upper part of the os pubis is larger in fuch women as have born children, being extended by the strong action of the musculi recti abdominis.

The cartilage between the two offa pubis, especially in women who have born children, is thicker than in men, by which the pelvis is more capacious in females.

The conjoined furfaces of the offa pubis, and of the offa innominata and facrum, are less, the angle under the symphysis of the offa pubis is much larger, and the arches • formed below and behind by the offa ilium and ischium are wider, which, with the straighter os facrum, and more distant tubera ischii, leave a larger passage for the exclusion of the child in birth.

The great tuberofity of the offa ischium is flatter in women than in men, because it is more pressed upon in

the fedentary life which females enjoy.

In consequence of the pelvis of women being wider, the articulations of their thigh-bones mult be farther removed from each other; and therefore a larger space is left for the procreation and, birth of children (b); which

R 2

(b) Albin. De offib. 9 339.

<sup>(</sup>a) Cheselden, Anatomy, book 1. chap. 1.

which distance of the thighs may be one reason why women in running shuffle more from one side to the other than men, to preserve the centre of gravity of their bodies from falling too far to a side of the joint of the thigh that supports them when the other is raised, which would endanger their tumbling to the ground.

# SYSTEM OF ANATOMY.

# PART II.


CONTAINING!A DESCRIPTION OF THE

# HUMAN MUSCLES,

Chiefly as they appear on Dissection.

Together with their feveral Uses, and the Synonyma of the best Authors.

By JOHN INNES.



# ALEXANDER MONRO, M.D.

PROFESSOR of ANATOMY and MEDICINE in the University of Edinburgh.

SIR,

by many of your Pupils, to publish a short Description of the Muscles as they appear upon Dissection of the Human body, I have at last ventured to comply with their request. Your anxiety to promote the Science of Anatomy, and to encourage every thing that may be useful to the gentlemen who attend your Theatre, was the principal motive which induced me to undertake this task. I have no knowledge of the subject but what I derived from you. If, therefore, this Treatise, which you have never seen till I now present it to you, should communicate any advantage to the gentlemen under your care, it is to you alone they are indebted for the obligation. I am, Sir, with respect, gratitude, and esteem, your much obliged and very humble servant,

EDINBURGH, }

JOHN INNES.

### PREFACE.

Several full and accurate descriptions of the muscles have already been published. But their size and prolixity have rendered them of less value to the dissector than the small treatise of Dr Douglas, which was first published about the beginning of this century; and since that time has undergone various impressions, without receiving any improvement, excepting the addition of the synonyma from Albinus. It is therefore presumed, that a simple and concise description of the muscles, which should contain all the improvements of the moderns, is still wanting.

R 4

To

To class the muscles according to their uses, may do very well in a large work, or in describing their compound actions: But this method can never answer the purposes of dissection. To remedy this inconvenience, the muscles in the following treatise are described chiefly as they appear in

diffecting the human body.

The describing of the muscles according to their origins and insertions prevents much circumlocution. This is the method pursued by Mr Douglas; and wherever his description seemed tolerably accurate, they have been followed with little alteration. But Dr Douglas's book is peculiarly desective with regard to the muscles of the abdomen, back, and neck: in describing these, therefore, the method of Albinus has been preserved.

Those who have not opportunity, or are averse from undergoing the labour of dissecting, may derive considerable advantages from comparing the descriptions now given with the beautiful and correct tables of Albinus; and, to facilitate still more the study of these intricate organs, I have caused to be published eight of Albinus's tables, with concise explanations, on a small scale adapted for the pocket.

For the benefit of those who wish to examine the history of the muscles more minutely, the synonyma of the best authors are added; and, for the sake of brevity, the compound action of the muscles, and the origin and insertion of seve-

ral inconsiderable fibres, are omitted.

The reader will observe, that, in general, the muscles of one side only are described; because all the muscles of the body, with very few exceptions, have correspondent ones on the opposite side.

# A fhort ACCOUNT of the AUTHOR, And of his WRITINGS.

R JOHN INNES, author of the following Treatife, was born at Callart, an obscure village in the Highlands of Scotland. He came to Edinburgh at an early period of life, where he obtained the patronage and protection of Dr Alex-

A D-

Alexander Monro, professor of anatomy; who instructed him in the knowledge of the human hody, and in the art of disfection. When about the age of eighteen years, Dr Monro appointed him dissector to the anatomical theatre. The functions of this important and dissicult office he continued to perform with much reputation for near twenty years.

But his abilities were not confined to the dexterity of diffecting the most minute parts of the human frame. He deferibed the various organs with ease and with perspicuity. This happy talent attracted the notice of the students; and, at their solicitation, and by the approbation of Dr Monro, he opened an evening-course of anatomical demonstrations.

The number of pupils who annually attended these demonstrations afforded the best evidence of his abilities, and of the advantages derived from his labours. During his last course, he was attended by near two hundred students.

For some time before his death, he was troubled with an affection of the lungs, which terminated in a phthisis pulmonalis, and proved fatal to him on the 12th of January 1777.

On the 15th of the fame month, the following account of

him appeared in the public papers:

" Mr John Innes, at an early period of life, had been educated in the diffecting art: He made a rapid progress in his profession; and his genius and industry were rewarded with the privilege of giving private lectures for his own emo-The utility of his lessons were foon perceived. Numbers of students reforted to him for instruction; and all of them acknowledged the advantages they had received. At that stage of life when men are most capable of benefiting themselves, and of being useful to the public, death hurried him out of the world. He has given two small specimens of what was to be expected from his anatomical skill. In the course of last year, he published a short description of the Human Muscles as they appear on diffection, together with their several uses, and the synonyma of the best authors. The merit of this work was univerfally acknowledged. Some months after, he published, as a vade mecum for students, eight anatomical tables, containing the principal parts of the skeleton and muscles represented in the large tables of Albinus, with accurate explanations. These are all the monuments he has left by which the public is to judge of his ability. To his numerous friends and acquaintance, it is unnecessary to mention the warmth of his heart, or the integrity of his disposition."

# ADVERTISEMENT.

DURING the illness of which Mr INNES died, he put into my hands the first edition of his Description of the Muscles, with a few, chiefly verbal, corrections of it.

On perusing that work lately, at the request of the Bookseller, I have found it necessary to make a very considerable number of alterations in what relates to the description, as well as to the uses, of the Muscles.

Edin. Sep. 5. }

ALEX. MONRO.

### DESCRIPTION

OFTHE

# HUMAN MUSCLES.

#### CHAP I.

Muscles of the Teguments of the Cranium.

THE skin that covers the cranium is moved by a fingle broad digastric muscle, and one small pair.

### 1. OCCIPITO-FRONTALIS,

Arises sleshy from the transverse protuberant ridge near the middle of the os occipitis laterally, where it joins with the temporal bone; and tendinous from the rest of that ridge backwards, opposite to the lateral sinus; it arises after the same manner on the other side: From thence it comes straight forwards, by a broad thin tendon, which covers the upper part of the cranium at each side, as low down as the attollens aurem, to which it is connected, as also to the zygoma, and co-

vers a part of the aponeurofis of the temporal muscle; when it comes as far forwards as near the hair of the front, it becomes slessly, and descends with straight sibres.

Inferted into the orbicularis palpebrarum of each fide, and into the skin of the eye-brows, sending down a slessly slip between them, as far as the compressor naris and levator labii superioris alæque nasi.

Use. Pulls the skin of the head backwards; raises the eye-brows upwards; and, at the same time, it draws

up and wrinkles the skin of the fore-head.

Epicranius, Albinus.

Frontalis et occipitalis, Winflow.

#### 2. CORRUGATOR SUPERCILII,

Arises sleshy from the internal angular process of the os frontis, above the joining of the os nasi, and nasal process of the superior maxillary bone; from thence it runs outwards, and a little upwards.

Inserted into the inner and inferior fleshy part of the occipito-frontalis muscle, where it joins with the orbicularis palpebrarum, and extends outwards as far as the

middle of the fuperciliary ridge.

Use. To draw the eye-brow of that fide towards the other, and make it project over the inner canthus of the eye: When both act, they pull down the skin of the forehead, and make it wrinkle, particularly between the eye-brows.

Musculus Supercilii, Winflow.

Musculus Frontalis verus, seu Corrugator, Douglas.

I.

it

n

r

e

#### C H A P. II.

Of the Muscles of the EAR.

THE muscles of the ear may be divided into three classes, viz. the common, proper, and internal. The common may move the whole ear; the proper only affect the particular parts to which they are connected; and the internal, the small bones within the tympanum.

The common muscles are,

#### 1. ATTOLLENS AUREM,

Arises thin, broad, and tendinous, from the tendon of the occipito-frontalis, from which it is almost inseparable, where it covers the aponeurosis of the temporal muscle.

Inserted into the upper part of the ear, opposite to the

antihelix.

Use. To draw the ear upwards, and make the parts into which it is inferted tenfe.

Superior auris, Winflow.

### 2. ANTERIOR AURIS,

Arises thin and membranous near the posterior part of the zygoma.

Inserted into a small eminence on the back of the he-

lix, opposite to the concha.

Use. To draw this eminence a little forwards and upwards.

3. RE-

### 3. RETRAHENTES AURIS,

Arises, sometimes by three, but always by two diffinct small muscles, from the external and posterior part of the root of the mastoid process, immediately above the insertion of the sterno-cleido mastoid muscle.

Inserted into that part of the back of the ear which is opposite to the septum that divides the scapha and

concha.

Use. To draw the ear back, and stretch the concha. Posterior auris, Winslow.

The proper muscles are,

### 1. HELICIS MAJOR,

Arises from the upper and acute part of the helix anteriorly.

Inserted into its cartilage a little above the tragus.

Use. To depress that part from which it arises a little downwards and forwards.

#### 2. HELICIS MINOR,

Arises from the inferior and anterior part of the helix.

Inserted into the crus of the helix, near the fissure in the cartilage opposite to the concha.

Use. To contract the fiffure.

### 3. TRAGICUS,

Arises from the middle and outer part of the concha; at the root of the tragus, along which it runs.

Inferted into the point of the tragus.

Use. Pulls the point of the tragus a little forwards.

### 4. ANTITRAGICUS.

Arises from the internal part of the cartilage that supports the antitragus; and, running upwards, is

Inserted into the tip of the antitragus, as far as the inferior part of the antihelix, where there is a fissure in the cartilage.

Use.

Use. Turns the tip of the antitragus a little outwards, and depresses the extremity of the antihelix towards it.

#### 5. TRANSVERSUS AURIS,

Arises from the prominent part of the concha on the dorsum of the ear; the fibres not so fleshy as in the former.

Inferted opposite to the outer side of the antihelix.

Use. Draws the parts to which it is connected towards each other, and stretches the scapha and concha.

The muscles of the internal ear are three:

#### I. LAXATOR TYMPANI,

Arises by a small beginning from the extremity of the spinous process of the sphenoid bone, behind the entry of the artery of the dura mater; then running backwards, and a little upwards, along with the nerve called chorda tympani, in a sissure of the os temporis near the sossa that lodges the condyle of the lower jaw.

Inserted into the long process of the malleus, within the tympanum, where it rests upon the edge of the fis-

fure between the pars squamosa and petrosa.

Use. To draw the malleus obliquely forwards towards its origin, confequently the membrana tympani, by which that membrane is made less concave, or is relaxed.

Externus mallei, Albinus. Anterior mallei, Winflow. Obliquus auris, Douglas.

### 2. TENSOR TYMPANI,

Arises, by a very small beginning, from the cartilaginous extremity of the Eustachian tube, just where it begins to be covered by the pars petrosa, and spinous process of the sphenoid bone, near the entry of the artery of the dura mater; from thence running backwards near the offeous part of the Eustachian tube, forms a very distinct sleshy belly, below a thin offeous plate, between the pars squamosa and labyrinth; and sends off a slender tendon, which makes a turn into the tympanum along with the nerve called chorda tympani.

Inferted into the posterior part of the handle of the malleus, a little lower than the root of its long pro-

cefs.

Use. To pull the malleus and membrana tympani inwards towards the pars petrofa, by which the membrane is made more concave and tenfe.

Internus mallei, Winflow. Internus auris, Douglas.

#### 3. STAPEDIUS,

Arises, by a small fleshy belly, from a little cavern in the pars petrosa, near the cells of the mastoid process, before the inferior part of the passage for the portio dura of the auditory nerve; its tendon passes straight thro' a small round hole in the same cavern, enters the anterior part of the tympanum, and is

Inserted into the posterior part of the head of the

stapes.

Use. To draw the stapes obliquely upwards towards the cavern, by which the posterior part of its base is moved inwards, and the anterior part outwards.

Musculus stapedis, Winslow.

Stapidaus, Douglas.

#### C H A P. III.

Of the Muscles of the Eye-Lids.

THE palpebræ, or eye-lids, have one muscle common to both, and the upper eye-lid one proper to itself.

#### 1. ORBICULARIS PALPEBRARUM.

Arifes, by a number of fleshy fibres, from the outer edge of the orbitar process of the superior maxillary bone, and from a tendon near the inner angle of the eye: these run a little downwards, then outwards, over the upper part of the cheek, below the orbit, covering the under eye-lid, and surround the external angle, being loosely connected only to the skin and fat; run over the superciliary ridge of the os frontis, towards the inner canthus, where they intermix with those of the occipito-frontalis and corrugator supercilii; then covering the upper eye-lid, descend to the inner angle opposite to the inferior origin of this muscle, firmly adhering to the internal angular process of the os frontis, and to the short round tendon which serves to fix the palpebræ and muscular sibres arising from it.

Inferted, by the short round tendon, into the nasal process of the superior maxillary bone, covering the anterior and upper part of the lachrymal sac; which tendon can be easily felt at the inner canthus of the eye.

Use. To shut the eye, by drawing both lids close together, the sibres contracting from the outer angle towards the inner, press the eye-ball, squeeze the lach-Vol. I.

rymal gland, and convey the tears towards the puncta lachrymalia.

The ciliaris of some authors is only a part of this muscle covering the cartilages of the eye-lids, called ci-

lia or tarfi.

There is often a small fleshy slip which runs down from the outer and inferior part of this muscle above the zygomaticus minor, and joins with the levator labii superioris alæque nasi.

#### 2. LEVATOR PALPEBRÆ SUPERIORIS,

Arises from the upper part of the foramen opticum of the sphenoid bone, through which the optic nerve passes, above the levator oculi, near the trochlearis muscle.

Inserted, by a broad thin tendon, into the cartilage

that supports the upper eye-lid, named tarsus.

Use. To open the eye, by drawing the eye-lid upwards; which it does completely, by being fixed to the tarfus, pulling it below the eye-brow, and within the orbit.

Aperiens palpebram rectus, Douglas.

### C H A P. IV.

Muscles of the EYE-BALL.

THE muscles which move the globe of the eye are fix, viz.

Four fraight and two oblique.

The four straight muscles very much resemble each other; all

Arifing, by a narrow beginning, a little tendinous and fleshy,

fleshy, from the bottom of the orbit around the foramen opticum of the sphenoid bone, where the optic nerve enters, so that they may be taken out adhering to this nerve; and all having strong sleshy bellies.

Inserted at the fore-part of the globe of the eye into the anterior part of the tunica sclerotica, and under the tunica adnata, at opposite sides, which indicates both their names and Use; so that they scarcely require any further description, but to name them singly.

#### I. LEVATOR OCULI,

Arises from the upper part of the foramen opticum of the sphenoid bone, below the levator palpebræ superioris; and runs forwards to be

Inserted into the superior and fore-part of the tunica sclerotica, by a broad thin tendon.

Use. To raise up the globe of the eye.

Atollens, Albinus. Elevator, Douglas.

#### 2. DEPRESSOR OCULI,

Arises from the inferior part of the foramen opticum.

Inserted opposite to the former.

Use. To pull the globe of the eye down.

Deprimens, Albinus.

### 3. ADDUCTOR OCULI,

Arises, as the former, between the obliquus superior and depressor, being, from its situation, the shortest.

Inserted opposite to the inner angle.

Use. To turn the eye towards the nose.

### 4. ABDUCTOR OCULI,

Arises from the bony partition between the foramen opticum and lacerum, being the longest from its situation; and is

Inserted into the globe opposite to the outer canthus. Use. To move the globe outwards.

S 2

The

The oblique muscles are two:

### , 1. OBLIQUUS SUPERIOR, SEU TROCHLEARIS,

Arises, like the straight muscles, from the edge of the foramen opticum at the bottom of the orbit, between the levator and adductor oculi; from thence runs straight along the pars plana of the ethmoid bone to the upper part of the orbit, where a cartilaginous trochlea is fixed to the inside of the internal angular process of the os frontis, through which its tendon passes, and runs a little downwards and outwards, inclosed in a loose membranous sheath.

Inserted, by a broad thin tendon, into the tunica sclerotica, about half-way between the insertion of the at-

tollens oculi and optic nerve.

Use. To roll the globe of the eye, and to turn the pupil downwards and outwards, so that the upper side of the globe is turned inwards, and the inferior part to the outside of the orbit, and the whole globe drawn forwards towards the inner canthus.

Obliquus major, Winflow.

### 2. OBLIQUUS INFERIOR,

Arises, by a narrow beginning, from the outer edge of the orbitar process of the superior maxillary bone, near its juncture with the os unguis; and running obliquely outwards, is

Inferted into the sclerotica, in the space between the abductor and optic nerve, by a broad thin tendon.

Use. To draw the globe of the eye forwards, inwards, and downwards; and, contrary to the superior, to turn the pupil upwards, towards the inner extremity of the eye-brow; at the same time, the external part of the globe is turned towards the inferior side, and the internal rolls towards the upper part.

Obliquus minor, Winflow.

### C H A P. V.

Of the Muscle of the Nose.

THERE is only one muscle on each side that can be called proper to the nose, though it is affected by several muscles of the face.

#### COMPRESSOR NARIS,

Arises, by a narrow beginning, from the root of the ala nasi externally, where part of the levator labii superioris alæque nasi is connected to it; it spreads into a number of thin disgregated fibres, which run up along the cartilage in an oblique manner, towards the dorsum of the nose, where it joins with its fellow, and is

Inferted flightly into the anterior extremity of the os nasi and nasal process of the superior maxillary bone, where it meets with some of the sibres descending from the assinite frontalis muscle

the occipito-frontalis muscle.

Use. To compress the ala towards the septum nasi, particularly when we want to smell acutely; but, if the sibres of the frontal muscle which adhere to it act, the upper part of this thin muscle assists to pull the ala outwards. It also corrugates the skin of the nose, and assists in expressing certain passions.

Rinaus, vel nasalis, Douglas.

### C H A P. VI.

Muscles of the Mouth and Lips.

THE mouth has nine pair of muscles, which are inferted into the lips, and a common one formed by the termination of these, viz. three above, three below, three outwards, and the common muscle surrounds the mouth.

The three above are,

#### 1. LEVATOR ANGULI ORIS,

Arises, thin and fleshy, from the hollow of the superior maxillary bone, between the root of the socket of the first dens molaris and the foramen infraorbitarium.

Inserted into the angle of the mouth and under lip,

where it joins with its antagonist.

Use. To draw the corner of the mouth upwards, and make that part of the cheek opposite to the chin prominent, as in smiling.

Elevator labiorum communis, Douglas.

Caninus, Winflow.

### 2. LEVATOR LABII SUPERIORIS ALÆQUE NASI,

Arises by two distinct origins; the first, broad and sleshy, from the external part of the orbitar process of the superior maxillary bone which forms the lower part of the orbit, immediately above the foramen infra orbitarium; the second portion arises from the nasal process

cess of the superior maxillary bone, where it joins the os frontis at the inner canthus, descending along the edge of the groove for the lachrymal sac. The first and shortest portion is

Inferted into the upper lip and orbicularis labiorum; the fecond and longest, into the upper lip and outer

part of the ala nafi.

Use. To raise the upper lip towards the orbit, and a little outwards; the second portion serves to draw the skin of the nose upwards and outwards, by which the nostril is dilated.

Elevator labii superioris proprius, Douglas. Incisivus lateralis, First portion; Pyramidalis, Second portion; Winslow.

### 3. DEPRESSOR LABII SUPERIORIS ALÆQUE NASI,

Arises, thin and fleshy, from the os maxillare superius, immediately above the joining of the gums with the two dentes incisivi, and the dens caninus; from thence it runs up under part of the levator labii superioris alæque nasi.

Inserted into the upper lip and root of the ala nasi.

Use. To draw the upper lip and ala nasi downwards and backwards.

Depressor alæ nasi, Albinus. Incisivus medius, Winslow.

Depressor labii superioris proprius, Douglas.

The three below are,

### 1. DEPRESSOR ANGULI ORIS,

Arises, broad and fleshy, from the lower edge of the maxilla inferior, at the side of the chin, being firmly connected to that part of the platysma myoides, which runs over the maxilla to the angle of the mouth, to the depressor labii inferioris within, and to the skin and sat without, gradually turning narrower; and is

5 4

C

Inferted into the angle of the mouth, joining with the zygomaticus major and levator anguli oris.

Use. To pull down the corner of the mouth.

Triangularis, Winflow.

Depressor labiorum communis, Douglas.

### 2. DEPRESSOR LABII INFERIORIS,

Arises, broad and fleshy, intermixed with fat, from the inferior part of the lower jaw next the chin; runs obliquely upwards; and is

Inferted into the edge of the under lip, extends along

one half of the lip, and is loft in its red part.

Use. To pull the under lip and the skin of the side of the chin downwards, and a little outwards.

Quadratus, Winflow.

Depressor labii inferioris proprius, Douglas.

### 3. LEVATOR LABII INFERIORIS,

Arises, from the lower jaw, at the roots of the alveoli of two dentes incisivi and of the caninus; is

Inferted into the under-lip and skin of the chin.

Use. To pull the parts into which it is inserted up-

Levator menti, Albinus. Incisivus inferior, Winslow. Elevator labii inferioris proprius, Douglas.

The three outward are,

### 1. BUCCINATOR,

Arises, tendinous and fleshy, from the lower jaw, as far back as the last dens molaris and fore-part of the root of the coronoid process; sleshy from the upper jaw, between the last dens molaris and pterygoid process of the sphenoid bone; from the extremity of which it arises tendinous, being continued between both jaws to the constrictor pharyngis superior, with which it joins; from thence proceeding with straight fibres, and ad-

adhering close to the membrane that lines the mouth, it is

Inferted into the angle of the mouth within the orbi-

cularis oris.

Use. To draw the angle of the mouth backwards and outwards, and to contract its cavity, by pressing the cheek inwards, by which the food is small between the teeth.

Retractor anguli oris, Albinus.

### 2. ZYGOMATICUS MAJOR,

Arises, fleshy, from the os malæ, near the zygoma-tic suture.

Inserted into the angle of the mouth, appearing to be lost in the depressor anguli oris and orbicularis oris.

Use. To draw the corner of the mouth and underlip towards the origin of the muscle, and make the cheek prominent, as in laughing.

Zygomaticus, Douglas.

### 3. ZYGOMATICUS MINOR,

Arises from the upper prominent part of the os malæ, above the origin of the former muscle; and, descending obliquely downwards and forwards, is

Inserted into the upper lip, near the corner of the

mouth, along with the levator anguli oris.

Use. To draw the corner of the mouth obliquely outwards, and upwards, towards the external canthus of the eye.

The common muscle is the

#### ORBICULARIS ORIS.

This muscle is, in a great measure, formed by the muscles that move the lips; the fibres of the superior descending, those of the inferior ascending, and decussating each other about the corner of the mouth, run along the lip to join those of the opposite side, so that the

the fleshy fibres appear to surround the mouth like a

fphincter.

Use. To shut the mouth, by contracting and drawing both lips together, and to counteract all the muscles that assist in forming it.

Sphincter labiorum, Douglas. Semi-orbicularis, Winflow. Constrictor oris, Cowper.

There is another small muscle described by Albinus, which he calls Nasalis labii superioris; but it seems to be only some sibres of the former connected to the septum nasi.

#### C H A P. VII.

Muscles of the Lower Jaw.

THE lower jaw has four pair of muscles for its elevation or lateral motions, viz. two, which are feen on the side of the face, and two concealed by the angle of the jaw.

### I. TEMPORALIS,

Arises, fleshy, from a semicircular ridge of the lower and lateral part of the parietal bone, from all the pars squamosa of the temporal bone, from the external angular process of the os frontis, from the temporal process of the sphenoid bone, and from an aponeurosis which covers it: from these different origins the sibres descend like radii towards the jugum, under which they pass; and are

Inserted, by a strong tendon, into the upper part of the coronoid process of the lower jaw; in the duplica-

ture

1

is

8

y

of

ture of which tendon this process is inclosed as in a sheath, being continued down all its fore-part to near the last dens molaris.

Use. To pull the lower jaw upwards, and press it against the upper, at the same time drawing it a little backwards.

N. B. This muscle is covered by a tendinous membrane, called its aponeurosis, which arises from the bones that give origin to the upper and semicircular part of the muscle; and, descending over it, is inserted into all the jugum, and the adjoining part of the os frontis.

The use of this membrane is to give room for the origin of a greater number of sleshy sibres, to fortify the muscle in its action, and to serve as a desence to it.

Crotaphite muscle, Winflow.

#### 2. MASSETER.

Arises, by strong, tendinous, and sleshy sibres, which run in different directions, from the superior maxillary bone, where it joins the os malæ, and from the inferior and interior part of the zygoma, its whole length, as far back as the tubercle before the socket for the condyle of the lower jaw; the external sibres slanting backwards, and the internal forwards.

Inserted into the angle of the lower jaw, and from that upwards to near the top of its coronoid process.

Use. To pull the lower to the upper jaw, and, by means of its oblique decussation, a little forwards and backwards.

### 3. PTERYGOIDEUS INTERNUS,

Arises, tendinous and sleshy, from the inner and upper part of the internal plate of the pterygoid process, filling all the space between the two plates; and from the pterygoid process of the os palati between these plates.

Inserted into the angle of the lower jaw internally.

Use. To draw the jaw upwards, and obliquely towards the opposite side.

Pterygoideus major, Winflow.

### 4. PTERYGOIDEUS EXTERNUS,

Arises from the outer fide of the external plate of the pterygoid process of the sphenoid bone, from part of the tuberofity of the os maxillare adjoining to it, and from the root of the temporal process of the sphenoid bone.

Inferted into a cavity in the neck of the condyloid process of the lower jaw; some of its fibres are inserted into the ligament that connects the moveable cartilage

and that process to each other.

Use. To pull the lower jaw forwards, and to the opposite side; and to pull the ligament from the joint, that it may not be pinched during these motions: when both external pterygoid muscles act, the foreteeth of the under-jaw are pushed forwards beyond those of the upper jaw.

Pterygoideus minor, Winflow.

### C H A P. VIII.

The Muscles which appear about the anterior part of the NECK.

ON the fide of the neck are two muscles or layers.

1. MUSCULUS CUTANEUS, PLATYSMA MYOIDES.

Arises, by a number of slender disgregated sleshy fibres, II.

0-

he

of

nd

oid

oid

ed

ge

p-

nt,

S:

rend

of

hy

S,

fibres, from the cellular fubstance that covers the upper parts of the deltoid and pectoral muscles; in their ascent they all unite to form a thin muscle, which runs obliquely upwards along the side of the neck, adhering to the skin.

Inserted into the lower jaw, between its angle and the origin of the depressor anguli oris, to which it is simply connected, and but slightly to the skin that covers the inserior part of the masseter muscle and parotid glands.

Use. To affift the depressor anguli oris in drawing the skin of the cheek downwards; and when the mouth is shut, it draws all that part of the skin, to which it is connected, below the lower jaw, upwards.

Platysma myoides, Galen.

Musculus cutaneus, Winflow.

Quadratus genæ, vel Latissimus colli, Douglas.

Latissimus colli, Albinus.

#### 2. STERNO-CLEIDO-MASTOIDEUS,

Arises by two distinct origins; the anterior, tendinous and a little sleshy, from the top of the sternum near its junction with the clavicle; the posterior, sleshy, from the upper and anterior part of the clavicle; both unite a little above the anterior articulation of the clavicle, to form one muscle, which runs obliquely upwards and outwards, to be

Inserted, by a thick strong tendon, into the mastoid process, which it surrounds; and, gradually turning thinner, is inserted as far back as the lambdoid suture.

Use. To turn the head to one side, and bend it forwards.

Sterno-mastoideus and Cleido-mastoideus, Albinus. Mastoideus, Douglas.

C

cl

h

to

C

### C H A P. IX.

Muscles situated between the Lower JAW and Os Hyoldes.

THERE are four layers before, and two muscles at the side.

The four layers are,

#### 1. DIGASTRICUS,

Arises, by a fleshy belly, intermixed with tendinous sibres, from the fossa at the root of the maltoid process of the temporal bone, and soon becomes tendinous; runs downwards and forwards: the tendon passes generally through the stylo-hyoideus muscle; then it is sixed by a ligament to the os hyoides; and, having received from that bone an addition of tendinous and muscular sibres, runs obliquely forwards, turns sleshy again, and is

Inserted, by this anterior belly, into a rough finushity at the inserior and anterior edge of that part of the lower jaw called the chin.

Use. To open the mouth, by pulling the lower jaw downwards and backwards; and, when the jaws are shut, to raise the larynx, and consequently the pharynx, upwards, as in deglutition.

Biventer maxillæ inferioris, Albinus.

### 2. MYLO-HYOIDEUS,

Arises, fleshy, from all the inside of the lower jaw, be-

between the last dens molaris and the middle of the chin, where it joins with its fellow.

Inserted into the lower edge of the basis of the os

hyoides, and joins with its fellow.

Use. To pull the os hyoides forwards, upwards, and to a fide.

### 3. GENIO-HYOIDEUS,

Arises, tendinous, from a rough protuberance in the middle of the lower jaw internally, or inside of the chin.

Inserted into the basis of the os hyoides.
Use, To draw this bone forwards to the chin.

#### 4. GENIO-HYO-GLOSSUS,

Arises, tendinous, from a rough protuberance in the infide of the middle of the lower-jaw; its fibres run, like a fan, forwards, upwards, and backwards; and are

Inferted into the tip, middle, and root of the tongue,

and base of the os hyoides, near its cornu.

Use. According to the direction of its fibres, to draw the tip of the tongue backwards into the mouth, the middle downwards, and to render its dorfum concave; to draw its root and os hyoides forwards, and to thrust the tongue out of the mouth.

The two muscles at the side are,

### I. HYO-GLOSSUS,

Arises, broad and fleshy, from the base, cornu, and appendix of the os hyoides; the sibres run upwards and outwards, to be

Inserted into the fide of the tongue, near the stylo-

gloffus.

3

Use. To pull the tongue inwards and downwards.

Basio-cerato-chondro-glossus, Albinus.

Cerato-glossus, Douglas.

#### 2. LINGUALIS,

Arises from the root of the tongue laterally; runs forwards between the hysso-glossus and genio-glossus, to be

Inserted into the tip of the tongue, along with part

of the stylo-glossus.

Use. To contract the substance of the tongue, and bring it backwards.

### C H A P. X.

Muscles situated between the Os Hyoides and Trunk.

THESE may be divided into two layers.

The first layer confists of two muscles.

### I. STERNO-HYOIDEUS,

Arises, thin and fleshy, from the cartilaginous extremity of the first rib, the upper and inner part of the sternum, and from the clavicle where it joins with the sternum.

Inserted into the base of the os hyoides. Use. To pull the os hyoides downwards.

#### 2. OMO-HYOIDEUS,

Arises, broad, thin, and fleshy, from the superior costs of the scapula, near the similunar nitch, and from the ligament that runs across it; thence ascending obliquely, it becomes tendinous below the sterno-cleidomastoid muscle; and, growing sleshy again, is

IL

ns

18,

art

nd

K.

x-

ne

n n

)-

7-

Inserted into the base of the os hyoides, between its cornu and the insertion of the sterno-hyoideus.

Use. To pull the os hyoides obliquely downwards.

Coraco-hyoideus, Albinus and Douglas.

The fecond layer confifts of three muscles.

#### 1. STERNO-THYROIDEUS,

Arises, fleshy, from the whole edge of the uppermost bone of the sternum internally, opposite to the cartilage of the first rib, from which it receives a small part of its origin.

Inserted into the surface of the rough line at the external part of the inferior edge of the thyroid cartilage.

Use. To draw the larynx downwards.

#### 2. THYREO-HYOIDEUS,

Inserted into part of the basis, and almost all the cornu of the os hyoides.

Arises from the rough line, opposite to the former. Use. To pull the os hyoides downwards, or the thyroid cartilage upwards.

Thyro-hyoideus, vel Hyo-thyroideus, Winflow.

## 3. CRICO-THYROIDEUS,

Arises from the fide and fore-part of the cricoid cartilage, running obliquely upwards.

Inserted by two portions; the first, into the lower part of the thyroid cartilage; the second, into its inserior cornu.

Use. To pull forwards and depress the thyroid, or to elevate and draw backwards the cricoid cartilage.

Vol. I.

T

CHAP.

#### C H A P. XI.

Muscles situated between the Lower JAW and Os Hyoldes laterally.

THEY are five in number. Three proceed from the styloid process of the temporal bone; from which they have half of their names; and two from the pterygoid process of the sphenoid bone.

The three from the styloid process are,

#### 1. STYLO-GLOSSUS,

Arises, tendinous and sleshy, from the styloid process, and from a ligament that connects that process to the angle of the lower jaw.

Inserted into the root of the tongue, runs along its

fide, and is infenfibly loft near its tip.

Use. To draw the tongue laterally and backwards.

#### 2. STYLO-HYOIDEUS.

Arises, by a round tendon, from the middle and inferior part of the styloid process.

Inserted into the os hyoides at the junction of the

base and cornu.

Use. To pull the os hyoides to one side, and a little

upwards.

N. B. Its fleshy belly is generally perforated by the tendon of the digastric muscle, on one or both sides. There is often another accompanying it, called stylo-hyoideus,

byoideus alter; and has the same origin, insertion, and use.

#### 3. STYLO-PHARYNGEUS,

Arises, fleshy, from the root of the styloid process.

Inserted into the side of the pharynx and back-part of the thyroid cartilage.

Use. To dilate and raise the pharynx and thyroid car-

tilage upwards.

m

m

to

ts

e-

ne

le

ne

S.

s,

The two from the pterygoid process are,

## 1. CIRCUMFLEXUS, OR TENSOR PALATI,

Arises from the spinous process of the sphenoid bone, behind the foramen ovale, which transmits the third branch of the sisth pair of nerves; from the Eustachian tube, not far from its osseous part: it then runs down along the pterygoideus internus, passes over the hook of the internal plate of the pterygoid process by a round tendon, which soon spreads into a broad membrane.

Inserted into the velum pendulum palati, and the se milunar edge of the os palati, and extends as far as the suture which joins the two bones. Generally some of its posterior sibres join with the constrictor pharyngis superior, and palato-pharyngeus.

Use. To stretch the velum, to draw it downwards, and to a side towards the hook. It has little effect upon the tube, being chiefly connected to its offeous part.

Circumflexus palati, Albinus.

Spheno-salpingo-staphilinus, seu Staphilinus externus, Winslow.

Musculus tubæ novus, Valsalva; vel Palato-salphingeus, Douglas.

## 2. LEVATOR PALATI,

Arises, tendinous and sleshy, from the extremity of the pars petrosa of the temporal bone, where it is perforated

forated by the Eustachian tube, and also from the membranous part of the same tube.

Inferted into the whole length of the velum pendulum palati, as far as the root of the uvula, and unites with its fellow.

Use. To draw the velum upwards and backwards, fo as to shut the passage from the fauces into the mouth and nose.

Levator palati mollis, Albinus.

Petro-salpingo-staphilinus, vel Salpingo-staphilinus internus, Winflow.

Salpingo-staphilinus, Valsalva. Pterigo-staphilinus externus, vulgo, Douglas.

Spheno-staphilinus, Couper.

Previous to the description of the muscles situated about the passage into the throat, it will be necessary to mention the principal parts to which they are connected.

Upon looking into any person's mouth, when wide opened, we see a soft curtain hanging from the palatebones, named velum pendulum palati. In the middle of which, we likewise observe a papilla projecting from the velom, named uvula, or pap of the throat. From each side of the uvula, at its root, two arches, or columns, are sent down; the anterior to the root of the tongue, the posterior to the pharynx. Between these arches, on each side, the cellular glands called amygdalæ, or almonds of the ears, are situated.

The common opening behind the anterior arch may be named fauces, or top of the threat, from which there are fix passages, viz. two upwards, being one to each nostril: two at the sides, or one to each ear, called the Eustachian tubes: two downwards; the anterior is the passage through the glottis and larynx, into the trachea, which terminates in the lungs; the posterior is the largest, named pharynx, or top of the assophagus,

which leads to the stomach.

#### C H A P. XII.

Muscles fituated about the entry to the FAUCES.

THE RE are two on each fide, and a fingle one in the middle.

The two on each fide are,

#### 1. CONSTRICTOR ISTHMI FAUCIUM,

Arises, by a slender beginning, from the side of the tongue, near its root; from thence running upwards, within the anterior arch, before the amygdala, it is

Inserted into the middle of the velum pendulum palati, at the root of the uvula anteriorly, being connected with its fellow, and with the beginning of the palato-pharyngeus.

Use. Draws the velum towards the root of the tongue, which it raises at the same time, and, with its fellow, contracts the passage between the two arches, by which it shuts the opening into the sauces.

Gloffo-staphilinus, Winflow and Douglas.

#### 2. PALATO-PHARYNGEUS.

Arises, by a broad beginning, from the middle of the velum pendulum palati, at the root of the uvula posteriorly, and from the tendinous expansion of the circumstexus palati. The fibres are collected within the posterior arch behind the amygdala, and run backwards to the top and lateral part of the pharynx, where the fibres are scattered, and mix with those of the stylopharyngeus.

Inferted

Inferted into the edge of the upper and back part of the thyroid cartilage; some of its sibres being lost between the membrane of the pharynx and the two inferior constrictors.

Use. Draws the uvula and velum downwards and backwards; and at the same time pulls the thyroid cartilage and pharynx upwards, and shortens it; with the constrictor superior and tongue, it assists in shutting the passage into the nostrils; and, in swallowing, it thrusts the food from the fauces into the pharynx.

Thyro-staphilinus, Douglas.

Thyro-pharyngo-staphilinus, Winflow.

SALPINGO-PHARYNGEUS of Albinus is composed of a few fibres of this muscle, which

Arise from the anterior and lower part of the cartilaginous extremity of the Eustachian tube; and are

Inferted into the inner part of the last-mentioned

muscle.

Ufe. To affist the former, and to dilate the mouth of the tube.

The one in the middle is the

## AZYGOS UVULÆ.

Arifes, fleshy, from the extremity of the suture which joins the palate-bones, runs down the whole length of the velum and uvula, resembling a small earth-worm, and adhering to the tendons of the circumslexi.

Inserted into the tip of the uvula.

Ufe. Raifes the uvula upwards and forwards, and shortens it.

Palato-stapbilinus, Douglas.

Staphilinus, or Epistaphilinus, Winslow.

#### C H A P. XIII.

Muscles fituated on the posterior part of the PHARYNK.

OF these there are three pair.

## 1. CONSTRICTOR PHARYNGIS INFERIOR,

Arises from the side of the thyroid cartilage, near the attachment of the sterno-hyoideus and thyreo-hyoideus muscles; and from the cricoid cartilage, near the crico-thyroideus. This muscle is the largest of the three, and is

Inferted into the white line, where it joins with its fellow; the superior fibres running obliquely upwards, covering nearly one half of the middle constrictor, and terminating in a point; the inferior fibres run more transversely, and cover the beginning of the cesophagus.

Use. To compress that part of the pharynx which it covers, and to raise it with the larynx a little upwards.

Thyro-pharyngeus, Crico-pharyngeus, Douglas.

#### 2. CONSTRICTOR PHARYNGIS MEDIUS.

Arises from the appendix of the os hyoides, from the cornu of that bone, and from the ligament which connects it to the thyroid cartilage; the fibres of the superior part running obliquely upwards, and, covering a considerable part of the superior constrictor, terminate in a point.

Inserted into the middle of the cuneiform process of T4

the os occipitis, before the foramen magnum, and joined to its fellow at a white line in the middle back part of the pharynx. The fibres at the middle part run more transversely than those above or below.

Use. To compress that part of the pharynx which it covers, and to draw it and the os hyoides upwards.

Hyo-pharyngeus, Syndesmo-pharyngeus, Douglas.

#### 3. CONSTRICTOR PHARYNGIS SUPERIOR,

Arises, above, from the cuneiform process of the os occipitis, before the foramen magnum, near the holes where the ninth pair of nerves passes out; lower down, from the pterygoid process of the sphenoid bone; from the upper and under jaw, near the roots of the last dentes molares; and between the jaws, it is continued with the buccinator muscle, and with some fibres from the root of the tongue and from the palate.

Inserted into a white line in the middle of the pharynx, where it joins with its fellow, and is covered by

the constrictor medius.

Use. To compress the upper part of the pharynx, and draw it forwards and upwards.

Gephalo-pharyngeus, Pterygo-pharyngeus, Mylo-pharyngeus, Glosso-pharyngeus, Douglas.

## C H A P. XIV.

Muscles situated about the GLOTTIS.

THEY confift generally of four pair of small muscles, and a single one.

1. CRI-

## r. CRICO-ARYTÆNOIDEUS POSTICUS,

Arises, fleshy, from the back-part of the cricoid cartilage, and is

Inserted into the posterior part of the base of the ary-

tenoid cartilage.

Use. To open the rima glottidis a little, and, by pulling back the arytenoid cartilage, to stretch the ligament so as to make it tense.

#### 2. CRICO ARTYÆNOIDEUS LATERALIS.

Arises, fleshy, from the cricoid cartilage, laterally, where it is covered by part of the thyroid, and is

Inserted into the fide of the base of the arytenoid car-

tilage near the former.

Use. To open the rima glottidis, by pulling the ligaments from each other.

#### 3. THYREO-ARYTÆNOIDEUS,

Arises from the under and back part of the middle of the thyroid cartilage; and, running backwards and a little upwards, alongst the side of the glottis, is —

Inferted into the arytenoid cartilage, higher up and farther forwards than the crico-arytænoideus latera-

lis.

Use. To pull the arytenoid cartilage forwards nearer to the middle of the thyroid, and consequently to shorten and relax the ligament of the larynx or glottis vera.

## 4. ARYTÆNOIDEUS OBLIQUUS.

Arises from the base of one arytenoid cartilage; and, crossing its fellow, is

Inferted near the tip of the other arytenoid carti-

lage.

Use. When both act, they pull the arytenoid cartilages towards each other.

N. B. Very often one of these is wanting.

Arytenoideus minor, Douglas.

The

1

The fingle muscle is the

## ARYTÆNOIDEUS TRANSVERSUS.

Arises from the fide of one arytenoid cartilage, from near its articulation with the cricoid to near its tip. The fibres run straight across; and are

Inserted, in the same manner, into the other aryte-

noid cartilage.

Use. To shut the rima glottidis, by bringing these two cartilages, with the ligaments, nearer on another.

Arytanoideus major, Douglas.

Besides these, there are a few disgregated muscular fibres on each side; which, from their general direction, are named,

#### 4. THYREO EPIGLOTTIDEUS,

Arises, by a few pale disgregated fibres from the thyroid cartilage; and is

Inferted into the epiglottis laterally.

Use. To draw the epiglottis obliquely downwards, or, when both act, directly downwards; and, at the same time, it expands that fost cartilage.

## 2. ARYTÆNO-EPIGLOTTIDEUS,

Arises, by a number of small sibres, from the lateral and upper part of the arytænoid cartilage; and, running along the outer side of the external rima, is

Inserted into the epiglottis along with the former.

Use. To pull that side of the epiglottis towards the external rima; or, when both act, to pull it close upon the glottis. It is counteracted by the elasticity of the epiglottis.

## C H A P. XV.

Muscles situated on the anterior part of the Abdomen.

THEY confift of three broad layers on each fide of the belly; always a long one, and generally also a short one, on each fide of the linea alba.

The three layers are,

II.

e

T

## 1. OBLIQUUS DESCENDENS EXTERNUS,

Arifes, by eight heads, from the lower edges of an equal number of inferior ribs, at a little distance from their cartilages: it always intermixes, in a ferrated manner, with portions of the serratus major anticus; and generally coheres to the pectoralis major, intercostals, and latissimus dorsi; which last covers the edge of a portion of it extended from the last rib to the spine of the os ilium. From these origins the sibres run down obliquely forwards, and terminate in a thin broad tendon, whose sibres are continued in the same direction.

Inferted into the whole length of the linea alba \*; becomes thicker towards the lower part of the abdomen, and is perforated in the middle by the umbilicus †. On the outside of the rectus muscle, the tendon

† The umbilious was originally the passage for the vessels that

<sup>\*</sup> The linea alba is formed by the tendinous fibres of the two oblique and transverse muscles, interlaced with those of the opposite side, the whole way from the cartilago ensistormis to the os pubis; so that some think they should be called three digastric muscles, with a broad middle tendon and two sleshy bellies.

b

tl

li

C

fe

don of the external oblique appears whiter than elfewhere, by its being there connected with the tendons of the internal oblique and transverse muscles; so that this part has been called 'Linea semilunaris, from its curved shape. The under part of the tendon divides into two columns, which leaves an oval space between them, named the ring sof the external oblique muscle, for the passage of the spermatic cord in the male, or round ligament of the womb: The anterior superior column passes over the cartilage between the ossa pubis, and is fixed to the opposite os pubis; the other is fixed to the os pubis of the same side. It is also inserted, tendinous and sleshy, into the middle of the spine of the ilium.

From that part, which is named its anterior fuperior spinous process, it is stretched tendinous to the os pubis, and is named Poupart's or Fallopius's ligament \*. From this ligament it sends a tendinous layer, which is

lost in the membranous fascia of the thigh.

Use. Supports and compresses the peritoneum and abdomen; assists the evacuations of faces and urine, and likewise in the exclusion of the foetus; thrusts the diaphragm upwards, and draws down the ribs in expi-

ra

connected the feetus to the fecundines; and is really a hole through the teguments and tendons, filled up only by a cellular fubstance,

and covered within by the peritoneum.

If the ring of the external oblique muscle is made somewhat circular, by a thin tendinous or rough cellular substance, which helps to fill it up; and though a few muscular sibres of the internal are separated, yet the stricture in hernize only happens in the tendon of the external.

\* Poupart, or Fallopius's ligament, is the inferior part of the tendon of the external oblique, extending from the anterior superior spinous process of the ilium to the os pubis, where it is thickest, in order to strengthen the inferior part of the abdomen: here it is not inserted into any bone, but passes over the blood-vessels of the inferior extremity; and in women, from the gerater size of the pelvis, is longer and looser, by which they are more subject to crural herniæ; but, by the size of the spermatic cord, men are more liable to the inguinal.

I.

e.

18

at

ts

es

10

r

1-

IS

1-

e

r

S

d

e

.

h

ration; bends the body obliquely when the ribs are fixed, and raises the pelvis obliquely.

Obliquus externus abdominis, Albinus. Obliquus descendens, Douglas.

## 2. OBLIQUUS ASCENDENS INTERNUS,

Arises from the spine of the ilium, the whole length between the posterior and superior anterior spinous process; from the os sacrum and the three undermost lumbar vertebræ, by a tendon common to it and to the serratus posticus inferior muscle; from Poupart's ligament, at the middle of which it sends off the beginning of the cremaster muscle; and the spermatic cord in the male, or round ligament of the womb, passes under its thin edge, except a few detached sibres.

Inferted into the cartilago ensiformis, into the cartilages of the seventh and those of all the salse ribs; but, at the upper part, it is extremely thin, resembling a cellular membrane, and only becomes sleshy at the cartilage of the tenth rib. Here its tendon divides into two layers \*; the anterior layer, with a great portion of the inferior part of the posterior layer, joins the tendon of the external oblique, and runs over the rectus to be inferted into the whole length of the linea alba. The posterior layer joins the tendon of the transversalis muscle as low as half-way between the umbilicus and os pubis; but, below this place, only a few sibres of the posterior layer are seen, and the rest of it passes before the rectus muscle, and is inserted into the linea alba; so that the whole tendon of the external oblique

<sup>\*</sup> To obtain a proper view of the two layers of the tendon of the internal oblique muscle, both the oblique muscles should be raised as far forwards as their joining near the linea semilunaris; then the tendon before the rectus must be cut parallel to the linea alba, and turned outwards as far as the outer edge of the rectus; by which the whole of the rectus is brought into view, and the tendons are preserved: But Douglas directs to cut the posterior layer of the internal oblique, where it joins with the transversalis; by this method the rectus is laid bare, but the structure of the tendinous sheath, which incloses it, is destroyed.

(

t

muscle, with the anterior layer of the internal oblique, passes before the rectus muscle; and the whole posterior layer of the internal oblique, together with the whole tendon of the transversalis muscle, excepting at the inferior part, pass behind the rectus, and are inferted into the linea alba. At its undermost part it is inserted into the fore-part of the os pubis.

Use. To affift the former; but it bends the trunk in

the reverse direction.

Obliquus internus abdominis. Albinus and Winflow. Obliquus afcendens, Douglas.

## 3. TRANSVERSALIS,

Arifes tendinous, but soon becoming sleshy from the inner or back part of the cartilages of the seven lower ribs, where some of its sibres are continued with those of the diaphragm and the intercostal muscles; by a broad thin tendon, connected to the transverse processes of the last vertebra of the back and the sour superior vertebræ of the loins; sleshy, from the whole spine of the os ilium internally, and from the tendon of the external oblique muscle, where it intermixes with some sibres of the internal oblique.

Inferted into the cartilago enfiformis, and into the whole length of the linea alba, excepting its lowermost

part.

Use. To support and compress the abdominal bowels; and it is so particularly well adapted for the latter purpose, that it might be called the proper constrictor of the abdomen.

Transversus abdominis, Albinus.

The long muscle in the middle is named,

## RECTUS ABDOMINIS,

Arises, by two heads, from the ligament of the cartilage which joins the two offa pubis to each other; runs upwards the whole length of, and parallel to, the linea

II.

ie,

te.

he

at

n-

18

in

ne

er

fe

a

·f-

e-

ne

le

ne st

r-

e

linea alba, growing broader and thinner as it ascends.

Inferted into the cartilages of the three inferior true ribs, and often intermixes with some fibres of the pectoral muscle.

It is generally divided by three tendinous interfections; the first is at the umbilicus, the second where it runs over the cartilage of the seventh rib, the third in the middle between these; and there is commonly a half intersection below the umbilicus: These seldom penetrate through the whole thickness of the muscle; they adhere firmly to the anterior part of the sheath, but very slightly to the posterior layer.

Use. To compress the fore-part, but more particularly the lower part of the belly; to bend the trunk forwards, or to raise the pelvis. By its tendinous intersections, it is enabled to contract at any of the intermediate spaces; and, by its connection with the tendons of the other muscles, it is prevented from changing place, and from rising into a prominent form when in action.

## The short muscle in the middle is named

## PYRAMIDALIS,

Arises along with the rectus; and, running upwards within the same sheath, is

Inferted, by an acute terminination, near half-way between the os pubis and umbilicus, into the linea alba and inner edge of the rectus muscle.

As it is frequently wanting in both fides, without any inconveniency, its

Use seems to be, to affist the inferior part of the rec-

# C H A P. XVI.

Muscles about the Male Organs of GENERATION.

THE testicles are said to have a thin muscle common to both, and have one proper to each.

The fupposed common muscle is called the

#### DARTOS.

This appears to be no more than a condensation of the cellular membrane lining the scrotum; yet the skin here is capable of being corrugated and relaxed in a greater degree than in other places.

The muscle proper to each testicle is the

#### CREMASTER.

Arifes from the internal oblique, where a few fibres of that muscle intermix with the transversalis, near the junction of the os ilium and pubis, over which part it passes, after having pierced the ring of the external oblique; and then it descends upon the spermatic cord.

Inferted into the tunica vaginalis of the testicle, upon which it spreads, and is insensibly lost.

Use. To suspend and draw up the testicle, and to compress it in the act of coition.

The penis has three pair of muscles,

I.

in

W

h

**C-**

1-

0

#### I. ERECTOR PENIS.

Arises, tendinous and sleshy, from the tuberosity of the os ischium, and runs upwards, embracing the whole crus of the penis.

Inferted into the strong tendinous membrane that covers the corpora cavernosa penis, near as far up as the union of these bodies.

Use. To compress the crus penis, by which the blood is pushed from it into the fore-part of the corpora cavernosa; and the penis is by that means more completely distended. The erectores feem likewise to keep the penis in its proper direction.

Ischio-cavernosus, Winslow.

# 2. ACCELERATOR URINÆ, SEU EJACULATOR SEMINIS,

Arises, fleshy, from the sphincter ani and membranous part of the urethra; and tendinous from the crus, near as far forwards as the beginning of the corpus cavernosum penis; the inferior sibres run more transversely, and the superior descend in an oblique direction.

Inserted into a line in the middle of the bulb, where it joins with its fellow, by which the bulb is completely inclosed.

Use. To drive the urine or semen forwards; and, by grasping the bulb of the urethra, to push the blood towards its corpus cavernosum and the glans, by which they are distended.

Bulbo-cavernofus, Winflow.

## 3. TRANSVERSUS PERINEI,

Arises from the tough fatty membrane that covers the tuberosity of the os ischium; from thence it runs transversely inwards, and is

Vol. I. U In-

Inserted into the accelerator urinæ, and into that part

of the sphincter ani which covers the bulb.

Use. To dilate the bulb, and draw the perineum and verge of the anus a little outwards and backwards.

Transversalis urethra, Winslow. Transversus perinei, Albinus. Levator parvus, seu externus, Douglas.

There is often a fourth muscle, named

## TRANSVERSUS PERINEI ALTER,

Arises behind the former, runs more obliquely forwards, and is

Inserted into that part of the accelerator urinæ which covers the anterior part of the bulb of the urethra.

Use. To affish the former.

Inferior prostate, Winslow.

Transversus perinei alter, Albinus.

## C H A P. XVII.

Muscles of the Anus.

THE anus has a fingle muscle, and one pair.

The fingle muscle is

## SPHINCTER ANI,

Arises from the skin and fat that surround the verge of the anus on both sides, near as far out as the tuber of the os ischium; the sibres are gradually collected into

rt

n

into an oval form, and furround the extremity of the rectum.

Inserted, before, by a narrow point, into the perineum, acceleratores urinæ, and transversi perinei; behind, by an acute termination, into the extremity of the os coccygis.

Use. Shuts the passage through the anus into the rectum; pulls down the bulb of the urethra, by which it assists in ejecting the urine and semen.

Sphincler externus, Albinus and Douglas,

Sphineter cutaneus, Winflow.

N. B. The sphincter internus of Albinus and Douglas, is only that part of the circular sibres of the muscular coat of the rectum, which surrounds its extremity.

## LEVATOR ANI,

Arises from the os pubis within the pelvis, as far up as the upper edge of the foramen thyroideum, and joining of the os pubis with the os ischium; from the thin tendinous membrane that covers the obturator internus and coccygeus muscles; from the spinous process of the os ischium; and its sibres run down like rays from a circumference to a centre.

Inserted into the sphincter ani, acceleratores urinæ, and anterior part of the two last bones of the coccygis; surrounds the extremity of the rectum, neck of the bladder, prostate gland, and part of the vesiculæ seminales; so that its sibres behind and below the os coccygis joining it with its fellow, they, together, very much resemble the shape of a funnel.

Use. To draw the rectum upwards after the evacuation of the fæces, and to assist in shutting it; to sustain the contents of the pelvis, and to help in ejecting the semen, urine, and contents of the rectum; and, perhaps, by pressing upon the veins, to contribute greatly

to the erection of the penis.

## C H A P. XVIII.

Muscles of the Female Organs of Generation.

THE clitoris has one pair,

#### ERECTOR CLITORIDIS.

Arises from the crus of the os ischium internally, and in its ascent covers the crus of the clitoris as far up as the os pubis.

Inserted into the upper part of the crus and body of

the clitoris.

Use. Draws the clitoris downwards and backwards; and may serve to make the body of the clitoris more tense, by squeezing the blood into it from its crus.

First muscle of the clitoris, Douglas.

The vagina has one pair,

## SPHINCTER VAGINÆ,

Arises from the sphincter ani, and from the posterior side of the vagina, near the perineum; from thence it runs up the side of the vagina, near its external orisice, opposite to the nymphæ, and covers the corpus cavernosum vaginæ.

Inferted into the crus and body, or union of the

crura clitoridis.

Use. Contracts the mouth of the vagina, and compresses its corpus cavernosum.

Constrictor cunni, Albinus.

Second muscle of the clitoris, Douglas.

The perineum has one pair.

II.

nd

up

of

S;

re

or

it

ri-

us

he

n-

he

#### TRANSVERSUS PERINEI.

Arises, as in the male, from the fatty cellular membrane which covers the tuberosity of the os ischium.

Inferted into the upper part of the sphincter ani, and into a white hardish tough substance in the perineum, between the lower part of the pudendum and anus.

Use. To fustain and keep the perineum in its proper place.

The anus, as in the male, has a fingle muscle, and one pair.

#### SPHINCTER ANI,

Arises, as in the male, from the skin and fat surrounding the extremity of the rectum.

Inserted, above, into the white tough substance of the perineum; and below, into the point of the os coccygis.

Use. To shut the passage into the rectum; and, by pulling down the perineum, to assist in contracting the mouth of the vagina.

## LEVATOR ANI,

Arises, as in the male, within the pelvis, and defeends along the inferior part of the vagina and rectum.

Inserted into the perineum, sphincter ani, extremity of the vagina, and rectum.

Use. To raise the extremity of the rectum upwards, to contract the inferior part of the rectum, and to assist in contracting and supporting the vagina; and, perhaps, by pressing on the veins, to contribute to the distention of the cells of the clitoris and corpus caver-

nofum of the vagina.

## C H A P. XIX.

Muscles situated within the PELVIS-

OF these there are two pair.

## 1. OBTURATOR INTERNUS,

Arises from more than one half of the internal circumference of the foramen thyroideum, formed by the os pubis and ischium: its inside is covered by a portion of the levator ani; and appears to be divided into a number of sasciculi, which unite and form a roundish tendon, that passes out of the pelvis, between the posterior sacro-ischiatic ligament and tuberosity of the os ischium: where it passes over the capsular ligament of the thigh-bone, it is inclosed, as in a sheath, by the gemini muscles.

Inserted, by a round tendon, into the large pit at the

root of the trochanter major.

Use. To roll the os femoris obliquely outwards.

Marsupialis, seu Obturator internus, Douglas.

N. B. The infertion of this muscle should not be profecuted, until the muscles of the thigh, to which it belongs, are diffected. Vid. Chap. xxix.

#### 2. COCCYGEUS.

Arises, tendinous and fleshy, from the spinous process of the os ischium, and covers the inside of the posterior sacro-ischiatic ligament; from this narrow beginning, it gradually increases, to form a thin sleshy belly, interspersed with tendinous sibres.

In-

II.

ir-

he

a

fh

0-

09

of

he

ne

0-

e-

e

y

3-

Inferted into the extremity of the os facrum, and near the whole length of the os coccygis laterally.

Use. To support and move the os coccygis forwards, and to tie it more firmly to the facrum.

## C H A P. XX.

Muscles fituated within the Cavity of the Abdomen.

THESE confist of a single muscle, and sour pair.

#### DIAPHRAGMA,

This broad thin muscle, which makes a complete septum between the thorax and abdomen, is concave below and convex above; the middle of it on each side reaching as high within the thorax of the skeleton as the fourth rib, and is commonly divided into two portions.

1. The Superior or Greater Muscle of the DIAPHRAGM,

Arises, by distinct sleshy fibres, from the cartilago ensistentials, from the cartilages of the seventh, and of all the inferior ribs on both sides: The sibres from the cartilago ensistentials, and from the seventh and eighth ribs, run obliquely upwards and backwards; from the ninth and tenth, transversely inwards and upwards; and from the eleventh and twelsth, obliquely upwards. From these different origins the sibres run, like radii from the circumference to the centre of a circle; and are

Inserted into a cordiform tendon, of a confiderable breadth, which is situated in the middle of the diaphragm,

phragm; and in which, therefore, the fibres from opposite sides are interlaced. Towards the right side the tendon is perforated, by a triangular hole, for the passage of the vena cava inferior; and to the upper convex part of it the pericardium and mediastinum are connected.

# 2. The Inferior, Leffer Mofele, or Appendix of the DIAPHRAGM,

Arises from the second, third, and sourth lumbar vertebræ, by eight heads; of which two in the middle, commonly called its crura, are the longest, and begin tendinous. Between the crura, the aorta and thoracic duct pass; and, on the outside of these, the great sympathetic nerves and branches of the vena azygos perforate the shorter heads. The muscular sibres run obliquely upwards and forwards, and form in the middle two sleshy columns, which decustate and leave an oval space between them for the passage of the cesophagus and eighth pair of nerves.

Inserted, by strong sleshy fibres, into the posterior

part of the middle tendon.

Use. The diaphragm is the principal agent in respiration, particularly in inspiration: for when it is in action, the fibres, from their different attachments, endeavour to bring themselves into a plane towards the middle tendon, by which the cavity of the thorax is enlarged, particularly at the fides, where the lungs are chiefly fituated; and as the lungs must always be contiguous to the infide of the thorax and upper fide of the diaphragm, the air rushes into them, in order to fill up the increased space. This muscle is assisted by the two rows of intercoftals, which elevate the ribs, and the cavity of the thorax is more enlarged. In time of violent exercise, or whatever cause drives the blood with unufual celerity towards the lungs, the pectoral muscles, the ferrati antici majores, the ferrati postici fuperiores, and scaleni muscles, are brought into action.

tion. And in laborious inspiration, the muscles which arise from the upper part of the thorax, when the parts into which they are inserted are fixed, likewise assist. In expiration, the diaphragm is relaxed and pushed up by the pressure of the abdominal muscles upon the viscera of the abdomen; and at the same time that they press it upwards, they also, together with the sternocostales and serrati postici interiores, pull down the ribs, and are assisted in a powerful manner by the elasticity of the cartilages that join the ribs to the sternum; by which the cavity of the thorax is diminished, and the air suddenly pushed out of the lungs: and, in laborious expiration, the quadrati lumborum, sacrolumbales, and longishimi dorsi, concur in pulling down the ribs.

The four pair are,

I.

)-

le f-

)-

e

ır

,

n

C

-

e e

1

S

#### 1. QUADRATUS LUMBORUM.

Arises, pretty broad, tendinous and sleshy, from the

posterior part of the spine of the os ilium.

Inserted into the transverse processes of all the vertebræ of the loins, into the last rib near the spine, and by a small tendon into the side of the last vertebra of the back.

Use. To move the loins to one fide, pull down the last rib, and, when both act, to bend the loins forwards.

Quadratus, seu Lumbaris externus, Winslow.

## 2. PSOAS PARVUS,

Arises, fleshy, from the sides of the two upper vertebræ of the loins, and sends off a small long tendon, which ends thin and slat, and is

Inserted into the brim of the pelvis, at the junction of

the os ilium and pubis.

Use. To affift the ploas magnus in bending the loins for-

forwards; and, in certain positions, to assist in raising the pelvis.

N. B. This muscle is very often wanting.

#### 3. PSOAS MAGNUS.

Arises, fleshy, from the side of the body, and transverse process of the last vertebra of the back; and, in the same manner, from all those of the loins, by as many distinct slips.

Inserted, tendinous, into the trochanter minor of the os femoris; and fleshy into that bone, a little below the

fame trochanter.

Use. To bend the thigh forwards; or, when the inferior extremity is fixed, to affift in bending the body. Psoas, seu Lumbaris internus, Winslow.

#### 4. ILIACUS INTERNUS.

Arises, fleshy, from the transverse process of the last vertebra of the loins, from all the inner lip of the spine of the os ilium, from the edge of that bone between its anterior superior spinous process and the acetabulum, and from most of the hollow part of the ilium. It joins with the psoas magnus, where it begins to become tendinous; and is

Inferted along with it.

Use. To affift the ploas in bending the thigh, and to bring it directly forwards.

N. B. The infertion of the two last muscles should not be prosecuted till the muscles of the thigh are dissected.

g

IS

e

e

,

e

1

1

## C H A P. XXI.

Muscles situated on the Anterior Part of the THORAX.

THESE may be divided into two layers. The first layer confists of one muscle, named

## PECTORALIS MAJOR,

Arises from the cartilaginous extremities of the fifth and fixth ribs, where it always intermixes with the external oblique muscle of the abdomen; from almost the whole length of the sternum, and from near half of the anterior part of the clavicle: The sibres run towards the axilla in a folding manner.

Inserted, by two broad tendons, which cross each other at the upper and inner part of the os humeri, above the insertion of the deltoid muscle, and outer side of the groove for lodging the tendon of the long head of the biceps.

Use. To move the arm forwards, and obliquely upwards, towards the sternum.

Pectoralis, Albinus.

The fecond layer confifts of three muscles;

#### 1. SUBCLAVIUS,

Arises tendinous from the cartilage that joins the first rib to the sternum.

Inferted, after becoming fleshy, into the inferior part of the clavicle, which it occupies from within an inch or so of the sternum, as far outwards as to its connection, by ligament, with the coracoid process of the scapula.

Use.

Use. To pull the clavicle downwards and forwards.

#### 2. PECTORALIS MINOR,

Arises, tendinous and sleshy, from the upper edge of the third, fourth, and sifth ribs, near where they join with their cartilages.

Inserted, tendinous, into the coracoid process of the

fcapula; but foon grows fleshy and broad.

Use. To bring the scapula forwards and downwards, or to raise the ribs upwards.

Serratus anticus, Albinus.

Serratus minor anticus, Douglas.

#### 3. SERRATUS MAGNUS,

Arises from the nine superior ribs, by an equal number of sleshy digitations, resembling the teeth of a saw.

Inserted, fleshy, into the whole base of the scapula internally, between the insertion of the rhomboid and the origin of the subscapularis muscles, being folded about the two angles of the scapula.

Use. To move the scapula forwards; and, when the scapula is forcibly raised, to draw upwards the ribs.

Serratus major anticus, Douglas.

## C H A P. XXII.

Muscles situated between the Ribs, and within the Thorax.

BETWEEN the ribs, on each fide, there are eleven double rows of muscles, which are therefore named intercostals. These decussate each other like the strokes of the letter X.

#### I. INTERCOSTALES EXTERNI,

Arise from the inferior acute edge of each superior rib, and run obliquely forwards, the whole length from the spine to near the joining of the ribs with their cartilages; from which, to the sternum, there is only a thin membrane covering the internal intercostals.

Inferted into the upper obtuse edge of each inferior rib, as far back as the spine, into which the posterior

portion is fixed.

II.

y

e

#### INTERCOSTALES INTERNI,

Arise in the same manner as the external: but they begin at the sternum, and run obliquely backwards, as far as the angle of the rib; and from that to the spine they are wanting.

Inserted in the same manner as the external.

Use. By means of these muscles, the ribs are equally raifed upwards during inspiration. Their fibres being oblique, give them a greater power of bringing the ribs near each other, than could be performed by straight ones. But, by the obliquity of the fibres, they are almost brought contiguous: and as the fixed points of the ribs are before and behind, if the external had been continued forwards to the sternum, and the internal backwards to the spine, it would have hindered their motion, which is greatest in the middle, though the obliquity of the ribs renders it less perceptible; and, instead of raising the fibres fixed to the sternum and spine, would have depressed the ribs.

N. B. The portions of the external intercostals which arise from the transverse processes of the vertebræ, where the ribs are fixed to them, and other portions that pass over one rib and terminate in the next below it, Albinus calls Levatores costarum longiores et breviores.

The portions of the internal that pass over one rib, and are inferted into the next below it, Douglas calls Costarum depressores proprii Cowperii.

Thefe

C

a

li

These portions of both rows affist in raising the ribs in the same manner as the rest of the intercostals.

Supra costales, and Infra costales, Winslow.

The muscles within the thorax are one pair, viz.

TRIANGULARIS, or STERNO-COSTALIS,

Arises, fleshy, and a little tendinous, from all the length of the cartilago ensisormis laterally, and from the edge of the lower half of the middle bone of the sternum, from whence its sibres ascend obliquely upwards and outwards.

Inferted, generally by three triangular terminations, into the lower edge of the cartilages of the third, fourth, and fifth ribs, near where these join with the ribs.

Use. To depress these cartilages, and the extremities of the ribs; and consequently to affist in contracting

the cavity of the thorax.

This muscle often varies; and is sometimes inserted into the cartilage of the second rib, sometimes into the cartilage of the fixth rib.

# C H A P. XXIII.

Muscies fituated on the Anterior Part of the Neck close to the Vertebræ.

THESE confist of one layer formed by four mus-

## 1. LONGUS COLLI,

Arises, tendinous and fleshy, from the bodies of the three superior vertebræ of the back laterally; and from the

the transverse process of the third, fourth, fifth, and fixth vertebræ of the neck, near their roots.

Inserted into the fore-part of the bodies of all the vertebræ of the neck, by as many small tendons, which are covered with flesh.

Use. To bend the neck gradually forwards, and to one fide.

## 2. RECTUS CAPITIS INTERNUS MAJOR,

Arises, from the anterior points of the transverse processes of the third, fourth, fifth, and sixth vertebræ of the neck, by four distinct beginnings.

Inserted into the cuneiform process of the os occipitis, a little before the condyloid process.

Use. To bend the head forwards.

Rectus anterior longus, Winflow.

#### 3. RECTUS CAPITIS INTERNUS MINOR,

Arises, fleshy, from the fore-part of the body of the first vertebra of the neck, opposite to the superior oblique process.

Inserted near the root of the condyloid process of the os occipitis, under, and a little farther outwards than, the former muscle.

Use. To nod the head forwards.

Rectus anterior brevis, Winflow.

## 4. RECTUS CAPITIS LATERALIS,

Arises, fleshy, from the anterior part of the point of the transverse process of the first vertebra of the neck.

Inserted into the os occipitis, opposite to the foramen stylo-mastoideum of the temporal bone.

Use. To bend the head a little to one side. Transversalis anticus primus, Winslow.

#### C H A P. XXIV.

Muscles situated on the Posterior Part of the TRUNK.

THESE may be divided into four layers, and a fingle pair.

The first layer consists of two muscles, which cover almost the whole posterior part of the trunk.

#### I. TRAPEZIUS, SEU CUCULARIS,

Arises, By a strong round tendon, from the lower part of the protuberance in the middle of the os occipitis behind; and, by a thin membranous tendon, which covers part of the splenius and complexus muscles, from the rough curved line that extends from the protuberance towards the mastoid process of the temporal bone; runs down along the nape of the neck, where it seems to arise from its fellow, and covers the spinous processes of the superior vertebræ of the neck; but arises from the spinous processes of the two inferior, and from the spinous processes of all the vertebræ of the back; adhering, tendinous, to its sellow, the whole length of its origin.

Inserted, fleshy, into the posterior half of the clavicle; tendinous and fleshy, into the acromion, and into almost

all the spine of the scapula.

Use. Moves the scapula according to the three different directions of its fibres; for the upper descending fibres draw it obliquely upwards, the middle transverse straight fibres draw it directly backwards, and the inferior ascending fibres draw it obliquely downwards and backwards.

N.B.

N. B. Where it is inseparably united to its fellow in the nape of the neck, it is named Ligamentum Nuchæ or Colli.

#### 2. LATISSIMUS DORSI.

Arises, by a broad thin tendon, from the posterior part of the spine of the os ilium, from all the spinous processes of the os sacrum and vertebræ of the loins, and from the seven inserior ones of the vertebræ of the back; also, tendinous and sleshy, from the extremities of the three or four inserior ribs, a little beyond their cartilages, by as many distinct slips. The inserior sibres ascend obliquely, and the superior run transversely, over the inserior angle of the scapula, towards the axilla, where they are all collected, twisted, and solded.

Inserted, by a strong thin tendon, into the inner edge of the groove for lodging the tendon of the long head of the biceps.

Use. To pull the arm backwards and downwards, and

to roll the os humeri.

a

15

75

1-

n,

18

m

16

k,

ne;

e-

æ

1c

f-

g

le

11-

ds

B.

N. B. The infertion of this muscle should not be prosecuted till the muscles of the os humeri, to which it belongs, are diffected.

The fecond layer confifts of three pair, two on the back, and one on the neck:

On the back,

## 1. SERRATUS POSTICUS INFERIOR,

Arises, by a broad thin tendon, in common with that of the latissimus dorsi, from the spinal processes of the two inferior vertebræ of the back, and from the three superior of the loins.

Inserted into the lower edges of the four inferior ribs, at a little distance from their cartilages, by as many

distinct sleshy lips.

Vol. I. X Use.

a

Ì

f

Use. To depress the ribs into which it is inserted.

#### 2. RHOMBOIDEUS,

This muscle is divided into two portions.

1. Rhomboideus major, arises, tendinous, from the spinous processes of the five superior vertebræ of the back.

Inferted into all the basis of the scapula below its

fpine.

Use. To draw the scapula obliquely upwards, and

directly inwards.

2. Rhomboideus minor, arifes, tendinous, from the spinous processes of the three inferior vertebræ of the neck, and from the ligamentum nuchæ.

Inferted into the base of the scapula, opposite to its

fpine.

Use. To assist the former.

On the neck,

#### 3. SPLENIUS.

Arises, tendinous, from the four superior spinous processes of the vertebræ of the back; tendinous and sleshy, from the five inferior of the neck, and adheres sirmly to the ligamentum nuchæ. At the third vertebra of the neck, the splenii recede from each other, so that part of the complexus muscle is seen.

Inserted, by as many tendons, into the five superior transverse processes of the vertebræ of the neck; and tendinous and sleshy, into the posterior part of the masterior process, and into the os occipitis, where it joins

with the root of that process.

Use. To bring the head and upper vertebræ of the neck backwards laterally; and, when both act, to pull

the head directly backwards.

N. B. Albinus divides this muscle into two; viz. That portion which arises from the five inferior spinous processes of the neck, and is inserted into the mastoid process and os occipitis, he calls Splenius Capitis; and that

that portion which arises from the third and fourth of the back, and is inserted into the five superior transverse processes of the neck, is called by him Splenius Colli.

The fingle pair,

#### SERRATUS SUPERIOR POSTICUS,

Arises, by a broad thin tendon, from the spinous processes of the three last vertebræ of the neck, and the two uppermost of the back.

Inferted into the second, third, fourth, and fifth

ribs, by as many fleshy slips.

Use. To elevate the ribs, and dilate the thorax.

The third layer confifts of three pair on the back, and three on the neck.

On the back,

#### 1. SPÍNALIS DORSI,

Arises from the spinous processes of the two uppermost vertebræ of the loins, and the three inferior of the back, by as many tendons.

Inserted into the spinous processes of the nine uppermost vertebræ of the back, except the first, by as many

tendons.

r

S

e

d

Use. To erect and fix the vertebræ, and to assist in raising the spine.

## 2. LONGISSIMUS DORSI,

Arises, tendinous without, and fleshy within, from the side, and all the spinous processes of the os facrum; from the posterior spine of the os ilium; from all the spinous processes; and from the roots of the transverse processes of the vertebræ of the loins.

Inserted into all the transverse processes of the vertebræ of the back, chiefly by small double tendons; also by a tendinous and sleshy slip, into the lower edge of all the ribs, except the two inferior, at a little distance from their tubercles.

Use. To extend the vertebræ, and to raise and keep

the trunk of the body erect.

N. B. From the upper part of this muscle there runs up a round sleshy portion which joins with the cervicalis descendens.

#### 3. SACRO-LUMBALIS,

Arises, in common with the longissimus dorsi.

Inserted into all the ribs, where they begin to be curved forwards, by as many long and thin tendons;

and.

From the upper part of the fix or eight lower ribs, arise as many bundles of thin sleshy fibres, which soon terminate in the inner-side of this muscle, and are named Musculi ad Sacro lumbatem Accessorii.

Use. To pull the ribs down, and affift to erect the

trunk of the body.

N. B. There is a fleshy slip which runs from the upper part of this muscle into the fourth, sisth, and fixth transverse processes of the vertebræ of the neck, by three distinct tendons: it is named Cervicalis Descendens; and its use is to turn the neck obliquely backwards, and to one side.

On the neck,

#### 1. COMPLEXUS.

Arises from the transverse processes of the seven superior vertebræ of the back, and sour inferior of the neck, by as many distinct tendinous origins; in its ascent it receives a slessly slip from the spinous process of the first vertebra of the back: from these different origins it runs upwards, and is every where intermixed with tendinous sibres.

Inserted, tendinous and fleshy, into the inserior edge of the protuberance in the middle of the os occipitis,

and into a part of the curved line that runs forwards from that protuberance.

Use. To draw the head backwards, and to one fide; and, when both act, to draw the head directly backwards.

N. B. The long portion of this muscle that is situated next the spinous processes, lies more loose, and has a roundish tendon in the middle of it; for which reason Albinus calls it Biventer cervuis.

## 2. TRACHELO-MASTOIDEUS,

Arises from the transverse processes of the three uppermost vertebræ of the back, and from the five lowermost of the neck, where it is connected to the transversalis cervicis, by as many thin tendons, which unite into a belly, and run up under the splenius.

Inferted into the middle of the posterior side of the

mastoid process, by a thin tendon.

Use. To affist the complexus; but it pulls the head more to a side.

Complexus minor, seu Mastoideus lateralis, Winslow. Trachelo-mastoideus, seu Capitis par tertium Fallopii, Douglas.

## 3. LEVATOR SCAPULÆ,

Arises, tendinous and sleshy, from the transverse processes of the five superior vertebræ of the neck, by as many distinct slips, which soon unite to form a muscle that runs downwards and outwards.

Inferted, fleshy, into the superior angle of the sca-

pula.

1

Use. To pull the scapula upwards, and a little forwards.

Angularis, vulgo Levator proprius, Winflow. Elevator, seu Musculus patientia, Douglas.

The fourth layer confists of two pair on the back, two on the posterior part of the neck, four small pair X 3 situated fituated immediately below the posterior par of the occiput, and three on the side of the neck.

On the back,

#### 1. SEMI-SPINALIS DORSI,

Arises, from the transverse processes of the seventh, eighth, ninth, and tenth vertebræ of the back, by as many distinct tendons, which soon grow sleshy, and then become tendinous again; and are

Inferted into the spinous processes of all the vertebræ of the back above the eighth, and into the two

lowermost of the neck, by as many tendons.

Use. To extend the spine obliquely backwards.

Semispinalis externus, seu Transverso spinalis dorsi,
Winslow.

#### 1. MULTIFIDUS SPINÆ,

Arises from the side and spinous processes of the os sacrum, and from the posterior part of the os ilium, where it joins with the sacrum; from all the oblique and transverse processes of the vertebræ of the loins; from all the transverse processes of the vertebræ of the back, and from those of the neck, except the three first, by as many distinct tendons, which soon grow sleshy, run in an oblique direction, and are

Inferted, by distinct tendons, into all the spinous processes of the vertebræ of the loins, of the back, and

of the neck, except the first.

Use. When the different portions of this muscle act on one side, they extend the back obliquely, or move it laterally; but, if they act together on both sides, they extend the vertebræ backwards.

Transverso-spinalis lumborum, veterib. Sacer. Semi-spinalis internus, sive Transverso-spinalis dorsi. Semi-spinalis, sive Transverso-spinalis colli, Pars interna, Winslow.

Trans-

Transversalis lumborum, vulgo Sacer. Transversalis dorsi. Transversalis colli, Douglas.

On the posterior part of the neck,

#### 1. SEMI-SPINALIS COLLI,

Arises, from the transverse processes of the uppermost fix vertebræ of the back, by as many distinct tendons, ascending obliquely under the complexus.

Inferted into the spinous processes of all the vertebræ

of the neck, except the first and the last.

Use. To extend the neck obliquely backwards.

Semi-spinalis, five Transverso-spinalis colli, Winslow.

Spinalis cervicis, Albinus.

Spinalis, Douglas.

d

e

e

S

#### 2. TRANSVERSALIS COLLI,

Arises from the transverse processes of the five uppermost vertebræ of the back, by as many tendinous and sleshy origins; runs between the trachelo-mastoideus, and splenius colli and cervicalis descendens.

Inferted into the transverse processes of all the cervi-

cal vertebræ, except the first and the last.

Use. To turn the neck obliquely backward, and a little to one fide.

Below the posterior part of the occiput,

## 1. RECTUS CAPITIS POSTICUS MAJOR,

Arises,, fleshy, from the external part of the spinous process of the second vertebra of the neck; and grows broader in its ascent, which is not straight, but obliquely outwards.

Inserted, tendinous and fleshy, into the os occipitis, near the rectus capitis lateralis, and the insertion of the

obliquus capitis superior.

U/e.

Use. To pull the head backwards, and to affift a little in its rotation.

Rectus major, Winflow and Douglas.

#### 2. RECTUS CAPITIS POSTICUS MINOR,

Arises, by a narrow beginning, close by its fellow, from a little protuberance in the middle of the backpart of the first vertebra of the neck, its outer edge being covered by the rectus major.

Inserted, pretty broad, into the sides of a dimple in

the os occipitis, near its foramen magnum.

Use. To affist the rectus major in moving the head backwards.

Obliquus minor, Winflow and Douglas.

## 3. OBLIQUUS CAPITIS SUPERIOR,

Arises from the transverse process of the first vertebra of the neck.

Inferted, tendinous and fleshy, into the os occipitis behind the back-part of the mastoid process of the temporal bone, and under the infertion of the complexus muscle.

Use. To draw the head backwards.

Obliquus major, Winflow.

Obliquus superior, Douglas.

## 4. OBLIQUUS CAPITIS INFERIOR,

Arises, sleshy, from the spinous process of the second vertebra of the neck, its whole length; and, forming a thick sleshy belly, is

Inserted into the transverse process of the first verte-

bra of the neck.

Use. To give a rotatory motion to the head.

On the fide of the neck,

## 1. SCALENUS ANTICUS,

Arises from the fourth, fifth, and fixth transverse processes

processes of the first vertebra of the neck, by as many tendons.

Inserted, tendinous and fleshy, into the upper side of the first rib, near its cartilage.

Scalenus prior, Albinus.

II.

tle

w,

ke-

in

ad

e-

is

IS

Anterior portion of the first scalenus, Winslow. First scalenus, Douglas.

#### 2. SCALENUS MEDIUS.

Arises, from all the transverse processes of the vertebræ of the neck, by as many strong tendons; the nerves to the superior extremity pass between it and the former.

Inserted into the upper and outer part of the first rib, from its root, to within the distance of an inch from its cartilage.

Posterior portion of the first scalenus, Winslow. Second scalenus, Douglas.

#### 2. SCALENUS POSTICUS.

Arises from the fifth and fixth transverse processes of the vertebræ of the neck.

Inferted into the upper edge of the second rib, not far from the spine.

Posterior portion of the second scalenus, Winslow.

Third scalenus, Douglas.

Use of the three scaleni: To bend the neck to one side; or, when the neck is fixed, to elevate the ribs, and to dilate the thorax.

There are a number of small muscles situated between the spinous and transverse processes of contiguous vertebræ, which are accordingly named,

## 1. INTERSPINALES COLLI,

The space between the spinous processes of the vertebræ of the neck, most of which are bisurcated, is silled up with sleshy portions; which

Arife,

Arise, double, from the spinous process of the infe-

rior vertebræ of the neck; and ascend to be

Inserted, in the same manner, into the spinous process of the superior vertebra. They are sive in number.

Use. To draw these processes nearer to each other.

#### 2. INTERTRANSVERSALES COLLI.

They begin from the transverse process of the first vertebra of the back, and fill up the spaces between the transverse processes of the vertebræ of the neck, which are likewise bifurcated; and, consequently, there are six distinct double muscles, which

Arise from the inferior transverse process of each vertebra of the neck, and first of the back, and are

Inserted into the superior transverse processes.

Use. To draw these processes towards each other, and turn the neck a little to one side.

# 3. 4, 5. INTERSPINALES DORSI et LUMBORUM, and the INTERTRANSVERSALES DORSI,

Are rather small tendons than muscles, serving to connect the spinal and transverse processes.

## 6. INTERTRANSVERSALES LUMBORUM,

Are four distinct small bundles of slesh, which sill up the spaces between the transverse processes of the vertebræ of the loins, and serve to draw them towards each other.

## C H A P. XXV.

Muscles of the Superior Extremities.

THESE may be divided into the muscles that are situated on the scapula, on the os humeri, on the cubit or fore-arm, and on the hand.

Muscles situated on the scapula.

These are called *muscles* of the os humeri; and are three behind, one along its inferior costa, two before, and one beneath it.

Behind,

ft e h

X

r-

ıd

nd

to

ill

ne

ds

P.

#### 1. SUPRASPINATUS,

Arises, fleshy, from all that part of the base of the scapula that is above its spine; also from the spine and superior costa; passes under the acromion, and adheres to the capsular ligament of the os humeri.

Inferted, tendinous, into that part of the large protuberance on the head of the os humeri that is next the groove for lodging the tendon of the long head of the biceps.

Use. To raise the arm upwards; and, at the same time, to pull the capsular ligament from between the bones, that it may not be pinched.

#### 2. INFRASPINATUS,

Arises, fleshy, from all that part of the base of the scapula that is between its spine and inferior angle; from

b

from the spine, as far as the cervix of the scapula. The fibres ascend and descend obliquely towards a tendon in the middle of the muscle, which runs forwards, and adheres to the capsular ligament.

Inferted, by a thick and short tendon, into the upper and middle part of the large protuberance on the head

of the os humeri.

Use. To roll the humerus outwards; to affift in raifing, and in supporting it when raised; and to pull the

ligament from between the bones.

N. B. These two muscles are covered with a tendinous membrane, from which a number of their slessly sibres arise. It serves besides to strengthen their actions, and keeps them from swelling too much outwardly when in action.

#### 3. TERES MINOR,

Arises, fleshy, from all the round edge of the inferior costa of the scapula, and runs forwards along the inferior edge of the infraspinatus muscle, and adheres to the ligament.

Inserted, tendinous, into the back part of the large protuberance on the head of the os humeri, a little behind and below the termination of the last-named

muscle.

Use. To roll the humerus outwards; to draw the humerus backwards; and to prevent the ligament from being pinched between the bones.

## Along the inferior costa of the scapula,

## TERES MAJOR,

Arises, fleshy, from the inferior angle of the scapula, and from all that portion of its inferior costa that is rough and thicker than the rest; its sleshy fibres are continued over part of the infraspinatus muscle, to which they firmly adhere.

Inserted, by a broad, short, and thin tendon, into

the

the ridge at the inner fide of the groove for lodging the tendon of the long head of the biceps, along with the latissimus dorsi.

Use. To roll the humerus inwards, and to draw it backwards and downwards.

The two before the fcapula.

n d

d

1-

e

1.

y

0

d

le

n

e

0

0

#### 1. DELTOIDES,

Arises, fleshy, from all the posterior part of the clavicle that the pectoralis major does not posses; tendinous and sleshy, from the acromion, and lower margin of almost the whole spine of the scapula opposite to the insertion of the cucultaris muscle: from these origins it runs in three different directions, i. e. from the clavicle outwards and downwards; from the spine of the scapula outwards, forwards, and downwards; and from the acromion straight downwards; and is composed of a number of sasciculi, which form a strong sleshy muscle that covers the anterior part of the joint of the os humeri.

Inferted, tendinous, into a rough protuberance in the outer fide of the os humeri, near its middle, where the fibres of this muscle intermix with some part of the brachialis externus.

Use. To pull the arm directly outwards and upwards, and a little forwards or backwards, according to the different directions of its fibres.

## 2. CORACO BRACHIALIS,

Arises, tendinous and sleshy, from the fore-part of the coracoid process of the scapula; adhering, in its descent, to the short head of the biceps.

Inferted, tendinous and fleshy, about the middle of the internal part of the os humeri, near the origin of the third head of the triceps, called brachialis externus, where it sends down a thin tendinous expansion to the internal condyle of the os humeri.

Use.

Ch

ed

the

of wi

ce

TI

ter

fca

m

be

fo

CC

jo of m of It

fr

iı

P

Use. To raise the arm upwards and forwards.

N. B. There passes a nerve through this muscle, called Musculo-cutaneus.

The one beneath the scapula,

#### SUPSCAPULARIS,

Arises, fleshy, from all the base of the scapula internally, and from its superior and inferior costæ, being composed of a number of tendinous and sleshy fasciculi, which make prints on the bone; they all join together, fill up the hollow of the scapula, and pass over the joint, adhering to the capsular ligament.

Inferted, tendinous, into the upper part of the in-

ternal protuberance at the head of the os humeri.

Use. To roll the humerus inwards, and to draw it to the fide of the body; and to prevent the capsular ligament from being pinched.

## C H A P. XXVI.

Muscles situated on the Os Humeri.

# THESE are called

Muscles of the Cubit or Fore-arm.

They confist of two before, and two behind.

Before,

## 1. BICEPS FLEXOR CUBITI,

Arises, by two heads. The first and outermost, called

ed longus, begins tendinous from the upper edge of the glenoid cavity of the scapula; passes over the head of the os humeri within the joint; and, in its descent without the joint, is inclosed in a groove near the head of the os humeri, by a membranous ligament that proceeds from the capsular ligament and adjacent tendons. The second, or innermost head, called brevis, arises, tendinous and sleshy, from the coracoid process of the scapula, in common with the coraco-brachialis muscle. A little below the middle of the fore-part of the os humeri, these heads unite.

Inserted, by a strong roundish tendon, into the tu-

bercle on the upper end of the radius internally.

Use. To turn the hand supine, and to bend the fore-arm.

N. B. At the bending of the elbow, where it begins to grow tendinous, it fends off an aponeurosis, which covers all the muscles on the inside of the fore-arm, and joins with another tendinous membrane, which is sent off from the triceps extensor cubiti, and covers all the muscles on the outside of the fore-arm, and a number of the fibres, from opposite fibres, decussate each other. It serves to strengthen the muscles, by keeping them from swelling too much outwardly, when in action; and a number of their sleshy fibres take their origin from it.

Biceps brachii, Albinus. Coraco radialis, seu biceps, Winslow. Biceps internus, Douglas.

## 2. BRACHIALIS INTERNUS,

Arises, sleshy, from the middle of the os humeri, at each side of the infertion of the deltoid muscle, covering all the inferior and fore part of this bone, runs over the joint, and adheres sirmly to the ligament.

Inserted, by a strong short tendon, into the coronoid

process of the ulna.

CI

fle

na

th

fo

8

W

b

1

it

ft

Use. To bend the fore-arm, and to prevent the capfular ligament of the joint from being pinched. Brachialis, Winflow.

Behind,

#### 1. TRICEPS EXTENSOR CUBITI,

Arises, by three heads; the first called longus, pretty broad and tendinous, from the inferior costa of the scapula, near its cervix. The second head, called brewis, arises by an acute, tendinous, and sleshy beginning, from the back part of the os humeri, a little below its head, outwardly. The third, called brachialis externus, arises by an acute beginning, from the back part of the os humeri. These three heads unite lower than the insertion of the teres major, and cover the whole posterior part of the humerus, from which they receive addition in their descent.

Inferted into the upper and external part of the process of the ulna, called olecranon, and partly into the condyles of the os humeri, adhering firmly to the ligament.

Use. To extend the fore-arm.

Anconeus major, Anconeus externus, and Anconeus internus, Winflow.

Biceps externus, and Brachialis externus, Douglas.

## 3. ANCONEUS,

Arises, tendinous, from the posterior part of the external condyle of the os humeri; it soon grows fleshy, and is continued from the third head of the triceps.

Inserted, fleshy, and thin, into a ridge on the outer and posterior edge of the ulna, being continued some way below the olecranon, and covered with a tendinous membrane.

Use. To affift in extending the fore-arm.

Anconeus minor, Winflow.

Anconeus, vel Cubitalis Riolani. Douglas.

CHAP.

## C H A P. XXVII.

Muscles fituated on the Cubit or Fore-ARM \*.

THESE may be divided into three classes: first, flexors and extensors of the whole hand; fecond, flexors and extensors of the fingers; and, third, supinators and pronators, or those that roll the radius on the ulna.

First class confists of three flexors, and three extenfors.

Flexors:

II.

p-

he re-

n-

e-

lis

ck

er he

ey

ohe

a.

us

S.

х•

y,

er

ne li-

P.

#### 1. PALMARIS LONGUS,

Arises, tendinous. from the internal condyle of the os humeri, soon grows sleshy, and, after a short progress, sends off a long slender tendon.

Inferted into the ligamentum carpi annulare, and into a tendinous membrane that is expanded on the palm of the hand, named aponeurofis palmaris; which, above, begins at the transverse or annular ligament of the wrist, and, below, is fixed to the roots of the fingers.

Use. To bend the hand, and to stretch the membrane that is expanded on the palm.

Ulnaris gracilis, Winflow. You. I. Y

N.B.

\* In the following description, the arm is supposed to hang by the side with the palm turned forwards; so that the radius and thumb are upon its outer side, and the ulna and little singer upon its inner side; whereas, when the muscles are described in the less straining posture of pronation, as has been generally done by authors, the utmost confusion is necessarily in roduced in the application of the terms outer and inner, from the decussation of the radius and ulna.

tl

C

fi

ra

b

m

0

is

is

lit

N. B. This muscle is fometimes wanting; but the aponeurosis palmaris is always to be found, and a small muscle named

#### PALMARIS BREVIS,

Arises from the ligamentum carpi annulare, and tendinous membrane that is expanded on the palm of the hand.

Inserted, by small bundles of sleshy fibres, into the skin and fat that covers the abductor minimi digiti, and into the os pisiforme.

Use. To affift in contracting the palm of the hand.

Palmaris cutaneus, Winflow.

#### 2. FLEXOR CARPI RADIALIS,

Arises, tendinous and sleshy, from the internal condyle of the os humeri, and from the anterior part of the upper end of the ulna, where it firmly adheres to the pronator radii teres.

Inferted, by a flat tendon, into the fore and upper part of the metacarpal bone that sustains the fore-finger, after running through a fossa in the os trapezium.

Use. To bend the hand, and to affift in its prona-

tion.

Radialis internus, Albinus and Winflow.

## 3. FLEXOR CARPI ULNARIS,

Arises, tendinous, from the internal condyle of the os humeri. It has likewise a small sleshy beginning from the outer side of the olecranon; between which and the condyle the ulnar nerve passes to the fore-arm; and a number of its sleshy sibres arise from the tendinous membrane which covers the fore arm.

Inserted, by a short strong tendon, into the os pisforme; at a little distance from its insertion, a small ligament is sent off to the metacarpal bone that sustains

the little finger.

Use.

Use. To assist the former in bending the hand.
Ulnaris internus, Albinus and Winslow.

#### Extenfors:

II.

the

nall

enthe

the

and

on-

of

to

per

er,

na-

the

ing ich

n;

di-

ifi-

li-

ins

Je.

#### 1. EXTENSOR CARPI RADIALIS LONGIOR,

Arises, broad, thin, and fleshy, immediately below the supinator radii longus, from the lower part of the external ridge of the os humeri, above its external condyle.

Inferted, by a round tendon, into the posterior and upper part of the metacarpal bone that sustains the fore-finger.

Use. To extend and bring the hand backwards.

Radialis externus longior, Albinus. Radialis externus primus, Winflow.

#### 2. EXTENSOR CARPI RADIALIS BREVIOR,

Arises, tendinous, from the external condyle of the os humeri, and from the ligament that connects the radius to it, and runs along the outside of the radius.

Inferted, by a round tendon, into the upper and back part of the metacarpal bone that fustains the middle finger.

Use. To affist the last mentioned muscle.

Radialis externus brevior, Albinus.

Radialis secundus, Winslow.

## EXTENSOR CARPI ULNARIS,

Arises, tendinous, from the external condyle of the os humeri; and, in its progress, sleshy from the middle of the ulna, where it passes over it. Its round tendon is inclosed by a membranous sheath, in a groove which is situated at the extremity of the ulna.

Inserted, by its round tendon, into the posterior and upper part of the metacarpal bone that sustains the little-finger.

Use.

Dse. To assist the former in extending the hand. Ulnaris externus, Albinus and Winslow.

## Second Class.

The flexors and extensors of the four fingers are, two long, and one small flexor to each finger, and one extensor.

#### 1. FLEXOR SUBLIMIS PERFORATUS,

Arifes, tendinous and fleshy, from the internal condyle of the os humeri; tendinous from the coronoid process of the ulna, near the edge of the cavity that receives the head of the radius; fleshy from the tubercle of the radius; and membranous and fleshy from the middle of the fore-part of the radius, where the flexor pollicis longus arises. Its fleshy belly sends off four round tendons before it passes under the ligament of the wrist.

Inferted into the anterior and upper part of the fecond bone of each finger, being, near the extremity of the first bone, divided for the passage of the perforans.

Use. To bend the second joint or phalanx of the fingers.

Sublimis, Albinus. Perforatus, Douglas.

## 2. FLEXOR PROFUNDUS PERFORANS,

Arises, fleshy, from the external side, and upper part of the ulna, for some way downwards, and from a large share of the interosseous ligament. It splits into sour tendons, a little before it passes under the ligamentum carpi annulare; and these pass through the slits in the tendons of the slexor sublimis.

Inferted into the fore and upper part of the third or last bone of all the four fingers.

Use. To bend the last joint of the fingers.

Pro-

d

e

tl

lo

fi

u

iı

f

C

0

f

t

Profundus, Albinus, Perforans, Douglas.

II.

re,

ne

n-

oid

at

u-

om he

off

nt

fe-

ity

fo-

he

er

to a-

he

or

ra-

The four finall flexors are named

#### LUMBRICALES,

Arise, thin and fleshy, from the outside of the tendons of the flexor profundus, a little above the lower edge of the ligamentum carpi annulare.

Inferted, by long slender tendons, into the outer sides of the broad tendons of the interossei muscles, about the middle of the first joint.

Use. To increase the flexion of the fingers while the long flexors are in full action.

#### Extensors:

#### EXTENSOR DIGITORUM COMMUNIS,

Arises, by an acute, tendinous, and sleshy beginning, from the external condyle of the os humeri, where it adheres to the supinator radii brevis. Before it passes under the ligamentum carpi annulare externum, it splits into four tendons; some of which may be divided into several smaller; and about the fore-part of the metacarpal bones they remit tendinous silaments to each other.

Inferted into the posterior part of all the bones of the four fingers, by a tendinous expansion.

Use. To extend all the joints of the fingers.

## Third Clafs,

Consists of four muscles, viz. two supinators, and two pronators.

## Supinators:

## 1. SUPINATOR RADII LONGUS,

Arises, by an acute and fleshy origin, from the external

Ci

d

15

t

fi

f

p

P

b

ternal ridge of the os humeri, above the external condyle, near as far up as the middle of that bone.

Inserted into the outer-side of the inferior extremity

of the radius.

Use. To roll the radius outwards, and consequently the palm of the hand upwards.

Supinator longus, Albinus, Winflow, and Douglas.

#### 2. SUPINATOR RADII BREVIS,

Arises, tendinous, from the external condyle of the os humeri; tendinous and sleshy, from the external and upper part of the ulna, and adheres firmly to the ligament that joins these two bones.

Inserted into the head, neck, and tubercle of the radius, near the insertion of the biceps, and ridge run-

ning from that downwards and outwards.

Use. To roll the radius outwards, and so bring the hand supine.

#### Pronators:

## 1. PRONATOR RADII TERES,

Arises, fleshy, from the internal condyle of the os humeri, and tendinous from the coronoid process of the ulna.

Inferted, thin, tendinous, and fleshy, into the middle of the posterior part of the radius.

Use. To roll the radius, together with the hand, in-

wards.

## 2. PRONATOR RADII QUADRATUS,

Arifes, broad, tendinous, and fleshy, from the lower and inner part of the ulna; the fibres run transversely, to be

Inferted into the lower and anterior part of the radius, opposite to its origin.

Use. To turn the radius, together with the hand,

inwards.

CHAP.

## C H A P. XXVIII.

Muscles situated on the HAND chiefly.

THESE may be divided into four classes, viz. muscles of the thumb, fore-finger, little-finger, and meta-carpal bones.

Muscles of the Thumb.

These consist of three flexors, three extensors, one abductor, and one adductor.

#### Flexors:

#### 1. FLEXOR LONGUS POLLICIS MANUS,

Arises, by an acute sleshy beginning, from the upper part of the radius, immediately below its tubercle, and is continued down for some space on the fore-part of this bone. It has likewise generally another origin from the internal condyle of the os humeri, which forms a distinct sleshy slip that terminates near the upper part of the origin from the radius.

Inserted into the last joint of the thumb, after having passed its tendon under the ligament of the wrist.

Use. To bend the last joint of the thumb.

Flexor tertii internodii, Douglas.

## 2. FLEXOR BREVIS POLLICIS MANUS,

Arises, from the os trapezoides, magnum, and unciforme of the carpus, and is divided into two portions by the tendon of the flexor pollicis longus.

Y 4

In-

ıd,

ra-

II.

on-

ity

tly

S.

the

nal

the

the

un-

the

OS

of

dle

in-

ver ly,

P.

0

t

Inserted into the offa sesamoidea and first bone of the thumb.

Use. To bend the first joint of the thumb. Flexor secundi internodii, Douglas.

## 3. FLEXOR OSSIS METACARPI POLLICIS, OR OPPONENS POLLICIS,

Arises, fleshy, from the os trapezium and ligamentum carpi annulare, lying under the abductor pollicis.

Inserted, tendinous and fleshy, into the under and anterior part of the metacarpal bone of the thumb.

Use. To bring the thumb inwards, opposite to the other fingers.

Flexor primi internodii, Douglas.

#### Extensors:

#### 2. EXTENSOR OSSIS METACARPI POLLICIS MANUS.

Arises, fleshy, from the middle and posterior part of the ulna, immediately below the insertion of the anconæus muscle, from the posterior part of the middle of the radius, and from the interosseous ligament.

Inferted, generally by two tendons, into the os trapezium, and upper back-part of the metacarpal bone of the thumb, and often joins with the abductor pol-

licis.

De. To extend the metacarpal bone of the thumb outwardly.

Abductor longus pollicis manus, Albinus. Extensor primi internodii, Douglas.

## 2. EXTENSOR PRIMI INTERNODII,

Arises, fleshy, from the posterior part of the ulna near the former muscle, and from the interosseous ligament.

Inserted, tendinous, into the posterior part of the first bone of the thumb; and part of it may be traced as far

as the fecond bone.

II.

he

n-

S.

nd

he

S,

of o-

of

a-

ne

1-

b

a

1-

It

Use. To extend the first bone of the thumb obliquely outwards.

Extensor minor pollicis manus, Albinus. This and the preceding muscle is called Extensor pollicis primus, Winslow. Extensor secundi internodii, Douglas.

#### 3. EXTENSOR SECUNDI INTERNODII,

Arises, by an acute, tendinous, and fleshy beginning, from the middle back part of the ulna, and from the interosseous ligament; its tendon runs through a small groove at the inner and back part of the lower end of the radius.

Inserted into the last bone of the thumb.

Use. To extend the last joint of the thumb obliquely backwards.

Extensor major pollicis manus, Albinus. Extensor pollicis secundus, Winslow. Extensor tertii internodii, Douglas.

#### ABDUCTOR POLLICIS MANUS,

Arises, by a broad, tendinous, and sleshy beginning, from the ligamentum carpi annulare, and from the os trapezium.

Inserted, tendinous, into the outer side of the root of

the first bone of the thumb.

Use. To draw the thumb from the fingers.

N. B. Albinus names the inner portion of this muscle abductor brevis alter.

Abductor, Thenar Riolani, Douglas.

## ADDUCTOR POLLICIS MANUS,

Arises, fleshy, from almost the whole length of the metacarpal bone that sustains the middle singer; from thence its fibres are collected together.

Inferted, tendinous, into the inner part of the root

of the first bone.

U/c.

Use. To pull the thumb towards the fingers.

Adductor ad minimum digitum, Douglas.

Fore-finger,

#### INDICATOR,

Arises, by an acute fleshy beginning, from the middle of the posterior part of the ulna; its tendon passes under the same ligament with the extensor digitorum communis, with part of which it is

Inserted into the posterior part of the fore-finger.

Extensor secundi internodii indicis proprius, vulgo indicator, Douglas.

#### ABDUCTOR INDICIS MANUS,

Arises, from the os trapezium, and from the superior part and inner side of the metacarpal bone of the thumb.

Inserted, by a short tendon, into the outer and back part of the first bone of the fore-finger.

Use. To bring the fore-finger towards the thumb. Semi-interosseus, Winslow,

Little-finger,

## ABDUCTOR MINIMI DIGITI MANUS,

Arises, fleshy, from the os pisiforme, and from that part of the ligamentum carpi annulare next it.

Inferted, tendinous, into the inner fide of the upper end of the first bone of the little finger.

Use. To draw this finger from the rest.

Hypothenar minor, Winflow.

Extensor tertii internodii minimi digiti, Douglas.

## ADDUCTOR METACARPI MINIMI DIGITI MANUS,

Arises, fleshy, from the thin edge of the os unciforme, and from that part of the ligament of the wrist next it. Inserted, tendinous, into the inner side and anterior part of the metacarpal bone of this singer.

Use. To bend and bring the metacarpal bone of this

finger towards the rest.

Metacarpus, Winflow.

Flexor primi internodii minimi digiti, Douglas.

## FLEXOR PARVUS MINIMI DIGITI,

Arises, fleshy, from the outer side of the os unciforme, and from the ligament of the wrist which joins with that bone.

Inserted, by a rounish tendon, into the inner and anterior part of the upper end of the first bone of this finger.

Use. To bend the little finger, and affift the adduc-

tor.

II.

le

nn-

n-

or

k

at

er

1-

Abductor minimi digiti, Hypothenar Riolani, Dou-

Between the metacarpal bones, there are four internal and three external muscles, named interossei.

## Interossei interni:

#### 1. PRIOR INDICIS,

Arises, tendinous and sleshy, from the upper and outer part of the metacarpal bone that sustains the fore-finger.

Inferted into the outside of that part of the tendinous expansion from the extensor digitorum communis, which covers the posterior part of the fore-finger.

Use. To draw the fore-finger inwards towards the

thumb, and extend it obliquely.

Extensor tertii internodii indicis, Douglas.

## 2. POSTERIOR INDICIS,

Arises, tendinous and fleshy, from the root and inner part

C

fr

tl

W

C

part of the metacarpal bone that fustains the fore-

finger.

Inserted into the inner side of the tendinous expansion which is sent off from the extensor digitorum communis, along the posterior part of the fore-singer.

Use. To extend the fore-finger obliquely, and to

draw it outwards.

First interosseus, Douglas.

#### 3. PRIOR ANNULARIS.

Arises, from the root of the outside of the metacar-

pal bone that fustains the ring-finger.

Inferted into the outfide of the tendinous expansion of the extensor digitorum communis which covers the ring-finger.

Use. To extend and pull the ring-finger towards the

thumb.

Fourth interoffeus, Douglas.

#### 4. INTEROSSEUS AURICULARIS,

Arises from the root and outer side of the metacarpal

bone of the little finger; and is

Inferted into the outside of the tendinous expansion of the extensor digitorum communis, which covers the posterior part of the little finger.

Use. To extend and draw the little finger outwards.

Sixth interoffeus, Douglas.

Interossei externi, seu bicipites:

## r. PRIOR MEDII,

Arises, by two origins, from the roots of the metacarpal bones that sustain the fore and middle singers externally, and next each other: Runs along the outside of the middle-singer; and, being conspicuous on both sides of the hand, is

Inferted into the outside of the tendinous expansion from

from the extensor digitorum communis, which covers the posterior part of the middle-finger.

Ufe. To extend, and to draw the middle finger in-

wards.

Π.

e-

1-

n-

to

r.

n

e

e

Second interoffeus, Douglas.

#### 2. POSTERIOR MEDII,

Arises, by two origins, from the roots of the metacarpal bones, next each other, that sustain the middle and ring-fingers.

Inserted into the inside of the tendinous expansion from the extensor digitorum communis, which runs

along the posterior part of the middle-finger.

Use. To extend and draw the middle-finger outwards.

Third interosseus, Douglas.

#### 3. POSTERIOR ANNULARIS,

Arises, by two origins, from the roots of the metacarpal bones that sustain the ring and little fingers next each other.

Inserted into the infide of the tendinous expansion of the extensor digitorum communis, which runs along the posterior part of the ring-finger.

Use. To extend and draw the ring-finger inwards.

Fifth interoffeus, Douglas.

N. B. The internal interoffei are only conspicuous on the palm of the hand; but the external are apparent on both the palm and back of the hand.

#### C H A P. XXIX.

Muscles of the Inferior Extremities.

THESE may be divided into the muscles situated on the outside of the pelvis, on the thigh, on the leg, and on the foot.

Muscles on the outside of the pelvis, which are called muscles of the thigh.

These are composed of one layer before and three layers behind.

The layer before confifts of five muscles:

1. PSOAS MAGNUS.
2. ILIACUS INTERNUS.

These were formerly deferibed.

Vid. p. 313. 314.

## 3. PECTINALIS,

Arises, broad and fleshy, from the upper and anterior part of the os pubis or pectinis, immediately above the foramen thyroideum.

Inferted into the anterior and upper part of the linea afpera of the os femoris, a little below the trochanter minor, by a flat and short tendon.

Use. To bring the thigh upwards and inwards, and to give it a degree of rotation outwards.

Pectineus, Albinus.

Ú.

n

d

rs

r

## 4. TRICEPS ADDUCTOR FEMORIS,

Under this appellation are comprehended three diflinct muscles;

#### 1. ADDUCTOR LONGUS FEMORIS,

Arises, by a pretty strong roundish tendon, from the upper and anterior part of the os pubis, and ligament of its synchondrosis, on the inner-side of the pectinalis.

Inserted, tendinous, near the middle of the posterior part of the linea aspera, being continued for some way down.

Adductor femoris primus, Douglas. Triceps minus, Winslow.

#### 2. ADDUCTOR BREVIS FEMORIS,

Arises, tendinous, from the os pubis near its joining with the opposite os pubis below and behind the former.

Inferted, tendinous and fleshy, into the inner and upper part of the linea aspera, from a little below the trochanter minor, to the beginning of the insertion of the adductor longus.

Adductor femoris secundus, Douglas. Triceps secundus, Winslow.

## 3. ADDUCTOR MAGNUS FEMORIS,

Arises, a little lower down than the former, near the symphysis of the ossa pubis; tendinous and sleshy, from the tuberosity of the os ischium; the sibres run outwards and downwards.

Inferted, into almost the whole length of the linea aspera; into a ridge above the internal condyle of the os femoris; and, by a roundish long tendon, into the upper part of that condyle, a little above which the femoral artery takes a spinal turn towards the ham, passing between this muscle and the bone.

Use of these three muscles or triceps. To bring the

Ch

the

101

to

18

do

wa

ce

fp

ar

ft

ba

b

b

thigh inwards and upwards, according to the different directions of their fibres; and, in some degree, to roll the thigh outwards.

Adductor femoris tertius, and Adductor femoris quartus, Douglas. Triceps tertius, Winflow.

#### 5. OBTURATOR EXTERNUS,

Arises, fleshy, from the lower fore-part of the os pubis, and fore-part of the inner crus of the ischium; surrounds the foramen thyroideum; a number of its sibres, arising from the membrane which fills up that foramen, are collected like rays towards a centre, and pass outwards around the root of the back-part of the cervix of the os semoris.

Inserted, by a strong tendon, into the cavity at the inner and back part of the root of the trochanter major, adhering in its course to the capsular ligament of the thigh-bone.

Use. To roll the thigh-bone obliquely outwards, and to prevent the capsular ligament from being pinched.

Behind:

## First Layer.

## GLUTEUS MAXIMUS,

Arises, fleshy, from the posterior part of the spine of the os ilium, a little higher up than the joining of the ilium with the os sacrum, from the whole external side of the os facrum, below the posterior spinous process of the os ilium; from the posterior facro-ischiatic ligament, over which part of the inferior edge of this muscle hangs in a folded manner; from the os coccygis. All the sleshy sibres run obliquely forwards, and a little downwards, to form a thick broad muscle, which is divided into a number of strong sasciculi. The upper part of it covers almost the whole of the trochanter major,

major, between which and the tendon of this muscle there is a large bursa mucosa, and where it is inseparably joined to the broad tendon of the tensor vaginæ femoris.

Inferted, by a strong, thick, and broad tendon into the upper and outer part of the linea aspera, which is continued from the trochanter major, for some way downwards.

Use. To extend the thigh, by pulling it directly backwards, and a little outwards.

Gluteus magnus, Albinus. Gluteus major, Cowper.

M.

ent

roll

u-

n;

its

nd

he

he

or,

he

nd

of

ne

le of

a-

is

y-

d

h

0.

er

r,

## Second Layer,

#### GLUTEUS MEDIUS,

Arises, fleshy, from the anterior superior spinous process of the os ilium, and from all the outer edge of the spine of the ilium, except its posterior part, where it arises from the dorsum of that bone.

Inserted, by a broad tendon, into the outer and poflerior part of the trochanter major.

Use. To draw the thigh-bone outwards, and a little backwards; to roll the thigh-bone outwards, especially when it is bended.

N. B. The anterior and upper part of this muscle is covered by a tendinous membrane, from which a number of its sleshy sibres arise, and which joins with the broad tendons of the gluteus maximus, tensor vaginæ semoris, and latissimus dorsi.

## Third Layer confifts of four muscles.

#### 1. GLUTEUS MINIMUS,

Arises, fleshy, from a ridge that is continued from the superior anterior spinous process of the os ilium, and from the middle of the dorsum of that bone, as far back as its great nitch.

Inserted, by a strong tendon, into the fore and upper part of the trochanter major.

Vol. I. Z Use.

Ch

tub

the

fm

for

tua

off

an

th

du

ce

lit

th

he

of

k

Use. To assist the former in pulling the thigh out. wards and backwards, and in rolling it.

Gluteus minor, Albinus.

#### 2. PYRIFORMIS,

Arises, within the pelvis, by three tendinous and fleshy origins, from the second, third, and sourth pieces of the os sacrum; from thence growing gradually narrower, it passes out of the pelvis along with the posterior crural nerve, below the nitch in the posterior part of the os ilium, where it receives a few sleshy fibres.

Inferted, by a roundish tendon, into the upper part of the cavity at the inner side of the root of the trochanter major.

Use. To move the thigh a little upwards, and roll it

outwards.

Pyriformis, seu iliacus externus, Douglas.

#### 3. GEMINI

Arises, by two distinct origins; the superior from the spinous process, and the inferior from the tuberosity of the os ischium; also, from the posterior facro-ischiatic ligament. They are both united by a tendinous and sleshy membrane, and form a purse for the tendon of the obturator internus muscle, which was formerly described.

Inferted, tendinous and fleshy, into the cavity at the inner side of the root of the trochanter major, on each side of the tendon of the obturator internus, to which

they firmly adhere.

Use. To roll the thigh outwards, and to preserve the tendon of the obturator internus from being hurt by the hardness of that part of the ischium over which it passes; also, to hinder it from starting out of its place, while the muscle is in action.

Gemelli, Winflow.

4. QUA-

II.

out.

and

oie.

ally

poior fhy

of

1 it

mo

ity ia-

nd

of

le-

he

ch

ch

he

he

ile

1.

## 4. QUADRATUS FEMORIS,

Arises, tendinous and fleshy, from the outside of the tuberosity of the os ischium; and, running transversely, is Inserted, fleshy, into a rough ridge, continued from the root of the large trochanter to the root of he small one.

Use. To roll the thigh outwards.

## C H A P. XXX.

Muscles situated on the Thigh.

THESE are called muscles of the leg; and confift of one, on the outside; two, on the inside; four, before; and, four behind.

Previous to the description of the muscles that are situated on the thigh and leg, it is necessary to take notice of a broad tendinous fascia or sheath, which is sent off from the back and from the tendons of the glutei and adjacent muscles.

It is a strong thick membrane on the outside of the thigh and leg; but, towards the inside of both, it gradually turns thinner, and has rather the appearance of cellular substance, than a tendinous membrane. A little below the trochanter major, it is sirmly fixed to the linea aspera; and, farther down, to that part of the head of the tibia that is next the sibula; where it sends off the tendinous expansion along the outside of the leg.

It serves to strengthen the action of the muscles, by keeping them firm in their proper places while in ac-

tion, particularly the tendons that pass over the joints, where this membrane is thickest, and it gives origin to a number of the sleshy fibres of the muscles.

Outfide,

#### TENSOR VAGINÆ FEMORIS,

Arises, by a narrow, tendinous, and fleshy beginning, from the external part of the anterior superior spinous process of the os ilium.

Inferted, a little below the trochanter major, into the inner fide of the membranous fascia which covers the

outfide of the thigh.

Use. To stretch the membranous fascia, to affist in the abduction of the thigh, and somewhat in its rotation inwards.

Musculus aponeurosis, vel Fasciæ latæ, Winslow.

Infide,

## 1. SARTORIUS,

Arises, tendinous, from the anterior superior spinous process of the os ilium, soon grows sleshy, runs down for some space upon the rectus, and, going obliquely inwards, it passes over the vastus internus, and, about the middle of the os semoris, over part of the triceps, it runs down further between the tendon of the adductor magnus, and that of the gracilis muscle.

Inserted, by a broad and thin tendon, into the inner fide of the tibia, near the inserior part of its tubercle.

Use. To bend the leg obliquely inwards, or to bring one leg across the other.

## 2. GRACILIS,

Arises, by a thin tendon, from the os pubis near the symphysis of these two bones; soon grows sleshy; and, descending by the inside of the thigh, is

Inferted, tendinous, into the tibia under the fartorius.

Ufe.

ce

th

th

th

fre

la.

pa

m

by

ol

ti

li

W

te

Co fi:

P

Use. To affist the fartorius.

Gracilis internus, sive Rectus internus, Winslow.

Before,

II.

ts,

to

ıg,

us

he

he

he

on

us

vn :ly

ut

it

or

er

ng

ne

d,

IS.

le.

#### I. RECTUS.

Arises, fleshy, from the inferior anterior spinous process of the os ilium, and tendinous from the dorsum of the ilium, a little above the acetabulum; runs down over the anterior part of the cervix of the os semoris, the sibres not being straight, but running down like the plumage of a feather obliquely outwards and inwards, from a tendon in the middle.

Inferted, tendinous, into the upper part of the patella, from which a thin tendon ruus down, on the forepart of this bone, to terminate in a thick strong ligament, which is sent off from the inferior part of the patella, and inserted into the tubercle of the tibia.

Use. To extend the leg, and in a powerful manner, by the intervention of the patella, like a pulley.

Rectus, five Gracilis anterior, Winflow.

#### 2. VASTUS EXTERNUS,

Arises, broad, tendinous, and sleshy, from the root of the trochanter major, and upper part of the linea aspera, its origin being continued from near the infertion of the gluteus minimus, the whole length of the linea aspera, by sleshy fibres which run obliquely forwards to a middle tendon, where they terminate.

Inserted into a large share of the upper part of the patella; and part of it ends in an aponeurosis, which is continued down to the leg, and, in its passage, is firmly fixed to the head of the tibia.

Use. To extend the leg.

## 3. VASTUS INTERNUS,

Arises, tendinous and sleshy, from between the forepart of the os semoris and root of the trochanter minor, Z<sub>3</sub> and

C

0

fl

10

lo

tl

b

u

if

n

n

g

t

I

] i

and from almost all the inside of the linea aspera, by fibres running obliquely forwards and downwards.

Inferted, tendinous, into the upper and infide of the patella, continuing fleshy lower than the vastus externus. Part of it likewise ends in an aponeurosis continued down to the leg, and fixed in its passage to the upper part of the tibia.

Use. To extend the leg.

#### 4. CRURALIS,

Arises, fleshy, from between the two trochanters of the os semoris, but nearer the minor, and firmly adhering to most of the fore-part of the os semoris, and connected to both vasti muscles.

Inserted, tendinous, into the upper part of the pa-

tella, behind the rectus.

Ufe. To affift in the extension of the leg.

Cruralis, Albinus.

N. B. These four muscles before, being inserted into the patella, have the same effect upon the leg, as if they were immediately inserted into it, by means of the strong tendon, or rather ligament, which is sent of from the inserior part of the patella to the tibia.

Behind,

#### 1. SEMITENDINOSUS.

Arises, tendinous and sleshy, in common with the long head of the biceps, from the posterior part of the tuberosity of the os ischium; and sending down a long roundish tendon, which ends slat, is

Inserted into the inside of the ridge of the tibia, a

little below its tubercle.

Use. To bend the leg backwards, and a little inwards. Seminervosus, Winslow and Douglas.

## 2. SEMIMEMBRANOSUS,

Arises, tendinous, from the upper and posterior part

of the tuberofity of the os ischium, sends down a broad flat tendon, which ends in a fleshy belly, and, in its descent, runs at first on the fore-part of the biceps, and, lower, between it and the semitendinosus.

Inferted, tendinous, into the inner and back part of

the head of the tibia.

tII.

by

the

cter-

onti-

the

s of ad-

and

pa-

into

s if

the

of

the

the

1, a

rds.

part

of

Use. To bend the leg, and bring it directly backwards.

N. B. The two last form what is called the inner ham-string.

#### 3. BICEPS FLEXOR CRURIS,

Arises, by two distinct heads. The first, called longus, arises, in common with the semitendinosus, from the upper and posterior part of the tuberosity of the os ischium. The second, called brevis, arises from the linea aspera, a little below the termination of the gluteus maximus, by a sleshy acute beginning, which soon grows broader as it descends to join with the first head, a little above the external condyle of the os semoris.

Inserted by a strong tendon, into the upper part of

the head of the fibula.

Use. To bend the leg.

Biceps cruris, Albinus.

Biceps, Winflow and Douglas.

N. B. This muscle forms what is called the outer ham-string; and between it and the inner, the nervous popliteus, and arteria and vena poplitea, are situated.

## 4. POPLITEUS,

Arises, by a round tendon, from the lower and backpart of the external condyle of the os semoris; then runs over the ligament that involves the joint, firmly adhering to it, and part of the semilunar cartilage. As it runs over the joint, it becomes sleshy, and the sibres run obliquely inwards, being covered with a thin tendinous membrane.

Inferted, broad, thin, and fleshy, into a ridge at the

th on med bid

d

C

a

a

the upper and internal edge of the tibia, a little below its head.

Use. To affist in bending the leg, and to prevent the capsular ligament from being pinched. After the leg is bent, this muscle serves to roll it inwards.

## C H A P. XXXI.

Muscles situated on the Leg.

THESE are called Muscles of the Foot; and may be divided into two classes, viz. Extensors and Flexors of the Foot. 2. Common Extensors and Flexors of the Toes.

## Firft Clafs.

Extensors.

These consist of three,

## 1. GASTROCNEMIUS EXTERNUS, SEU GEMELLUS,

Arises, by two distinct heads. The first head arises from the upper and back-part of the internal condyle of the os semoris, and from that bone, a little above its condyle, by two distinct tendinous origins. The second head arises tendinous from the upper and back-part of the external condyle of the os semoris. A little below the joint, their sleshy bellies unite in a middle tendon; and, below the middle of the tibia, it sends off a broad thin tendon, which joins a little above the extremity of the tibia with the tendon of the following.

2. SO-

II.

Mc

he

is

#### 2. SOLEUS, SEU GASTROCNEMIUS INTERNUS.

Arises by two origins. The first is from the upper and back part of the head of the fibula, continuing to receive many of its sleshy fibres from the posterior part of that bone for some space below its head. The other origin begins from the posterior and upper part of the middle of the tibia; and runs inwards along the inferior edge of the popliteus towards the inner part of the tibia, from which it receives sleshy fibres for some way down. The slesh of this muscle, covered by the tendon of the gemellus, runs down near as far as the extremity of the tibia; a little above which the tendons of both gastrocaemii unite, and form a strong round chord, which is called tendo Achillis.

Inserted into the upper and posterior part of the os calcis; by the projection of which the tendo Achillis is at a considerable distance from the tibia.

Use. To extend the foot, by bringing it backwards and downwards.

Gemellus and Soleus, Albinus.

Gastrocnemii and Soleus, Winflow.

Extensor tarsi suralis, vel Extensor magnus, Douglas.

# 3. PLANTARIS,

Arises, thin and sleshy, from the upper and back part of the root of the external condyle of the os femoris, near the inferior extremity of that bone, adhering to the ligament that involves the joint in its defeent. It passes along the second origin of the soleus, and under the gemellus, where it sends off a long, slender, thin tendon, which comes from between the great extensors, where they join tendons; then runs down by the inside of the tendo Achilles.

Inferted into the infide of the posterior part of the os

calcis, below the tendo Achillis.

Use. To affist the former, and to pull the capsular ligament

ligament of the knee from between the bones. It feems likewise to affist in rolling the foot inwards.

Tibialis gracilis, vulgo Plantaris, Winflow. Extensor tarsi minor, vulgo Plantaris, Douglas.

N. B. This muscle, though seldom, has been found wanting on both sides.

#### Flexors:

These consist of four; two that belong to the tibia, and two to the fibula.

#### 1. TIBIALIS ANTICUS,

Arises, tendinous and fleshy, from the middle of that process of the tibia, to which the sibula is connected above; then it runs down fleshy on the outside of the tibia; from which, and the upper-part of the interosfeous ligament, it receives a number of distinct fleshy fibres; near the extremity of the tibia, it sends off a strong round tendon, which passes under part of the ligamentum tars annulare near the malleolus internus.

Inferted, tendinous, into the infide of the os cuneiforme internum, and posterior end of the metatarsal

bone that fultains the great toe.

Use. To bend the foot, by drawing it upwards, and, at the same time, to turn the toes inwards.

#### 2. TIBIALIS POSTICUS,

Arises, by a narrow sleshy beginning, from the fore and upper part of the tibia, just under the process which joins it to the sibula; then passing through a perforation in the upper part of the interosseus ligament, it continues its origin from the back-part of the sibula next the tibia, and trom near one half of the upper-part of the last-named bone; as also, from the interosseous ligament, the sibres running towards a middle tendon, which sends off a round one that passes in a groove behind the malleolus internus.

II.

ms

nd

ia,

at

ed

he

of-

a

li-

i-

al

d,

s a

1-

e

e

e

a

S

Inferted, tendinous, into the upper and inner part of the os naviculare, being further continued to the os cuneiforme internum and medium; besides, it gives some tendinous filaments to the os calcis, os cuboides, and to the root of the metatarsal bone that sustains the middle toe.

Use. To extend the foot, and to turn the toes inwards.

# 3. PERONEUS LONGUS,

Arises, tendinous and fleshy, from the fore-part of the head of the perone, or sibula, the sibres running straight down; also from the upper and external part of the sibula, where it begins to rise into a round edge; as also, from the hollow between that and its anterior edge, as far down as to reach within a handbreadth of the ankle, by a number of sleshy sibres, which run outwards towards a tendon, that sends off a long round one, which passes through a channel at the outer ankle, in the back-part of the inferior extremity of the sibula; then, being reslected to the sinussity of the os calcis, it runs along a groove in the os cuboides, above the muscles in the sole of the foot.

Inserted, tendinous, into the outside of the root of the metatarsal bone that sustains the great toe, and by some tendinous sibres in the os cuneiforme internum.

Use. To move the foot outwards, and to extend it a little.

Peroneus maximus, vulgo Peroneus posterior, Winslow. Peroneus primus, seu Posticus, Douglas.

#### 4. PERONEUS BREVIS,

Arises, by an acute fleshy beginning, from above the middle of the external part of the sibula; from the outer side of the anterior spine of this bone; as also, from its round edge externally, the sibres running obliquely outwards towards a tendon on its external side: it sends off a round tendon which passes through the groove

364

groove at the outer ankle, being there included under the same ligament with that of the preceding muscle; and a little further, it runs through a particular one of its own.

Inserted, tendinous, into the root and external part of the metatarsal bone that sustains the little toe.

Use To affift the former in pulling the foot outwards, and extending it a little.

Peroneus medius, vulgo Peroneus anticus, Winflow. Peroneus fecundus, seu Anticus, Douglas.

## Second Class.

Common Extensors.

These confist of two.

#### 1. EXTENSOR LONGUS DIGITORUM PEDIS,

Arises, tendinous and sleshy, from the upper and outer part of the head of the tibia, and from the head of the sibula where it joins with the tibia, and from the interosseous ligament; also from the tendinous sascia, which covers the upper and outside of the leg by a number of sleshy fibres; and tendinous and sleshy, from the anterior spine of the sibula, almost its whole length, where it is inseparable from the peroneus tertius. It splits into four round tendons, under the ligamentum tarsi annulare.

Inserted, by a flat tendon, into the root of the first joint of each of the four small toes; and is expanded over the upper-side of the toes, as far as the root of the last joint.

Uje. To extend all the joints of the four small toes.

Extensor longus, Douglas.

N. B. A portion of this muscle, which

Arises from the middle of the fibula, continues down to near its inferior extremity, and fends its fleshy fibres

for

Ch

fuf

pe div

fm no th

1.

te ca

le

t

forwards to a tendon, which passes under the annular

ligament, and is

r

if

rt

5,

Inserted into the root of the metatarsal bone that sustains the little toe: it is called by Albinus peroneus tertius; and by others, the Nonus Vesalii.

Use. To affift in bending the foot.

#### z. EXTENSOR BREVIS DIGITORUM PEDIS,

Arises, fleshy and tendinous, from the fore and upper part of the os calcis; and soon forms a sleshy belly, divisible into four portions, which sends off an equal number of tendons that pass over the upper part of the foot, under the tendons of the former.

Inferted, by four flender tendons, into the tendinous expansion from the extensor longus, which covers the small toes, except the little one; also into the tendinous expansion from the extensor pollicis, that covers the upper part of the great toe.

Use. To extend the toes. Extensor brevis, Douglas.

Flexors:

These may be reckoned three.

# 1. FLEXOR BREVIS DIGITORUM PEDIS, SUBLIMIS, PERFORATUS,

Arises, by a narrow fleshy beginning, from the inferior and posterior part of a protuberance of the os calcis, between the abductors of the great and little toes; soon forms a thick fleshy belly, which sends off four tendons that split for the passage of the flexor longus.

Inserted into the second phalanx of the four lesser toes. The tendon of the little toe is often wanting.

Use. To bend the second joint of the toes. Perforatus, seu Sublimis, Douglas.

Cha

the

vifi

the

COV

the

tl

f

# 2. FLEXOR LONGUS DIGITORUM PEDIS, PROFUNDUS, PERFORANS,

Arises, by an acute tendon, which soon becomes fleshy, from the back part of the tibia, some way below its head, near the entry of the medullary artery; which beginning, is continued down the inner edge of this bone by short sleshy sibres, ending in its tendon; also, by tendinous and sleshy sibres, from the outer edge of the tibia; and between this double order of sibres the tibialis posticus muscle lies inclosed. Having passed under two annular ligaments, it then passes through a sinuosity at the inside of the os calcis; and, about the middle of the sole of the foot, divides into four tendons, which pass through the slits of the perforatus; and, just before its division, it receives a considerable tendon from that of the slexor pollicis longus.

Inferted into the extremity of the last joint of the

four leffer toes.

Use. To bend the last joint of the toes.

This mufcle is affifted by the

# FLEXOR DIGITORUM ACCESSORIUS, SEU MASSA CARNEA JACOBI SYLVII,

Arises, by a thin fleshy origin, from most part of the sinuosity at the inside of the os calcis, which is continued forwards for some space on the same bone; also, by a thin tendinous beginning, from before the tuberosity of the os calcis externally; and, soon becoming all fleshy, is

Inserted into the tendon of the flexor longus, just at

its division into four tendons.

Use. To affift the flexor longus.

## 3. LUMBRICALES PEDIS,

Arise, by four tendinous and fleshy beginnings, from

the tendon of the flexor profundus, just before its division, near the infertion of the massa carnea.

Inserted, by four slender tendons, into the inside of the first joint of the four lesser toes, and are lost in the tendinous expansion that is fent from the extensors to cover the upper part of the toes.

Use. To increase the flexion of the toes, and to draw

them inwards.

I.

V-

S:

f

r

1,

0

.

e

#### C H A P. XXXII.

Muscles which are chiefly fituated on the Foot.

THESE may be divided into the muscles of the great toe, of the little toe, and of the metatarsal bones.

Muscles of the great toe.

These are five:

# 1. EXTENSOR PROPRIUS POLLICIS PEDIS,

Arises, by an acute, tendinous, and sleshy beginning, some way below the head and anterior part of the sibula, along which it runs to near its lower extremity, connected to it by a number of sleshy sibres, which descend obliquely towards a tendon.

Inserted, tendinous, into the posterior part of the

first and last joint of the great toe.

Use. To extend the great toe. Extensor longus, Douglas.

#### 2. FLEXOR LONGUS POLLICIS PEDIS,

Arises, by an acute, tendinous, and sleshy beginning, from the posterior part of the sibula, some way below its head, being continued down the same bone, almost to its inferior extremity, by a double order of oblique sleshy sibres; its tendon passes under an annular ligament at the inner ankle.

Inserted into the last joint of the great toe, and gene-

rally fends a small tendon to the os calcis.

Use. To bend the last joint of this toe.

Flexor longus, Douglas.

#### 3. FLEXOR BREVIS POLLICIS PEDIS,

Arises, tendinous, from the under and fore-part of the os calcis, where it joins with the os cuboides, from the os cuneiforme externum, and is inseparably united with the abductor and adductor pollicis.

Inserted into the external os sesamoideum and root of

the first joint of the great toe.

Use. To bend the first joint.

## 4. ABDUCTOR POLLICIS PEDIS,

Arises, fleshy, from the inside of the root of the protuberance of the os calcis, where it forms the heel; and tendinous from the same bone, where it joins with the os naviculare.

Inferted, tendinous, into the internal os fesamoideum, and root of the first joint of the great toe.

Use. To pull the great toe from the rest.

Thenar, Winflow.

## 5. ADDUCTOR POLLICIS PEDIS,

Arises, by a long thin tendon, from the os calcis, from the os cuboides, from the os cuneiforme externum, and from the root of the metatarfal bone of the second toe.

ſ

b

f

b

Inserted into the external os sesamoideum, and root of the metatarsal bone of the great toe.

Use. To bring this toe nearer the rest.

Antithenar, Winflow.

Muscles of the little toe.

These, besides the common extensors and slexors, are two, viz.

#### 1. ABDUCTOR MINIMI DIGITI PEDIS,

Arises, tendinous and sleshy, from the semicircular edge of a cavity on the inferior part of the protuberance of the os calcis, and from the root of the metatarsal bone of the little toe.

Inferted into the root of the first joint of the little toe

externally.

I.

n.

ay

e, of

u.

e-

of

m

ed

of

ro-

nd

OS

oi-

cis,

er-

the

In-

Uses. To draw the little toe outwards from the rest. Parathenar major, and Metatarseus, Winslow.

#### 2. FLEXOR BREVIS MINIMI DIGITI PEDIS,

Arises, tendinous, from the os cuboides, near the sulcus or surrow for lodging the tendon of the peroneus longus; sleshy from the outside of the metatarsal bone that sustains this toe, below its protuberant part.

Inserted into the anterior extremity of the metatar-

fal bone, and root of the first joint of this toe.

Use. To bend this toe.

Parathenar minor, Winflow.

Muscles from the metatarfal bones.

These are four external and three internal interossei, and one muscle which is common to all the metatarfal bones.

Vol. I.

Aa

In-

T

fi

fe

fi

it

fi

ta

ri

ar

ac

of lit

# Interoffei Pedis Externi, Bicipites.

#### 1. ABDUCTOR INDICIS PEDIS,

Arises, tendinous and sleshy, by two origins, from the root of the inside of the metatarsal bone of the fore-toe, from the outside of the root of the metatarsal bone of the great-toe, and from the os cuneiforme internum.

Inserted, tendinous, into the inside of the root of the

first joint of the fore-toe.

Use, To pull the fore-toe inwards from the rest of the small toes.

#### 2. ADDUCTOR INDICIS PEDIS,

Arises, tendinous and fleshy, from the roots of the metatarsal bones of the fore and second toe.

Inserted, tendinous, into the outside of the root of the first joint of the fore-toe.

Use. To pull the fore-toe outwards towards the rest.

# 3. ADDUCTOR MEDII DIGITI PEDIS,

Arises, tendinous and fleshy, from the roots of the metatarsal bones of the second and third toes.

Inserted, tendinous, into the outside of the root of the first joint of the second toe.

Use. To pull the second toe outwards.

# 4. ADDUCTOR TERTII DIGITI PEDIS,

Arifes, tendinous and fleshy, from the roots of the metatarfal bones of the third and little toe.

Inferted, tendinous, into the outfide of the root of the first joint of the third toe.

Use. To pull the third toe outwards.

# Interossei Pedis interni.

#### 1. ABDUCTOR MEDII DIGITI PEDIS,

Arises, tendinous and fleshy, from the inside of the root of the metatarfal bone of the middle toe internally.

Inserted, tendinous, into the inside of the root of the first joint of the middle toe.

Use. To pull the middle toe inwards.

#### 2. ABDUCTOR TERTII DIGITI PEDIS,

Arises, tendinous and fleshy, from the inside and inferior part of the root of the metatarsal bone of the third toe.

Inserted, tendinous, into the inside of the root of the first joint of the third toe.

Use. To pull the thirdstoe inwards.

f

f

### 3. ADDUCTOR MINIMI DIGITI PEDIS,

Arises, tendinous and sleshy, from the inside of the toot of the metatarsal bone of the little toe.

Inserted, tendinous, into the infide of the root of the first joint of the little toe.

Use. To pull the little toe inwards.

The muscle which brings the extremities of the metatarfal bones towards each other, is named

## TRANSVERSALIS PEDIS,

Arises, tendinous, from the under part of the anterior extremity of the metatarfal bone of the great toe, and from the internal os sesamoideum of the first joint, adhering to the adductor pollicis.

Inferted, tendinous, into the under and outer part of the anterior extremity of the metatarfal bone of the little toe, and ligament of the next toe.

Aa2

Use.

Use. To contract the foot, by bringing the great toe and the two outermost toes nearer each other.

N. B. The muscles situated on the sole of the foot are covered by a strong tendinous aponeurosis, which is extended from the os calcis to the first joints of all the toes, and serves to preserve the subjacent parts from being compressed in standing and walking.

A

L

H

at

ot

all m

A

# SYSTEM OF ANATOMY.

## PART III.

CONTAINING

EIGHT ANATOMICAL TABLES

OFTHE

# HUMANBODY;

Exhibiting the PRINCIPAL PARTS of the

SKELETON AND MUSCLES

REPRESENTED IN THE

LARGE TABLES OF ALBINUS.

TO WHICH ARE ADDED,

CONCISE EXPLANATIONS.

By JOHN INNES.

# MEDICAL SOCIETY.

GENTLEMEN,

I EMBRACE this opportunity of expressing my gratitude for the many favours I have received from you, both in your collective and individual capacity; but I am unwilling to offend, either by encomiums on the utility of your institution, or by exhibiting the private friendships with which I

have heen honoured by many of your number.

In compliance with your folicitations, I lately published a short Description of the Human Muscles, which has been fortunate enough to receive your approbation. It was, however, your opinion, that a set of Tables would render the descriptions still more perfect and useful. I have, therefore, caused the following plates to be engraven; and I hope they will not altogether disappoint your expectations.

The tables of Albinus, though accurate and complete, are not, perhaps, on account of their fize, the prolixity of the descriptions, and the number of references, so fully suited to the purposes of private diffection as could be wished.

To remedy this inconvenience, I have copied eight of Albinus's tables on a smaller scale. Two of them contain a fore and back view of the skeleton; and the other fix are representations of the muscles.

To each plate I have prefixed an explanation; and I have avoided, as much as was confiftent with perspicuity, loading the figures with an unnecessary quantity of referring letters. For this purpose, I have all along made use of one alphabet only.

To prevent the defacing of the muscles, and to enable the learner to distinguish them without the assistance of referen-

ces, the figures are only lettered on one fide.

If this little work be of use in facilitating your anatomical studies, it will give the greatest pleasure to,

Gentlemen,

Your very much obliged humble fervant, and Fellow-member,

JOHN INNES.

# EXPLANATIONS

OFTHE

# SKELETON AND MUSCLES,

AS REPRESENTED IN THE

# TABLES

# TABLE I.

THIS figure represents a front view of the human skeleton, with some of the ligaments and cartilages, which connect the bones to each other.

#### HEAD and NECK.

a, Os frontis.

ide

ng sti-

da een owthe ore, ney

the ted

Alore

re-

ave

rs. bet

he

en-

cal

ind

S.

b, Os parietale.

Between a and b, part of the coronal future.

c, The pars squamosa of the temporal bone. Between b and c, the squamous suture.

Below the pars squamosa, the zygoma; and, lower down, above f, the mastoid process.

Between the pars fquamofa and the cavity, which contains the eye-ball, called *orbit*, the temporal process of the sphenoid bone is seen.

Aa4

d, Os

T

f,

b

a,

b,

c,

d

f

g:

d, Os malæ.

Above d, a portion of the transverse suture.

e, Os maxillare fuperius, with the eight teeth of the

right fide.

The nasal process of the superior maxillary bone has theos nasi, joined by the lateral nasal suture, to its inside; and at the outside, within the orbit, the os unguis.

The offa nasi joined to each other before, by the

anterior nafal future.

f, Os maxillare inferius with fixteen teeth; the four anterior named incifores, the two corner ones canini, and the five posterior on each side molares.

Opposite to f, the angle of the lower jaw; above f, the condyloid process, by which the jaw is connected to the temporal bone, at the root of the zygoma; and behind the os malæ, the coronoid process.

g, The feven cervical vertebræ, with their intermediate

cartilages.

Opposite to g, their transverse processes.

#### TRUNK.

a, Sternum.

a, Its middle piece, to which one half of the cartilage that connects the fecond rib, the whole of the cartilages of the third, fourth, fifth, fixth, and one half of the feventh, are fixed.

Above a, the first or upper triangular piece, to which the clavicle and one half of the cartilage

that connects the fecond rib are fixed.

Below a, the extremity, or third piece of the sternum, named cartilago ensiformis, to which one half of the cartilage that connects the seventh rib is sixed.

b, The feventh, or last true rib.

c, The twelfth, or last of the five false ribs.

d, The

d, The five lumbar vertebræ, with their intermediate cartilages.

Opposite to d, their transverse processes.

e, The os facrum, with its five divisions.

f, Os innominatum, divided into

g, Os ilium,

le

IS

S

e

le

7-

f,

1-

le

d

e

1-

of

d

e

e e

b

e

h, Os pubis,

i, Os ischium.

Opposite to i, the foramen thyroideum.

#### SUPERIOR EXTREMITY.

a, The clavicle fixed before, to the first piece of the sternum, and outwards to the acromion of the scapula.

b, The fcapula.

Above b, the cervix of the scapula.

Opposite to it, the inferior costa; and below the outward extremity of the clavicle, the superior costa, and coracoid process, are seen.

c, The os humeri.

The upper end of it, which is connected to the cavity of the scapula, named glenoid, below the acromion, is named its head or ball; on each side of which is seen a tubercle, named external and internal; and between these, a groove for lodging the long head of the biceps slexor cubiti.

d, The internal condyle.

e, The external condyle.

Between d and e, the trochlea, upon which the ulna moves.

f, The radius.

The upper end, which moves on the external condyle of the os humeri, is named its *head*; below that, the tubercle for the infertion of the biceps flexor cubiti, and between these the cervix.

The inferior end of it is connected to the carpus.

g, Ulna.

The

The upper end of it forms the coronoid process, for the insertion of the brachialis muscle.

The inferior end has a process named flyloid, which is connected to the carpus by a ligament.

b, The carpus, formed of eight bones.

i, Metacarpal bone of the thumb.

k, The metacarpal bones of the four fingers.

I, The two joints of the thumb.

m, The three joints or phalanges of the fore-finger; and the same are seen in each of the other three.

#### INFERIOR EXTREMITY.

a, Os femoris.

The upper end of it is named its *head* or *ball*, which is lodged in a deep tocket of the os innominatum, named *acetabulum*.

Between the head and trochanter major, the cervix.

- b, Trochanter major.
- c, Trochanter minor.
- d, Internal condyle.
- e, External condyle.

f, Patella.

The place where it moves upon the os femoris, is named trochlea.

g, Tibia.

Between the tibia and the condyles of the os femories the femilunar cartilages are feen; and below the joint, the tubercle of the tibia-

n, Fibula.

- i, Malleolus internus.
- k, Malleolus externus.

1, Os calcis.

Between land m, the other fix bones of the tarfus.

m, Metatarfal bones of the four toes.


n, The three joints, or phalanges, of the four toes-

o, Metatarfal bone of the great toe.

p, The two joints of the great toe.

TA

TAB. I.



E

h ı,

is

0-)W

The Donaldon delinit et soulet

T

w T

a

в

C

-

-

# TABLE II.

THIS Table represents the first Layer of Muscles situated on the anterior part of the whole Body, immediately under the common Teguments, and tendinous Fasciæ.

Muscles situated on the HEAD and NECK.

a, The anterior fleshy belly of the occipito-frontalis fituated on the os frontis-

Above a, the tendinous aponeurofis of the occipitofrontalis, covering the upper part of the parietal bones.

b, Attollens aurem.

Under it, the tendinous aponeurosis covering the temporal muscle.

Anterior auris between c and the ear.

c, Orbicularis palbebrarum.

Its tendon is seen at the inner canthus, fixed to the nasal process of the superior maxillary bone.

Levator labii superioris alæque nasi-

Seen divided into two portions running down along the fide of the nofe; and on the outfide of it, the levator anguli oris:

Next this, the

Zygomaticus minor.

Further outwards,

Zygomaticus major.

On the ala and tip of the nofe, the

Compressor naris.

d, Depressor anguli oris-

And

And beneath it, a portion of the depressor labii inferioris.

e, Orbicularis oris.

f, Platysma-myoides.

Behind f, the sterno-cleido-mastoidæus is seen thro' the platysma-myoides-

#### TRUNK.

a, Pectoralis major.

The upper part of it is covered by the origin of the platysma-myoides-

b, Serratus magnus.

The other portions refemble this.

c, Latissimus dorsi.

d, Obliquus externus descendens.

e, Linea semilunaris.

f, Linea alba.

Below f, umbilicus.

Between e and f, the rectus abdominis; and, at the inferior part of the linea alba, opposite to g, the pyramidales appear through the tendons of the oblique muscles.

g, Ring of the external oblique muscle; with the spermatic chord, passing through it, and covered by the

cremaster muscle.

## SUPERIOR EXTREMITY.

a, Deltoides.

Above the clavicle, a portion of the trapezius is seen.

b, Biceps flexor cubiti-

At the bending of the arm are seen its tendon going towards the radius, and the part, from which the tendinous aponeurosis that covers the fore-arm, cut off.

On the infide of the biceps, part of the triceps extenfor cubiti; and, on the outfide, part of the brachialis internus.

c, Supinator radii longus.


d, Pro-

I. ii e 0 0 0

•

.

TAB. 11



The Donaldson deline a

- d, Pronator teres.
- e, Palmaris longus.

f, Palmaris brevis.

On the palm of the hand, the aponeurosis palmaris is seen extended from the annular ligament at the wrist, to the roots of the metacarpal bones of the four fingers.

g, Flexor carpi radialis.

h, Part of the flexor sublimis perforatus.

i, Infertion of the flexor carpi ulnaris.

k, Abductor pollicis.

#### INFERIOR EXTREMITY.

a, Tensor vaginæ semoris, the vagina or tendinous fascia being cut off,

On the outfide of it, a portion of the glutæus maximus.

b, Part of the iliacus internus.

On the infide of it, between b and c, part of the ploas magnus.

c, Pectinalis.

d, Triceps longus.

e, Gracilis,

f, Sartorius.

g, Rectus cruris.

Its tendon is feen inferted into the patella, from which a strong tendon is fent to be fixed to the tubercle of the tibia.

h, Vastus externus.

i, Vastus internus.

k, Tibialis anticus.

l, Peronæus longus.

On the outfide of it, a portion of the folæus.

m, Extensor longus digitorum pedis, with the peronæus tertius, and extensor proprius pollicis pedis.

n, Gastrocnemius externus, seu Gemellus.

o, Solæus

Ta

b,

c,

f, g,

a,

b,

1,

d

f

p, Ligamentum tarfi annulare.

q, Abductor pollicis pedis.

# TABLE III.

THIS Table represents the Second Layer of Muscles on the anterior Part of the whole Body.

Muscles situated on the HEAD and NECK.

a, Corrugator fupercilii.

b, Temporalis.

c, Masseter.

d, Levator anguli oris.

e, Buccinator.

f, Orbicularis oris.

Opposite to the right ala nasi, the portion of this muscle which Albinus names

Nafalis labii fuperioris.

g, Depressor labii inferioris.

b, Sterno-cleido-mastoidæus, which is

Seen below, arifing from the sternum and clavicle, by two heads.

i, Sterno-hyoidæus.

On the outfide of it, the

Omo-hyoidæus.

Further out, a portion of the

Hyo-thyroidæus.

k, Levator scapulæ.

TRUNK.

a, Subclavius.

b, Pecto-

b, Pectoralis minor.

c, Serratus magnus.

d, Rectus abdominis, divided into several sleshy portions by its tendinous intersections.

e, Pyramidalis.

111.

of

ole

118

f, Obliquus ascendens internus.

g, Spermatic cord, with the origin of the cremaster muscle.

#### SUPERIOR EXTREMITY.

a, Biceps flexor cubiti.

b, Short head of the biceps.

Beneath the upper part of it, a portion of the coracobrachialis.

Beneath the under part, a portion of the brachialis internus.

c, Long head of the biceps.

At the bending of the arm, the tendon of the biceps, and the place where the tendinous aponeurofis was cut from it, are feen.

d, Extenfor carpi radialis longior.

Beneath it a portion of the

Extensor carpi radialis brevior.

e, Flexor sublimis perforatus.

f, Insertion of the extensor carpi ulnaris.

g, Extensors of the thumb.

b, Opponens pollicis.

On the infide of it, a portion of the

Flexor pollicis brevis.

i, Tendon of the flexor longus pollicis manus, after passing through the flexor brevis pollicis manus.

k, Abductor minimi digiti manus.

1, Flexor parvus minimi digiti manus.

m, Ligamentum carpi annulare.

TAB.

#### INFERIOR EXTREMITY

- a, Iliacus internus. Between a and b, part of the ploas magnus.
- b, Pectinalis.
- c, Triceps longus.

d, Gracilis.

e, Rectus cruris cut off near its origin.

- f, Tendon of the rectus cruris cut off above the patel. la, from which a strong tendon is sent to be inserted into a tubercle of the tibia.
- g, Portion of the glutæus medius. On the infide of it, part of the glutæus minimus.
- h, Vastus internus.
  i, Vastus externus.

k. Cruræus.

- 1, Infertion of the biceps flexor cruris into the fibula.
- m, Tendons of the gracilis and femitendinofus inferted into the tibia.
- n, Solæus.

o, Peronæus longus.

- p, Extensor longus digitorum, with the peronæus tertius on the outfide, and extenfor pollicis proprius on the infide.
- q, Solæus.

r, Flexor longus digitorum.

- f, Tendons of the tibialis posticus and flexor longus digitorum pedis.
- t, Flexor brevis digitorum pedis-

TAB. III

Thos Donaldson deline at rought

111.


eled

ed

eron

i-

E



# TABLE IV.

# Fig. 1.

THIS Figure represents the right Eye-ball, the fixMuscles which move it, taken out of the Orbit, with the Optic Nerve.

a, The eye-ball.

b, Optic nerve.

c, Musculus trochlearis, seu obliquus superior.

d, The trochlea, or pulley, with a bit of the os frontis, through which the tendon of the muscle passes towards the eye-ball.

e, Obliquus inferior, with a bit of the superior maxil-

lary bone, from whence it arises.

f, Levator oculi.

g, Depressor oculi.

h, Adductor oculi.

i, Abductor oculi.

# Fig. 2.

Represents the anterior Part of the Cartilage of the right Ear, with its proper Muscles.

a, Helicis major.

b, Helicis minor.

c, Tragicus.

d, Antitragicus.

Transversus auris, vid. Tab. viii. Fig. 3.

# Fig. 3.

Represents the third Layer of Muscles, with fome of the Ligaments, Cartilages, and naked Bones on the anterior Part of the whole Body.

a, Depreffor labii superioris alæque nasi.

b, Orbicularis oris, after most of the muscles, which are fixed to it, and assist to form it, have been taken away.

c, Buccinator.

Above c, part of the pterygoidæus externus is seen passing behind the coronoid process of the lower jaw.

d, Levator labii inferioris.

e, Sterno-thyroidæus.

Immediately above, and seemingly the continuation of it, the

Hyo-thyroidæus.

f, Scalenus medius.

Contiguous to it, on the infide, the

Scalenus anticus.

Above it, a portion of the

Trachelo-mastoidæus.

Between the scalenus anticus, and sterno-thyroideus, and hyo-thyroideus, the

Rectus capitis anterius major, and Longus colli.

## TRUNK.

a, Third row of external intercostal muscles.

The rest appear in the same manner between the other ribs.

b, Third row of internal intercostal muscles.

The rest appear between the other ribs.

c, Transversalis abdominis.

d, The

d, The place from which the inferior part of the tendon of the transversalis, that passes hefore the rectus and pyramidalis muscles, is cut off.

Between these portions of each side, the peritonæum is laid bare, and the ligaments of the bladder which were formerly the umbilical arteries and urachus.

Between this portion and os pubis, the spermatic cord is seen cut.

e, The inferior edge of the upper part of the tendon of the transversalis, which passes behind the rectus, and immediately adheres to the peritonæum.

f, The anterior lamella of the internal oblique, which joined the tendon of the external to pass over the rectus.

Between f and g, the posterior lamella of the internal oblique, joining with the tendon of the transversalis, to pass behind the rectus.

g, The place at the linea alba, from which the tendon of the external oblique, and anterior lamella of the internal, were cut off.

At g, Umbilicus.

## Superior Extremity.

- a, Subscapularis.
- b, Teres minor.
- c, Coraco-brachialis:
  - The part from which the short head of the biceps slexor cubiti was cut off from it, is seen at its upper end.
- d, Brachialis internus.
- e, Brachialis externus, or third head of the triceps.
- f, Extensor carpi radialis longior, and with it the extensor carpi radialis brevior.

Both these are distinctly seen in the right hand.

Between the tendon of the brachialis internus and extensor radialis, the

Supinator radii brevis is feen.

B b 2

g, Flexor

- g, Flexor longus pollicis manus, with the fleshy portion of it which arises from the internal condyle of the os humeri.
  - b, Flexor profundus perforans, which splits into four tendons, which pass under the ligamentum carpi annulare.

i, Pronator quadratus.

k, Adductor minimi digiti manus.

1, One of the lumbricales.

The other three appear in the fame manner, along the tendons of the flexor profundus.

Behind these, the internal interossei are seen.

#### INFERIOR EXTREMITY.

- a, Glutæus minimus.
- b, Iliacus internus.
  On the infide of it, between b and c, the ploas magnus.
- c. Obturator externus.
- d, Adductor brevis femoris.
- e, Adductor magnus femoris.

f, Gracilis; which is

Seen inferted into the infide of the head of the

g, The short head of the biceps flexor cruris.

h, Peronæus longus.


i, Peronæus brevis.


Between these two peronæi and tibia, the tibialis

posticus is seen.

k, Tendon of the tibialis posticus, covering the tendon of the flexor longus digitorum pedis.

1, Extensor brevis digitorum pedis.





The Donaldson delint at

## Fig. 4.

Represents the Levatores Ani cut off from the Bones before, with their Connection to the Extremity of the Rectum, and Bulb of the Urethra.

a, The urethra, and its corpus cavernosum cut off.

b, Bulb of the urethra.

c, The circular fibres which furround the verge of the anus; by some named fphincler internus.

d, The anterior edge of the levator ani, cut off from the os pubis, and fide of the pelvis.

## Fig. 5.

Represents the anterior Part of the Sphincter Ani, and Bulb of the Urethra.

a, The urethra, and its corpus cavernofum cut.

b, Bulb of the urethra.

c, Left half of the sphincer ani, running obliquely upwards, to join with the right half.

d, The acute portion, which is inferted into the perinaum.

## Fig. 6.

Represents the Corpora Cavernosa of the Penis, Corpus Cavernosum Urethræ, Acceleratores Urinæ, Transversales, and Erectores Penis.

a, Corpus cavernosum penis cut.

b, Urethra, and its corpus cavernofum cut.

 $B_3$ 

c, Erector

- c, Erector penis covering the crus.
- d, Transversalis penis.
- e, Accelerator urinæ.
- f, Transversalis penis alter, running along the inside of the left erector penis.

### TABLE V.

REPRESENTS a Back View of the Human Skeleton, with some of the Ligaments and Cattilages which connect the Bones.


#### HEAD and NECK.

- a, Os parietale, joined to its fellow by the fagittal future.
- b, The os occipitis, joined to the parietal bones by the lambdoid future, which is between a and b.
- c, Os malæ.
- d, Maxilla inferior, with a view of the teeth of both jaws from behind.
- e, The feven cervical vertebræ.


#### TRUNK.

- a, The feventh, or last true rib.
- b, The twelfth, or last rib.
- c, The five lumbar vertebræ.
- d, Os facrum.
- e, Os coccygis.
- f, Os innominatum, divided into
- g, Os ilium.
- b, Os pubis.
- i, Os ischium.

AB. V.



Thos Donaldson delint at souls



### SUPERIOR EXTREMITY.

- a, The clavicle, joined outwards to the acromion of the scapula.
- b, The scapula.
- c, Os humeri.
- d, Internal condyle.
- e, External condyle-
- f, Radius.
- g, Ulna, its upper end, named olecranon; and near the wrist, its styloid process.
- b, The eight bones of the carpus.
- i, The metatarfal bone of the thumb.
- k, The metatarfal bones of the four fingers.
- 1, The two joints of the thumb.
- m, The three joints or phalanges of the four fingers.

### INFERIOR EXTREMITY.

- a, Os femoris.
- b, Trochanter major, and at the infide of it the cervix.
- c, Trochanter minor.
- d, Internal condyle.
- e, External condyle.
- f, Tibia.
- g, Fibula.
- h, Malleolus internus.
- i, Malleolus externus.
- k, The feven bones of the tarfus.
- l, The metatarfus.
- m, The joints or phalanges of the toes.

TA

## TABLE VI.

REPRESENTS a Back View of the Muscles which are immediately situated below the common Integuments.

### HEAD and NECK.

- a, Part of the occipito-frontalis muscle, with its aponeurosis.
- b, Attollens aurem.
- c, Anterior auris.
- d, Retrahentes auris.

#### TRUNK.

a, Trapezius, seu eucularis.

- b, Its tendinous joining with its fellow in the nape of the neck, which is called *ligamentum nuchæ* feu colli.
- c, The fleshy belly of the latissimus dorsi.
- d, The tendon of the latissimus dorsi, which arises in common with the ferratus posticus inferior.
- e, Part of the obliquus externus abdominis.

### SUPERIOR EXTREMITY.

a, Deltoides.

b, Infraspinatus, with a portion of the teres minor and major below it.

Its tendon is feen inferted into the head of the ulna,

TAB. VI

The Donaldson delint et sculpt.

a

e

1 8 1

called olecranon; and, on the infide of it, the an-

d, Extensor carpi radialis longior, covered by a portion of the supinator radii longus; and, under it, a portion of the extensor carpi radialis brevior.

- e, Extensor digitorum communis manus, which splits into sour tendons, and pass, with the indicator, under the ligamentum carpi annulare externum, at the extremities of the metacarpal bones, under ligaments proper to themselves; and are lost in a broad tendon, which covers the back of the four fingers.
- f, Extensor ossis metacarpi pollicis manus.
- g, Extensor primi internodii pollicis manus.
- b, Extensor secundi internodii pollicis manus.
- i, Extenfor carpi ulnaris.
- k, Part of the flexor carpi ulnaris.
  - Under it, part of the
  - Flexor profundus perforatus.
    - And on the infide, part of the
  - Flexor sublimus persoratus, which are more distinctly seen on the right fore-arm. Likewise, on the right hand, are seen part of the abductor pollicis manus, abductor minimi digiti manus, and the aponeurosis palmaris.

## INFERIOR EXTREMITY.

- a, Glutæus maximus.
- b, Part of the glutæus medius.
- c, Part of the tenfor vaginæ femoris.
- d, Vastus externus.
- e, The long head of the biceps flexor cruris:
  And beneath it,
- f, Part of the short head.
- g, Semitendinosus:
  - And beneath it, on each fide,
  - A portion of the semimembranosus is seen.
- b, Gracilis.

Tab.

a, R

b, R

c, T

d, S

e, P f, P

g, I

*b*,

c, d,

f

g

A

On the outfide of it,

A portion of the adductor magnus is feen.

i, A small part of the vastus internus.

k, Gastrocnemius externus, seu gemellus:

And within its outer head, A portion of the plantaris.

1, Solæus seu gastrocnemius internus.

m, Tendo Achillis, with the plantaris.
n, Peronæus longus.

o, Peronæus brevis, between it and the tendo Achillis, a portion of the flexor longus digitorum pedis.

p, Tendons of the extensor longus digitorum pedis, with the peronæus tertius passing under the ligamentum tarsi annulare; and the slexor brevis digitorum pedis is seen beneath them.

7, Abductor minimi digiti pedis; and above it the tendons of the peronæus longus and brevis, passing

under proper ligaments of their own.

## T A B L E VII.

REPRESENTS the fecond Layer of the Muscles on the Back part of the Body.

### HEAD and NECK.

a, Temporalis; its tendon is feen passing below the zygoma.

b, Masseter.

c, Splenius capitis et colli-

d, Portion of the complexus.

e, Levator scapulæ, seu musculus patientiæ.

TRUNK.

### TRUNK.

a. Rhomboides major.

III.

is,

is,

a-

li-

n-

g

b. Rhomboides minor:

And immediately above it, the upper edge of the ferratus posticus superior is seen.

t, The ferratus posticus superior on the right side.

d, Serratus posticus inferior.

e, Part of the spinalis dorsi.

f, Part of the longissimus dorsi-

g, Part of the facro-lumbalis.

b, Serratus magnus.

i, The broad tendon, by which the latissimus dorsi begins, and from which the tendon of the serratus posticus inferior is inseparable.

k, Part of the obliquus internus ascendens abdominis.

l, The sphincter ani, fixed to the point of the os coccygis; at the side of which the coccygæus, and a portion of the levator ani, are seen; and lower down, opposite to l, part of the transversalis penis.

### SUPERIOR EXTREMITY.

a, Supra-spinatus.

b, Infra-spinatus.

c, Teres minor.

d, Teres major.

e, Triceps extensor cubiti.

f, Its head, called longus.

g, The brevis: And,

- b, A small portion of the third head, named brachialis externus.
- i The tendon of the triceps, inferted into the olecra-

k, Part of the brachialis internus.

- l, Anconæus, which seems to be continued from that part of the brachialis externus immediately above it.
- m, Extensor carpi radialis longior; and beneath it, the brevior:

brevior: both are seen at the wrist, inserted into the metacarpal bones of the fore and middle fingers.

n. Flexor carpi ulnaris.

o, Part of the supinator radii brevis.

p, Extensor offis metacarpi pollicis manus. q, Extenfor primi internodii pollicis manus. r, Extensor secundi internodii pollicis manus.

f, Indicator, inferted into the root of the first joint of

the fore-finger.

t, One of the three external interoffei manus. other two are distinctly seen without letters.

u, One of the tendons of the extensors of the fingers cut; and the same is seen in each of the other three fingers, joining with the tendons and aponeuroses of the interossei and lumbricales, and fpread upon the back of the fingers.

N. B. On the right hand, part of the flexors of the fingers, the abductor pollicis, and minimi digiti,

are feen.

### INFERIOR EXTREMITY.

a, Glutæus medius.

b, Pyriformis.

c, The two muscles called gemini, between which the tendon of the obturator internus passes over the tuberosity of the os ischium; and its sleshy belly is feen within the pelvis, partly covered by the coccygæus and levator ani.

d, Quadratus femoris.

e. Vastus externus.

f, f, Parts of the triceps magnus.

g, Long head of the triceps flexor cruris, and beneath it part of the short head is seen.

b, Semitendinofus, and beneath it parts of the femimembranofus are feen on each fide of it.

i. Gracilis.

k, A small portion of the vastus internus.

B. VII

The Donaldson beling it sculpt

II.

to

of

rs er d

e i,

2

Tab.

l, P

n, S

p, 7 q, 1

7,

ſ, I

Fof

a, b, 1, Poplitæus.

m, The fleshy belly of the plantaris; and its long flender tendon is seen passing over the inside of the solæus.

n, Solæus.

off; but the flesh of the solwus runs further down.

t, Tendo Achillis, with the plantaris.

q, Peronæus longus, passing at the outer ankle to the sole of the foot; beneath it, the peronæus brevis to the root of the metatarsal bone of the little toe; and between it and the tendo Achillis, a portion of the slexor longus digitorum pedis.

r, Tendons of the extensor longus digitorum pedis, with the peronæus tertius; and beneath these, the

f, Flexor brevis minimi digiti pedis.

## TABLE VIII.

Fig. 1.

REPRESENTS the third Layer of Muscles on the posterior Part of the Body, with some of the Ligaments and naked Bones.

Muscles on the HEAD and NECK.

a, Part of the buccinator.

b, Complexus.

c, Trachelo-mastoidæus; on the outside of it, the transversalis colli.

b,

d

f

ghi, kl,

77

n

I

0

- d, Scalenus medius.
- e, Scalenus posticus.

#### TRUNK.

a, Spinalis dorfi; and beneath it, the multifidus spina,

b, Longissimus dorsi, which sends off a sleshy slip to the trachelo-mastoidæus.

- c, Sacro-lumbalis, with the cervicalis descendens sent off from it along the side of the neck, and outside of the transversalis colli.
- d, Semispinalis dorsi-

e, Transversalis abdominis.

N. B. The spaces between the spinous processes of the vertebræ have muscular fasciculi between them, particularly those of the neck; and are named interspinales colli, dorsi, and lumborum; but those of the back seem to be tendinous and ligamentous.

## SUPERIOR EXTREMITY.

a, Teres major.

b, Part of the coraco-brachialis.

c, Part of the brachialis internus.

- d, The third head of the triceps extensor cubiti, called brachialis externus, after the longus and brevis have been cut off.
- e, Extenfor radialis longior.
- f, Extensor radialis brevior.
- g, Part of the flexor profundus perforans.

b, Supinator radii brevis.

i, Part of the adductor pollicis manus.

k, One of the three external interoffei; the other two may be easily distinguished without letters.

I, Tendons of the extensors of the fingers, joining with those of the lumbricales and interossei, which form a tendinous expansion on the back of the four fingers.

N. B. On the right hand, part of the flexors of the fingers

fingers and thumb, part of the adductor pollicis, and the whole of the adductor minimi digiti, are feen.

#### INFERIOR EXTREMITY.

a. Glutæus minimus.

b, Obturator internus; its fleshy belly is seen within the pelvis.

Beneath b, the tendon of the obturator externus.

c, Semimembranofus.

d, The short head of the biceps flexor cruris.

e, Triceps magnus.

f, Gracilis.

t III

inæ.

the

fent

tfide

s of veen

are but

iga-

lled

evis

wo

vith

rm

our

the ers In the ham, the origins of the two heads of the gaftrocnemius externus and plantaris are feen.

g, Poplitæus.

b, Tibialis posticus.

i, Flexor longus digitorum pedis.

k, Flexor pollicis longus.

I, Peronæus longus, running down to be inferted into the metatarfal bone of the little toe.

Beneath it, the peronæus brevis, passing to the sole of the foot.

m, Extensor brevis digitorum pedis-

n, Part of the flexor longus digitorum pedis-

## Fig. 2.

Represents the fourth Layer of Muscles on the posterior Part of the Neck.

a, Rectus capitis posticus major.

b, Rectus capitis posticus minor.

c, Obliquus capitis superior.

d, Obliquus capitis inferior:

e, Scalenus medius.

f, Part

T

R

a

f, Part of the multifidus spinæ, covered by the semispinalis colli-

Between the spinous processes of the vertebræ, the interspinales colli are seen double; because these processes are bifurcated.

Between the transverse processes, the posterior row

of the intertransversales colli is seen.

## Fig. 3.

Represents a Portion of the posterior Part of the Cartilage of the Ear, to show,

a, Transversus auris,
Situated on the parts opposite to the antihelix and
scapha.

## Fig. 4.

Represents an Outline of the Base of the Os Occipitis, and inside of the Lower Jaw; with a View of the Muscles which surround the back Part of the Larynx and Pharynx.

- a, The upper part of the membrane of the pharynx.
- b, The trachea, cut.

c, The œfophagus, cut.

- d, The inner transverse fibres of the cosphagus, laid bare.
- e, The outer fibres descending obliquely backwards.
- f, Constrictor inferior pharyngis.
- g, Constrictor medius pharyngis.
- b, The cornu of the os hyoides.
- i, Constrictor superior pharyngis.
  k, The part of it which joins with the buccinator.

1, Stylo-pharyngæus.

Fig.

# Fig. 5.

Represents an Outline of the Inside of the Os Pubis, Os Ischium, and Back of the Os Coccygis, after the Os Sacrum and Ligaments have been taken away; with a posterior View of the Levatores Ani, and Extremity of the Rectum, resembling the Shape of a Funnel.

a, The anterior portion of the levator ani, viewed on its infide within the pelvis, arifing from the os pubis and upper part of the foramen thyroidæum.

b, Its origin from the spinous process of the os ischium.

c, The posterior part seen on its outer side.

d, Its infertion into the os coccygis, below which the fleshy fibres are continued with its fellow.

7, The circular fleshy fibres surrounding the extremity of the rectum, which authors name sphincter internus ani.

m, The anus.

tIII.

ispi-

the

nese

WO

the

ind

C-

th he

id

# Fig. 6.

Represents an Outline of the Back of the Os Coccygis, and posterior Part of the Sphincter Ani.

a, The infertion of the sphincter ani into the extremity of the os coccygis; to which, from

b, The anus,

The fibres meet from each fide, in angles; which are more acute as they point upwards.

## Fig. 7.

Represents the first Layer of Muscles on the Sole of the Foot, after the Aponeurosis Plantaris has been taken away.

a, Abductor pollicis pedis.

bb, Abductor minimi digiti pedis.

c, Flexor brevis digitorum pedis, which splits into four tendons, and are perforated by the tendons of the flexor longus digitorum pedis.

Between these tendons the lumbricales are seen.

d, Tendon of the flexor pollicis longus:
And beneath it, the flexor pollicis brevis.

e, Transversalis pedis.

## Fig. 8.

Represents the second Layer of Muscles on the Sole of the Foot, after the Abductor Pollicis Pedis, and Abductor Minimi Digiti Pedis, and the Flexor Brevis Digitorum Pedis, have been taken away.

a, Tendon of the flexor longus digitorum pedis.

bb, Flexor accessorius, seu massa carnea Jacobi Sylvii.

ecce, The four lumbricales arising from the tendons of

the flexor longus digitorum.

dd, Tendon of the flexor pollicis longus; and, in its progress, joins the tendon of the flexor longus digitorum pedis.

e, Tendon of the tibialis anticus.

f, Tendon of the peronæus longus.

g, Tendon of the peronæus brevis.

b Flexor



Tal

b,

k, ]

m,

0

b, Flexor pollicis brevis, with a portion of the abductor pollicis, on the infide of the great toe.

On the outfide of i, is the adductor pollicis.

k, Flexor brevis minimi digiti pedis.

l, A ligament which supports the bones of the tar-

Cc 2

Before I, two of the interossei are seen.

m, Transversalis pedis.

A

r terror las anv

The car position fraction and a sequence of a solution of the second of the care of the ca

en en la language de la companya de la language de Language de la 
A

ten cor ove is l eac

mic ing but fuit bor

thi

ATE ON THE SHIP OF THE STATE OF THE

effette tracks from the control according

## S Y S T E M OF A N A T O M Y.

## PART IV.

CONTAINING

An Enumeration of the Cartilages, Ligaments, &c. of the Joints.

#### OF THE HEAD.

THE condyloid processes of the os occipitis, the glenoid cavities or articular fossulæ of the ossa temporum, the eminences next these cavities, and the condyloid processes of the lower jaw, are all covered over with very white and smooth cartilages; and there is likewise an interarticular or moveable cartilage in each articulation of the lower jaw with the temporal bones.

This cartilage is thick near the circumference; very thin and transparent, and sometimes perforated in the middle. The lower side is uniformly concave, answering to the oblong convexity of the maxillary condyle; but the upper side is partly concave, and partly convex, suited to the fossula and eminence of the temporal bone.

The remaining cartilages of the bones of the head, viz.

viz. the cartilaginous feptum, and other cartilages of the nose, the small cartilaginous pulley in each orbit, the cartilages of the outward ear, and those which are joined to the os hyoides, must be referred to the description of the viscera.

The ligaments of the bones of the head are, Those of the articulation of the lower jaw with the temporal bones; those between the occipital bone and vertebra of the neck; those by which the os hyoides is con-

nected to the styloid process.

The ligaments of the articulation of the lower jaw are very firm and strong; and are fixed by one extremity round the glenoid cavity or articular fosfula and eminence of each temporal bone, by their middle round the interarticular cartilage, and by the other extremity round each condyle of the lower jaw, in such a manner as to allow the intermediate cartilage to sollow the motions of the condyles, and to change their situation from the glenoid cavities to the tubercles of the zygomatic processes, and to return again, as was mentioned in the description of the bones.—Besides the ligaments of the articulation of this bone, another small ligament is extended between the inner side of each angle of the jaw and the styloid process of the temporal bone.

The ligaments of the occipital condyles and superior oblique processes of the first vertebra of the neck, are—The circular ligament which arises from the inferior margin of the foramen magnum, to be fixed to the superior margin of the first vertebra:—The capsular ligaments which arise from the edges of the condyloid processes of the os occipitis, and are fixed to the edges of the superior oblique processes of the first vertebra:—The two lateral, or moderator ligaments, which are fixed by one end to the inner side of the first vertebra and to the os occipitis, and by the other end to the side of the processus dentatus of the second vertebra:—The perpendicular ligament, which arises from the point of

IV.

of bit,

are

de-

e of

oral bræ

on-

jaw

tre-

and

exuch

fol-

neir s of

was

des

her

e of

em-

rior

e—

fu-

ular

oid

ges

ra:

are

bra

fide

The

t of

the

the processus dentatus, and is firmly inserted to the anterior edge of the foramen magnum occipitis:

The ligamentum nuchæ, which arises from the protuberance on the back part of the occipital bone, and descends over the spinous processes of the vertebræ of the neck, to which it strongly adheres.

The bones of the head, as well as all the other bones of the body, are covered by a particular membrane; of which that part which belongs to the skull is termed pericranium; and that which covers the bones of the face, or of the two jaws, is called simply periosteum.

The fynovial fubstances of the occipital and maxillary articulations, have nothing particular to them.— They are proportioned to the joints to which they belong, and lie between the capsular ligaments and circumference of the cartilages.

#### OF THE TRUNK.

THE cartilages of the vertebræ in general are of two kinds; one proper to each vertebra, the other common to the two vertebræ that lie next each other. The first may be termed cartilages of articulation; the others,

cartilages of symphysis.

The proper articular cartilages of each vertebra of the spine, are those four which cover the surfaces of the oblique processes. In the natural state they are very white and smooth, and much thicker than in dry bones. Their circumference is the same with that of the articulated sides of the processes, except in those places where there are small superficial notches. In the first vertebra of the neck and vertebræ of the loins these cartilages are thicker than in the rest.

The cartilages of the two inferior oblique processes of the first vertebra, and of the two superior oblique of the second, seem to be disproportionate, though not so much as in dry boncs; and in some subjects we find moveable or interarticular cartilages between the pro-

cesses of these two vertebræ.

C 4

The

The first vertebra of the neck has a cartilaginous covering on the back part of the anterior arch, corresponding with another on the fore-side of the processus dentatus of the next vertebra; so that each of these two vertebræ has sive articular cartilages, besides the interarticular ones already mentioned.

The vertebræ of the back, besides the four cartilages of their oblique processes, have others which do not belong to their articulations with each other, viz. those that cover the lateral fossulæ in the bodies of these vertebræ, and the fossulæ of their transverse processes, by both of which they are articulated with the ribs.

The cartilages of fymphysis lie between the bodies of the vertebræ, uniting them closely together; the breadth and circumference of them answering exactly to that of the surfaces to which they are connected; but their height or thickness is different in each class of the vertebræ. Between the vertebræ of the loins, they are a quarter or third part of an inch in thickness, according to the fize of the subject: In those of the neck, they are not so thick; and the thinnest of all are between the vertebræ of the back.

These cartilages are not of an equal thickness in all their parts. Those on the neck and loins appear to be thickest on the foreside, and those of the back rather thickest on the back-side; but these differences are most remarkable in the vertebræ that lie near the middle of each class.

The internal structure of these cartilages is different from that of all the other cartilages of the body, and indeed they resemble the rest in nothing but in whiteness and elasticity. When we view their circumferences only, they seem to be one uniform mass as the others generally are; but when they are divided by an incision parallel to that surface of the vertebræ to which they are joined, we see they are composed of many cartilaginous concentrical rings contained within each other, a small distance being lest between them. The

outer

0-

e-

18 *fe* 

le

es

ot

le

r-

y

of h

of

It

r

c

0

outer part of these cartilages is the most solid and hard; and they gradually become fofter, till at last they are almost in the form of a glairy liquor in the centre.

These rings do not form an entire circumference; being turned inwards on the back-fide, answering to the polterior cavity in the body of each vertebra. lie horizontally, between the vertebræ. The interstices of the rings are filled with a mucilaginous fubstance, less fluid than that of the joints. Each lamina taken separately is very pliable according to its length; but taken together, they are not fo eafily bent, partly because of their circular figure, and partly because of their vicinity and multiplicity. They yield, however, in the inflections of the spine; and their external surface, which is the ordinary situation of the spine, is even with the surface of the vertebræ, becomes prominent or juts out on that fide toward which the inflection is made, the cartilages being then compressed by the vertebræ.

They likewise yield on all sides, without any inflection of the spine, to the weight of the upper part of the body; but this is done by very fmall and imperceptible degrees, and most of all when the body is loaded with

any exterior weight.

They restore themselves afterwards merely by being freed from compression: So that a man is really taller after lying, than after he has walked or carried a burden for a length of time; the most natural and simple reason that can be given for the different height of the fame person at different times, first observed in England, and afterwards confirmed by Mr Morand, a member of the Royal Academy of Sciences, being the different state of the intervertebral cartilages. The intervertebral cartilages of the neck lying for the most part between the convex fide of one vertebra, and the concave fide of another, are of a greater extent in proportion to the fize of these vertebræ, than those of the back and loins. Without this convexity and hollowness in these vertebræ, which are the least of all, the carticartilages could not have been made large enough to be

able to refift strains and great exertions.

The os facrum has no cartilage, excepting that between its upper end and the last vertebra of the loins, and those by which it is connected with the offa innominata.

The intervertebral cartilages of this bone are for the most part entirely obliterated in the adult subject.

The cartilages which join the different portions of the os coccygis are preserved in some subjects to a very great age, in others they soon become entirely offisied.

The vertebræ are strongly connected together by three kinds of ligaments. Each vertebra is connected to that above and below it by a great number of very short and strong ligaments, which cross each other obliquely, and are fixed round the edges of the body of each vertebra. These crucial ligaments cover the circumference of the intervertebral cartilages, and adhere They feem to be loofer in the cerviclosely to them. cal and lumbar vertebræ, than in those of the back; and by that means yield to the cartilages in the different flections of the spine already mentioned. The bodies of all the vertebræ, from the first of the neck to the os facrum, are covered by a strong ligamentous band, which runs down chiefly over their anterior convex furface, and fends off many processes to be fixed to their fpongy fubstance.

All the vertebræ are likewise strongly connected by a ligamentous tube, which lines the inner surface of the medullary canal from the occipital hole to the os facrum, representing a kind of strong flexible funnel, its cavity at the upper part being equal to that of the occipital foramen, and ending in a small point at the os facrum. This ligament is composed of several strata of longitudinal and oblique sibres, interwoven together, adhering closely to the inside of the spinal canal through its whole length, by a great number of silaments detached from it to the porous substance of the vertebræ.

IV.

be

be-

ns,

10-

he

he

ry

y

d

y )-

of

-

e

The first vertebra is not only fixed to the os occipitis by a portion of this ligamentous funnel; but also by a distinct and very strong ligamentous covering, which furrounds and adheres very closely to that funnel. This covering is fixed, above, round the great occipital foramen, where it begins to adhere to the funnel; and below, quite round the circumference of the first vertebra.

The fecond vertebra has two ligaments peculiar to it; one which connects the processus dentatus to the os occipitis; and another transverse, which confines this process within the anterior portion of the cavity of the first vertebra. The first may be termed the perpendicular, and the fecond the transverse ligament of the processus dentatus.

The perpendicular ligament is very strong and thick, and adheres in a very fingular manner to the three planes of the apex of the process; and is afterwards divided into two or three portions which are fixed in a fimilar manner, in the anterior edge, of the foramen magnum, and in the inequalities of the cuneiform process near that hole.

The transverse ligament may be said more justly to belong to the first vertebra, both ends of it being inferted in the lateral impressions of the inner surface of that vertebra mentioned in the description of the skeleton; but it is ranked among the ligaments of the fecond vertebra, because of its use, and because of the in-

fertion of its middle portion.

This thick ligament is stretched from one side of the inner surface of the first vertebra to the other side, and prevents the processus dentatus from wounding the spinal marrow in the circumgyration of the head. About the middle of the fore-fide, its texture is very close where the processus dentatus plays upon it. Along the whole bony canal of the spine, between the basis of each spinal process, lies a flat and very elastic ligament, of a yellowish colour, which fills up the posterior great notches notches of the vertebræ, adhering to their edges; and likewise to the neighbouring portions of the funnel or

great ligamentous tube.

Between the extremities or apices of the spinal processes we find small ligamentous cords, which go from one spine to that next it; and which are really double, though they seem to be single in the vertebræ of the back and loins.

Between all the spinous processes, from their apices to the middle of their basis, lies a ligamentous membrane passing between each two processes; and thereby distinguishing the right side from the left. A ligament of the same kind is found between the transverse processes.

These are intermuscular ligaments or ligamentous septa, which divide the muscles on opposite sides, as will appear more particularly in describing the muscles.

The first kind may be termed interspinales, the other

intertransversales.

The articular ligaments of the spine are, Those which tie the superior oblique processes of the first vertebra to the condyles of the os occipitis, those that join the cartilaginous surface of the processus dentatus to the anterior cavity of the first vertebra, and those by which all the oblique processes are connected together.

These are small, short, strong ligamentous fasciculi, fixed by both extremities round the cartilaginous surfaces of the processes, surrounding very closely all the

capfular ligaments of these articulations.

The vertebral ligaments of the ribs, or those which connect the ribs to the bodies and transverse processes of the vertebræ of the back, are of the same kind, being inserted round the cartilaginous fossulæ in the body and processes of each vertebra.

Besides all these ligaments of the spine, there is one which goes in form of a membrane, from the os occipitis, all the way to the last two vertebræ of the neck.

fi

d

It is broad at the upper part, and from thence diminishes gradually. By its upper broad extremity, it is fixed along the occipital spine; and, by one edge, in the posterior tubercle of the first vertebra, between the two spinal surce of the following vertebræ, and in the apices of the spinal processes of the lowest vertebræ; but the other edge is loose. This is a true intermuscular ligament, and has the name of ligamentum cervicale posserius a nuchæ. There are two lateral ligaments of the same kind, fixed to the transverse processes of the vertebræ of the neck.

#### OF THE STERNUM AND RIBS.

THE sternum of an adult has commonly sixteen cartilages; fourteen of which are articular, the other two symphyses. Of the articular cartilages, two belong to the articulations of the clavicles; and twelve to those of the true ribs, from the second to the seventh inclusive. The two symphyses are those between the sternum and the sirst rib on each side.

There is likewise another symphysis, by which the upper portion of the sternum is connected to the lower; the cartilage of which is often obliterated in an advan-

ced age.

The cartilago ensiformis is often long toward the sternum, and more or less cartilaginous at the other end. In very aged persons it has been found entirely ossified, and sometimes wholly cartilaginous even in adults.

All the ribs have cartilaginous portions, which differ from each other in length, breadth, incurvation, adhefion, and in their extremities; all which were explained in the description of the skeleton. We have only to observe here, that these cartilages are whiter, more polished, broader, and thicker in the natural state than when they are dried. The cartilages of the false ribs are naturally more slender and pliable than those of the true ribs: the middle or inner substance acquires the

con-

confistence of bones in old age; and their extremities fometimes offify, and are immoveably fixed to the ster-

414

The sternum has several ligaments, by which it is connected with the clavicles and ribs. It is joined to the clavicles by strong short ligaments, fixed by one extremity round the edges of its two superior notches; by the other, to the extremity of each clavicle; and by the middle, to the interarticular cartilages, already explained; furrounding the particular ligaments which go between the edges of these cartilages and the sternum, and the capfular ligaments between them and the clavicles.

All the ribs are connected to the bodies of the vertebræ by strong short ligamentous fasculi, fixed by one end round the small pits in the vertebræ, and by the other round the head of each rib. The ribs are likewise tied to each other by their ligaments, which go obliquely from the cartilage of each rib to that of the The ten uppermost ribs on each side are connected to the transverse processes of the vertebræ of the back by ftrong short articular ligaments, fixed to the tuberofities of the ribs, and round the fosfulæ of the processes, much in the same manner with those that go between the heads of the ribs and bodies of the verte-Both these articulations are provided with capfular ligaments. The eleventh rib on each fide having no articulation with the transverse processes, is connected to them by auxiliary strong short ligaments fixed in its neck. The last rib is only joined by its head to the body of the twelfth vertebra of the back; but it is connected in a particular manner to the transverse process of the first vertebra of the loins by a broad ligament fixed to the whole upper edge of the process, and to the lower edge of the rib for about two thirds of its length.

The first true rib has no ligamentous connection with the sternum, the cartilaginous symphysis being

Part

fuffi

fma

not

of

but

nul

the

fro

rib

wi

the

the

th

th

th

ti

n

fufficient. The rest are closely joined to that bone by fmall ligamentous portions, fixed by one end round the extremity of the cartilage, and by the other round the notches in the sternum. On the upper and lower sides of each articulation, these ligaments are very short; but on the fore-fide, they are expanded over the ster-

num in a radiated manner.

The cartilage of the first false rib is joined to that of the last true rib by several short filaments, which go from the lower edge of the one to the upper edge of the other, near its small extremity. The other false ribs are connected together much in the same manner: with this difference only, that the filaments by which the fourth is connected to the third are longer than those above them; and those between the fifth rib and the fourth much longer than any of the rest; and for this reason, these two ribs are less steady than any of the rest.

The ligamentous expansions of the vertebræ are in place of a periosteum, both outwardly and inwardly. The sternum and bony portions of the ribs have a periofteum like the other bones. The cartilaginous portions of the ribs are covered by a membrane of the fame kind, termed perichondrium. As the internal structure of these bones is cellular or spongy, they contain only small separate portions of marrow, or a red medullary juice, like that in the vertebræ.

The fynovial substances of all these articulations are very fmall, but are accompanied by many other fatty moleculæ lying round each joint. The inner furface of the ligamentous tube, which lines the bony canal of the spine, is lubricated by an oily or adipose substance,

which shall be mentioned hereafter.

#### OF THE PELVIS.

THE cartilages of the offa innominata are not fo numerous as one might imagine on examining the skeleton. We are apt to think we see the dried remains of carticartilages on the spines of the os ilium, on the tuberofity of the os ischium, and on the grooves and notches which give passage to the tendons of muscles: but none of these incrustations are true cartilages, being for the most part tendinous, aponeurotic, or ligamentous; which being dried, look more like cartilages than the

true cartilages themselves.

The crust which covers the spine of the os ilium is chiefly tendinous, and a small part of it aponeurotic in adult bodies; but in children, and in very aged persons, it appears cartilaginous. In children, the parts which are not completely offisied, are easily taken for true cartilages; and in old age, the tendons are often hardened to so great a degree, as to have the very same apearance. The substance which covers the tuberosity of the os ischium is almost entirely tendinous; and that which lines the grooves and notches of the tendons is chiefly ligamentous.

The true cartilages of the offa innominata are five in number; three common, and two proper. The first and principal common cartilage is that which makes the symphysis of the offa pubis. It reaches from the interval between the spines of these two bones to the angle formed by the crura where they begin to separate. It is something thicker or broader at its upper part than for a considerable space lower down; but the inferior part is by much the broadest. It fills the angle already mentioned; and forms a kind of arch, which is

more confiderable in women than in men.

The two other common cartilages join the offa ilium to the os facrum, but are thinner than that of the offa

pubis.

The proper cartilages are those that line the acetabula. Concerning these, it has been already observed in the description of the Skeleton, that in the edge of each there is a notch or opening between the anterior and inferior parts; and that, in the cavity itself, there is a broad unequal shallow depression, reaching from

the

Pa

the

fm

ed

is

to

m

ar

in

te

of

It

P

ir

b

fi

li

a

t

0

1

IV.

ro.

nes

ne

he

s;

he

18

m

er-

rts

for

en

ne

fi-

nd

ns

ve

he

es

he

he

e.

irt

n-

le

18

m

la

u-

in

of

70

rc

m

e

the notch beyond the middle of the cavity. All the rest of the surface is covered with a very white shining smooth cartilage, which terminates precisely at the edge of the cavity.

The circumference of the acetabulum has, besides, a border of a particular kind; the substance of which is neither wholly cartilaginous, nor wholly ligamentous; but it may be rather placed among the liga-

ments.

The ligaments of the offa innominata are of two kinds; common, and proper. The common ligaments are those which go between these and the neighbouring bones, of which there is a considerable number, viz.

One superior ligament, fixed by one end to the internal labium of the posterior part of the spine of the offa ilium about an inch above the angle of that spine. It is about an inch in breadth, and is fixed by its other extremity to the whole inferior edge of the transverse process of the last vertebra of the loins.

One inferior and anterior, fixed by one end to the inner side of the angle of the spine of the os ilium; and by the other, to the superior and anterior part of the first false transverse process of the os sacrum. In this ligament there are transverse openings, which make it

appear more or less complex.

Several inferior and posterior, fixed by one end along the internal labium of the tuberosity of the spine of the os ilium, and by the other to the first three salse transverse processes; and from thence extending laterally over the marks of the salse oblique processes of the os facrum. To these must be added the ligaments by which the os semoris is joined to the os innominatum, which shall be described among the other ligaments of the thigh.

The principal proper ligaments are four in number: two called facro-fciatic, one broad and external, the other small and internal; one obturator; and one in-

guinal.

VOL. I.

D d

The

f

The broad facro-sciatic, or internal sciatic ligament, is flightly fixed to the infide of the tuberofity of the fpine of the os ilium; covers exteriorly the two posterior fpinous processes of that bone; and continues to be inferted along the anterior and external edge of the falle transverse processes of the os facrum. From thence this ligament, diminishing in breadth, descends oblique. ly towards the tuberosity of the os ischium; and is inferted immediately below the finus which lies between that tuberofity and the sciatic spine. This infertion is afterwards continued over the whole internal labium of the interior portion of the os ischium, and of the crus of that bone, and the inferior portion of the crus of the neighbouring os pubis. Thro' all this latter course of its infertion, that is, after its arrival at the tuberofity of the ischium, it produces a kind of ligamentous falx, one end of which is fixed to the bones, the other lies loofe; and by this fituation of the falx, it forms, together with the bones, a kind of deep channel or groove.

The internal facro-sciatic ligament adheres closely to the inside of the posterior portion of the former.—It is fixed internally to the edge of the inserior part of the fourth false transverse process of the os sacrum; and from thence all the way to the upper part of the os

coccygis.

From this infertion, it runs up a little obliquely to the spine of the ischium, to the sharp points and upper part of which it is fixed. During this course, it crosses the broad ligament, being closely united to the inside thereof, and loses but very little of its breadth.

By these two ligaments two distinct openings are formed; a large one, with the superior sciatic sinus; and

a small one, with the inferior sciatic notch.

The obturator ligament fills up all the great foramen, except the oblique notch at its upper part. It is fixed precisely to the edge of the circumference of that hole, from the anterior part of the oblique notch as far as the symphysis between the os pubis and os ischium. From thence

V.

nt,

the

101

be

he

CC

1e-

ın-

en

18

of

us

he

its

he

ne

ies

e-

.

to

-It

of

1;

OS

to

er

es

de

re

nd

n,

d

e,

10

m

:6

thence to the posterior part of the inferior notch, it is fixed to the internal labium of the edge of the circumference, forming a kind of small channel with the external labium; and afterwards it is fixed to the common edge of the foramen ovale, and cotyloid notch or opening.

By this disposition, an opening is left between this ligament and the superior oblique notch; and immediately below this common opening there are two small

perforations in the ligament alone.

On the infide of the upper and anterior part of the os pubis, there is a transverse ligament fixed by its upper part to the os pubis, from the oblique notch of the foramen ovale, all the way to the lower part of the fymphysis, at a small distance from the circumference of the last mentioned hole. This transverse ligament is about half an inch in breadth in an adult body; and posteriorly, below the superior oblique notch of the foramen ovale, it joins the obturator ligament, by means of a particular fold; and by parting from it afterwards, a deep narrow groove is formed between them, the transverse ligament being at this place supported by li-The inguinal ligagamentous fræna of different fizes. ment, called from the discoverer ligamentuum Fallopii, is an aponeurotic or ligamentous band, fixed by one end to the anterior superior spinous process of the os ilium, and by the other to the spine of the os pubis. The middle portion of it is very narrow, but it expands confiderably towards both extremities; it is closely joined to the muscles of the abdomen, and to the aponeurotic fascia of the thigh.

Besides these ligaments, peculiar to each os innominatum, there is another small, slat, and very long ligament, which runs transversely between the two angles of the cotyloid notch of the acetabulum, and may be

termed the proper or transverse ligament thereof.

The elastic border of the acetabulum may likewise be reckoned among the ligaments. It is a fort of additional

P

T

la

W

W

f

1

a

t

ſ

i

1

1

tional piece strongly united to the edge of that cavity, but easily yields both ways to any pressure. It may be stretched out by pulling, and recovers and contracts again when that force is removed. It is of a very singular texture, being composed of elastic sibres, interwoven together through its whole circumference; and which in several places, are by degrees inclined to the bony edge of the cavity. It makes an entire circle; and where it passes over the notch, the transverse ligament formerly mentioned serves to support it, as the bony edge of the cavity does through all the rest of the circumference.

Of the ligaments connecting the os femoris to the os innominatum, one is called the orbicular or capfular li-

gament, the other the round ligament.

The capfular ligament is very strong, and unequally thick. It surrounds the whole convex circumference of the supercilium of the acetabulum, to which it is strongly fixed for the breadth of near a quarter of an inch from the sharp edge outwards. Its insertion at the sharp edge of the acetabulum, joins that of the elastic border; the rest of the ligament is distinct from the border, and only touches it quite round; and where it passes over the notch, it is fixed in the transverse ligament.

The ligament which lies in the joint is not round as the name expresses. It is a flat cord, broad at one end and narrow at the other; and therefore somewhat of a triangular shape. By its narrow end it is inserted at the two angles of the notch of the acetabulum, and by the

other to the head of the os femoris.

It is broad, and may be reckoned the basis of the ligament; and from thence arise some distinct ligamentous silaments, which are inserted at different distances in the circumference of the rough impression of the acetabulum.

The rough unequal depression at the bottom of the accetabulum is filled by a broad flat substance, and covered by

ty, be

as

ın-

er-

nd he

e;

he

os li-

lly

ce

18

an he

ic

he

li-

as nd

ne

ne

i-

1-

e-

1.

by a fine membrane, through which amucilaginous liquor passes to moisten the joint, and facilitate its motions. This membrane likewise gives a kind of covering to the round ligament. As these bones have no internal large cavity, their substance being cavernous, they contain no medullary mass, but a medullary juice, which distils continually through the membrane with which they are lined.

#### OF THE SUPERIOR EXTREMITIES.

THE scapula in many subjects has a small cartilaginous border along its whole basis; which in children is

pretty remarkable, but in adults it disappears.

The glenoid cavity of this bone is covered with a cartilage, which is thicker toward the circumference than in the middle, and a little raifed above the edge of the bone. This thickness of the cartilaginous circumference makes the cavity greater than it appears in the skeleton; and sometimes, in place thereof, there is an additional border, which is thickest at the circumference of the cavity than toward the bottom, and very narrow. It is of a pliable slippery substance, yet something different from that of a cartilage; resembling, in some measure, the border of the acetabulum of the os innominatum.

The fmall cartilaginous furface of the acromion, mentioned in the treatile of the dry bones, is thicker in the

natural state, and very little convex.

The small triangular surface at the extremity of the spine of the scapula, near the basis, is covered with a very thin cartilaginous lamina; but, being transparent, it does not appear very white. There are no other cartilages commonly found in the scapula: though we sometimes observe in dry bones several places which seem to have been cartilaginous; but this is owing to the dried remains of ligaments and tendons.

The sternal extremity of the clavicle is crusted over with a cartilage which is a little convex, and covers its

Dd3

whole

whole triangular furface; besides which it has another moveable interarticular cartilage, resembling that at the articulation of the lower jaw. The small cartilaginous surface of the scapular extremity of the clavicle, answering to that of the acromion, is much thicker in fresh than in dry bones, and appears, like that of the acromion, to be a little convex.

Between these two cartilages of the clavicle and acromion, there is in some subjects a thin interarticular car-

tilage very smooth on both fides.

The articulation of the acromion with the extremity of the clavicle, is strengthened quite round by several small strong ligaments, which go from one bone to the other. These ligaments lie very near each other; and are withal so tightly braced over the joint, as to hide it altogether; and they appear more like a cartilaginous covering than a ligamentous texture. The internal surface of these ligaments is lined with the capsula of the joint. When the small interarticular cartilage is found, its whole circumference is connected to these ligaments.

The articulation of the clavicle with the sternum is fustained by several ligaments fixed by one end, round the sternal extremity of the clavicle, near the edge of the triangular surface; and from thence passing over the interarticular cartilage, are inserted by the other end in

the sternum already mentioned.

There is a long narrow strong ligament, which goes from one clavicle to the other behind the surca of the sternum, being fixed to the internal angle of the contiguous extremities of the bones. It may be called the interclavicular ligament.

The neck of the scapula at a small distance from the edge of the glenoid cavity, gives insertion to the edge of the capsular ligament or mucilaginous bag, and to the articular ligaments of the joint of the scapula and os

humeri.

Besides these articular ligaments of the scapula there are

V.

her

us

er-

ni-

0-

ır-

ty

al

he

1;

to

a-

np-

ge

le

is

d

of

e

n

S

e

e

are three ligamentous cords fixed to the tuberofity of the coracoid proceses; two of which, by their other extremities, are inserted into the oblique eminence on the lower side of the humeral extremity of the clavicle; the third under the acromion. There is likewise a thin slat broad ligament, reaching between the crista of the spine of the scapula, and the edge of the inserior costa. The cartilage by which the head of the os humeri is covered, is gradually thicker toward the middle than toward the edges.

The four furfaces of the tuberofities which appear cartilaginous in dry bones, ferve only for the infertion of the tendons of four muscles which move the os hu-

meri on the scapula.

The channel or finus between the tuberofities is partly covered by a thin crust, which appears rather ligamentous than cartilaginous; and partly by a tendinous stratum.

The trochlea, and small head of the lower extremity of the os humeri, are covered by a common cartilage, in which the same proportion of thickness is observable, as in that of the upper extremities. This holds pretty generally of all the convex articular cartilages.

The fosfulæ near the pulley and small head are covered with a kind of cartilaginous or ligamentous var-

niff

The capfular or mucilaginous ligament loofely furrounds the whole articulation of the scapula with the head of the os humeri. From its insertion round the edge of the glenoid cavity already mentioned, it is continued over the hemisphere of the head of the os humeri, and fixed near its edges towards the muscular surfaces of the great and small tuberosities. Afterwards parting from them on both sides in the large space lest between the two tuberosities, that is, between the small tuberosity and the lowest surface of the great tuberosity, it runs down gradually on the neck of the bone below the lowest part of the cartilaginous hemi-

Dd4

fpere. In all this course, the capsula is closely fixed in the bone, excepting in the small space lest between the two tuberosities, that is, at the channel or sinus already mentioned; where it forms a production like the tube of a funnel, proportioned to the capacity of the channel, and strongly fixed in the upper portion thereof. This membranous tube is the vagina of the interarticular tendon of the biceps muscle. The true ligament appears to be composed of two kinds of ligaments closely united; namely, of a capsular ligament which surrounds the whole articulation; and of several true ligaments, which run over, and closely adhere to, the former at different distances.

Thus the capfula, or mucilaginous bag of this articulation, is in part strongly united to four flat tendons inserted in the two tuberosities; and in part covered by true ligamentous bands, which, between the four tendons, and on both sides of the first and last, form a considerable thickness. The rest of the space between the first or superior plane of the great tuberosity and the small tuberosity, is so little provided with ligamentous sibres, that it has been believed to be altogether without them; and anatomists have satisfied themselves with telling us, that in these places the capsular ligament is very rough on the outside, but shining and smooth on the inside.

The tendon of the long head of the biceps, which has been already mentioned in speaking of the production of the capsular ligament of the head of the os humeri, is contained in the articulation after the same manner as the ligamentum rotundum of the head of the os femoris.

On the body of the os humeri, there are two particular intermuscular ligaments, which are long, narrow, thin, but strong, fixed by one end along the two lower thirds of the bone, and reaching to both condyles.

The

IV.

l in

the

ady

ıbe

an.

of.

cu.

ent nts

ich li-

or-

ti-

ons by

n-

a

en

nd

n-

er

es a-

nd

h

C-

l-

of

0

.

e

The lower extremity of the os humeri is joined to the bones of the fore-arm by two fasciculi of ligamentous fibres; one fixed to the internal condyle, the other to the external. Each fasciculus is composed of fibres, closely joined together at the condyle, and afterwards parting in distinct fibres like the foot of a goose.

The capfular ligament is fixed to the condyles, and there covers them; and afterwards it is fixed round both fides of this lower extremity above the fosfulæ. Its infertion in these fides is archwise; so that it is there at a much greater distance from the articulation than at the condyles. The fosfulæ are lined with a very

thin cartilage.

This capfula appears to be strengthened by a ligamentous web, the sibres whereof cross each other in different directions: but we must not take for ligamentous silaments some tendinous sibres of muscles, to which the capsula adheres very closely. It appears larger and looser when the muscles are separated from it, than in its natural state when closely united to the muscles.

The two figmoid cavities in the upper extremity of the ulna are covered by a cartilage common to both; which is a little interrupted about the middle of the edges of the cavities by the transverse notches mentioned in the Description of the Bones. This cartilaginous crust seems to be thicker at the edges than in the middle.

The inferior extremity, or small head of the ulna, is crusted over by a cartilage round its cylindrical border, in the notch near the styloid process, and for some

space on the process itself.

The cartilage which covers the head of the radius is likewife turned over its cylindrical border; and a lateral portion of the muscular tuberosity, immediately below the neck, is also covered with a thin shining cartilage.

All the concave fide of the basis of the radius is likewise wife cartilaginous, and often divided by a small cartilaginous prominent line. The lateral notch of the bafis is also covered by a continuation of the same carti-

lage.

At the basis of the radius there is likewise a particular additional cartilage, or triangular production, longer than it is broad, very thin, and rather flat than concave on both its fmooth fides. It is fixed by its basis, or shortest fide, to the lateral figmoid notch of the basis of the radius, in fuch a manner, that one fide of it is on a evel with the large cartilaginous furface of the basis of the bone, and its apex directly opposite to the styloid The other fide touches the flat extremity of the small head of the ulna, but is not fixed to it. This cartilage may be termed the interarticular cartilage of the joint of the wrift. It is tied to the radius by very short ligaments; and, playing on the small head of the ulna, it follows all the motions of the radius. It is therefore a fort of particular production of the lower fide of the basis of the radius, and fills, in the natural flate, the void space which, in the skeleton, appears between the end of the ulna and the neighbouring bone of the carpus.

Some of the ligaments of the bones of the fore-arm are common to them with the os humeri; fome common to them with the bones of the hand; and fome are proper. These last are two in number; one called the interoffeus ligament of the fore-arm, and one which may be termed the coronary ligament of the radius. To these may be added the annular ligaments, which only ferve for the passage of tendons; and other ligamentous expansions, which may be called muscular liga-

ments.

The interoffeous ligament is fixed by one edge along the sharp angle of the ulna, and by the other along that of the radius. It is principally composed of two very strong planes of fibres, which cross each other at oblique angles, and leave holes at different distances for the pal-

fo

n

re

0

C

0

u t ti-

Da-

ti-

u-

er

ve

or

of

a

of

id

of

lis

of

ry

ne

is

er

al

rs

g

n

1.

C

d

h

0

.

passage of the blood-vessels. This ligament ties the two bones closely together, and the two planes ferve for the infertion of the feveral muscles. In the supination of the hand, it is very tightly braced; but in pronation, it is folded a little lengthwife.

The coronary ligament of the radius, is a fort of ligamentous hoop, furrounding the circular circumference of the head of that bone, reaching from one fide of the small lateral figmoid cavity of the ulna to the other, in an arch which is about three quarters of a circle. It is very strong, and comes near the folidity of a cartilage. The fide next the radius is very smooth; and though it connects that bone very closely to the ulna, yet it leaves it room enough to turn in the mo-

tions of pronation and supination. The capfular ligament of the joint of the elbow runs down from its infertion in the os humeri, already defcribed; and is fixed in the olecranon round the edge of the great figmoid cavity, including both the apex of the olecranon and of the coronoid process. It likewife runs over the head of the radius; and is fixed to the coronary ligament quite round. Thus it completely furrounds the articulation of these three bones; and ferves to contain the mucilaginous liquor or fynovia furnished by the bodies called the glands of the joint, and by fatty fubstances, both of which are found in abundance near the extremity of the ulna.

The true common ligaments by which the os humeri is connected to the bones of the fore-arm, called lateral ligaments, are the two fasciculi which, after being inferted in the condyles of the os humeri, are expanded like the foot of a goofe. That which is fixed to the inner condyle may be called brachio-cubitale, and the other brachio-radiale.

The brachio-cubital ligament running down over the capfula, to which it closely adheres, below the great edge of the trochlea of the os humeri, is inferted like radii (of which its other extremity fixed to the condyles '

is

is the centre) on the fide of the great figmoid cavity of the ulna. It is covered on the outfide by feveral tendons, which adhere closely to it, and feem to

strengthen it.

The brachio-radial ligament is disposed much after the same manner, but is of a greater extent. It is expanded from the external condyle of the os humeri, as from a centre; and is inserted round the coronary ligament, and from thence all the way down to the neck of the radius, and also in the neighbouring parts of the ulna. Through all this passage it covers the scapular ligament, and is covered by several tendons, adhering closely to both.

Of the ligaments by which these bones are connected to those of the hand, one is like a roundish cord, fixed to the styloid process of the ulna, and from thence passes directly over the os cuneiforme of the carpus; to which, and to other bones, it is inserted. Another pretty broad ligament is fixed to the point of the radius, and by its other extremity to the bones of the

carpus.

From this styloid ligament of the radius, along each edge of the basis of that bone, are ranks of ligamentous sibres, lying much in the same direction with the ligament itself, and continued all the way to the styloid ligament of the ulna: those next the ulna inclose the interarticular cartilage of the basis of the radius; and, near the styloid ligament of the ulna, there is a particular fasciculus inserted to the point of that cartilage.

All these ligaments surround and cover the capsular ligament so closely, that they can hardly be distinguished from it. The capsula is likewise in part covered by a portion of a great oblique ligament; which being, by a very broad insertion, fixed to the large extremity of the radius, about two singers breadth above the styloid apex, afterwards crosses obliquely, partly over the convex side of the basis radii, and partly over that of the carpus; and then turning toward the os orbiculare,

10

t

1

t

ł

1

V.

ity

ral

to

er x.

as

a-

k

le

ar

d,

c

r

1-

e

h

C

is inferted into it. It is called the external transverse ligament of the carpus; and may likewise be named the great oblique ligament of the wrist.

There are several small annular ligaments placed at different distances on the convex side of the basis radii, from its styloid process to its articulation with the extremity of the ulna. They are at least six in number;

some of them being often double or triple.

The first is fixed in the styloid process; the second, in the groove near that apex; the third, in the small narrow or middle groove; the fourth, in the groove next the former; the sifth, in the corner of the semilunar notch of the basis, at its articulation with the ulna; and the sixth, to the extremity of the ulna near the styloid

process.

These particular ligaments are almost wholly covered by the great oblique ligament already mentioned; and are fixed as strongly to it by one side, as they are to the bones by the other. They are all very strong; and their concave sides, serving for fræna to the tendons of several muscles that pass over them, are very smooth, and accompanied with thin mucilaginous vaginæ, which allow them to move easily.

To these we may add the ligamentous expansions with which several muscles are covered, and separated from each other as by so many distinct septa; which are all very thick and strong where they are fixed to the bones. One kind of them may be termed ligamentous bands, or muscular vaginæ; the other, ligamentous

fepta, intermufcular ligaments, &c.

All the bones of the carpus, metacarpus, and fingers, are crusted over with cartilages at the places which play upon each other; but in fresh subjects, they are thicker, softer, and whiter, than in the skeleton. In adults, their figure remains the same in both; but it changes in the dry bones of younger subjects, and in those of children it is quite different. The impressions and notches in which the bodies called mucilaginous glands

glands are lodged, are most sensible in the cartilages of

fresh bones, because of their thickness.

The ligaments of the carpus are very numerous. Some of them tie each bone to one or two neighbouring bones in the fame rank; and these are composed of a great number of filaments, but so very short as to allow these bones only a small degree of motion. Some of them tie the bones of one row to those of the other; which are likewise composed of many filaments, but not so short as the former; and therefore allow these bones a more manifest motion, as we see in bending the wrist. Lastly, there are other ligaments of the carpus, by which the three bones of the first row are connected to the bones of the fore-arm: and to these may be added the ligaments by which the bones of the second row are joined to those of the metacarpus and first phalanx of the thumb.

We have already described all the ligaments belonging to the articulation of the carpus with the bones of the fore-arm, except their insertions in the carpus. The styloid ligament of the radius is fixed round the neighbouring tuberosity of the os scaphoides. The styloid ligament of the ulna is fixed, first to the os cuneiforme, and then to the os unciforme; from whence it is a little stretched over the fourth bone of the meta-

carpus.

The ligaments which lie between the two former round the basis of the radius, and a small portion of the head of the ulna, are fixed round the common convexity of the three first carpal bones, as is also the mucilaginous capsula by which these ligaments are lined.

Besides these small short ligaments belonging to each bone in both rows, the rough surfaces of all the bones, especially those which form the convexity of the carpus, give insertion to a great many ligamentous fasciculi, stretched over and closely united to the former small ligaments, and serving probably to strengthen them. Some fasciculi of the same kind are found on the concave

of

us.

ng a

al-

ne

r;

out

ese

ng

he

re

efe

he

nd

g-

of

S.

he

ne

1-

ce

a.

er

e

1-

1-

h

١,

e

cave fide of the carpus; but they are fewer in number, and not fo strong.

Here is likewife a confiderable ligament, called the inner transverse ligament of the carpus. It was formerly called an annular ligament; and may still very justly

retain that name on account of its use.

The bones of the metacarpus, besides the short ligaments by which they are tied to the fecond row of the bones of the carpus, have feveral others, by which both their bases and heads are connected together. The bases of the third and fourth bones are not so closely tied as the rest; and therefore they have a very fensible motion; which, however, is greater in the fourth than in the third.

The heads of these bones are firmly tied to each other by a strong transverse ligament, situated in the palm of the hand, and fixed by distinct productions in the neighbouring part of the heads, in fuch a manner as to form, in the spaces between the heads, a kind of perforated fræna, through which the tendons of the flexor muscles of the fingers have a free passage; and these fræna are also supported by other aponeurotic expations.

The first phalanx of the thumb is fixed to the os trapezium by fhort ligaments, which pass obliquely over the articulation. The first phalanges of the other four fingers are joined to the heads of the metacarpal bones almost in the same manner, and by ligaments like the former; which are itrengthened by adhering to the transverse ligament already mentioned. The fecond phalanx of the thumb is joined to the first by ligaments

of the same kind.

The third phalanx of the thumb is joined to the fecond, the fecond phalanges of the other fingers to the first, and the third to the second, by lateral ligaments, almost in the same manner as the bones of the forearm to the os humeri; that is, these ligaments spread from a point fixed in the lateral tubercles of the heads

of the phalanges, and are inferted by their other extremity like radii to the basis of the neighbouring pha-

langes.

The two first phalanges of each finger have a very strong ligamentous vagina inserted in the rough lines, or ridges, on their flat sides: these vaginæ are lined with a mucilaginous membrane, which runs like a tube from one phalanx to the other over the articulation. They serve for fræna to the slexor muscles of the singers; the tendons of which pass through them.

### OF THE INFERIOR EXTREMITIES.

No part of the os femoris is covered with cartilage, excepting the uniform convexity of its head, and the articular portion of the lower extremity. The trochanters have no true cartilage; what looks like it being only the remains of tendinous infertions, as was observed of the spine of the os ilium. The cartilaginous substance which, to a certain age, unites the epiphyses to the body of the bone, does not belong to this place, because it is only found in the time of youth, and in adults is converted into bone. The cartilaginous matter by which the head of the os femoris is cemented, deserves, however, to be observed; because that epiphysis has been separated by violent falls.

The convexity of the head of the os femoris, all the way to its symphysis with the neck, is covered by a very smooth shining cartilage. We have already remarked, in describing the dry bones, that a little below the middle of this convexity, and somewhat towards the back-part, there is a depression in form of a crescent, the cartilage being here interrupted by the insertion of the interarticular ligament of the head of the os fe-

moris.

The cartilage which covers the lower extremity of this bone, is exactly fitted to the femi-oval convexity of the inferior furface of each condyle, and to the pulley formed by their union.

In

Con

Part

face T to the

rou

of t

gel

hur

the who fem the par fibrilique

ferification for the

mic two her to

diff two wit

In the posterior part of the lateral tuberosity of each condyle, there is another kind of cartilaginous surface.

The os femoris is connected by its upper extremity to the os innominatum, and by the lower to the bones of the leg by several ligaments. The ligaments of the upper extremity are two in number; one which surrounds the whole articulation with the acetabulum, and one contained in it.

The first is termed the orbicular or capsular ligament of the head of the os semoris; the other, the round or

internal ligament.

The capfular ligament is the most considerable, largest, and strongest, of all the articular ligaments of the human body. It is fixed quite round the border of the acetabulum, and from thence largely surrounds the whole head and superior portion of the neck of the os semoris, and is closely inserted to the lower portion of the neck that is between its basis and middle narrow part. This ligament is made up of several sorts of sibres, the chief of which are longitudinal and oblique; and it is much thicker and stronger in some parts than in others. It is very thick between the inferior anterior spine of the os ilium all the way to the small tuberosity, which unites, as it were, the basis of the great trochanter with the basis of the neck.

It is likewise very thick between the same and the middle part of the oblique rough line, observable between the tuberosity and the little trochanter. And here likewise it is strengthened by a bundle connected to the passage of the tendon of the iliac muscle, and to the inferior portion of the oblique rough line. The disposition of the ligamentous sibres, of which these two thick portions are composed, form a fort of triangle with the oblique rough line, which terminates the basis

of the neck.

At the posterior and upper portion of this ligament, there is another thick portion formed by oblique fibres;

Vol. I. E e one

one end of which is fixed between the inferior edge of the acetabulum and the passage of the tendon of the external obturator muscle; the other to the upper part of the small tuberosity of the great trochanter, already mentioned.

The portion and lower part of it is thinner and shorter than the rest: but even this is strengthened by a band of pretty strong sibres, which, from the whole crists of the os pubis, runs down obliquely near the fore-side of the notch of the acetabulum; and is sixed in the upper part of the basis of the neck of the os semoris, immediately above the small anterior tuberosity of the great trochanter.

The internal ligament resembles a slat cord, being composed of a bundle of sibres closely interwoven. One end of it is in a manner divided into two slat bands, which are fixed to the corners of the notch of the ace-

tabulum.

From this infertion it runs obliquely backward, and a little upward, between the glandular-like substance within the acetabulum and the cartilaginous convexity of the head of the os femoris, and ends in the upper part of the small semilunar fosfula; which may be reckoned the pole of that convexity. This insertion is oblique, a little rounded on the upper part, and slat on the lower; and in some subjects there is a fort of depression in the head of the bone for the passage of the ligament.

The ligaments of the lower end of the os femoris, by which this bone is connected with those of the leg, are fix in number; one posterior, two lateral, two middle

or crucial, and one capfular.

The crucial ligaments lie within the joint, and are fixed by one end to the back part of the notch which parts the two condyles. They are furrounded by the capfular ligament; but all the rest lie on the outside of it, being closely joined to it. Of the two lateral ligaments,

Part

mentube T

a litt

T

three the iftand of the greathe the i

in the cluft netre ramiftain

form

runt addi T peri

The depinal vext post mid the

whi

by t

the

ments, one is internal and broad, being fixed to the

tuberofity of the internal condyle.

The posterior ligament is broad and thin, being fixed a little above the convexity of the external condyle; from whence it descends obliquely behind the great

notch and internal condyle.

The capfular ligament, glued, as it were, to the three former, as has been faid, is fixed quite round the inferior extremity of the os femoris, at a small distance above the anterior, lateral, and posterior parts of the cartilage, and above the posterior part of the great notch; and from the cartilage and notch through the small space upward, already mentioned, it covers the bone; and afterwards is inverted downwards, to form the capsula for the synovia of the joint.

The marrow of the os femoris lies in a large mass, in the middle cavity of the bone; and in small distinct clusters, in the cells of each extremity. The first is penetrated at different distances by the bony silaments or ramifications of the reticular texture; and thereby suffained in violent motions and shocks, as in leaping,

running, &c.

The tibia has four or five proper cartilages, and two

additional ones.

The two proper cartilages which cover the two fuperior furfaces of the head of the tibia are the thickest. They are both gently hollow; but the internal is more depressed than the other. The back-part of the external is insensibly depressed, and thereby a fort of convexity is formed. Anteriorly they join each other; posteriorly they are parted by a shallow notch; in the middle they are separated by the articular tuberosity of the head of the tibia, which is likewise partly covered by them on each side.

The third proper cartilage covers the small surface which lies on the lower part of the external condyle.

The fourth covers the lower surface of the basis of the tibia, being continued over the outside of the inner E e 2 ankle.

There are likewise superficial cartilaginous incrustations on the back-part of this basis behind the inner ankle, and also on the back-side of the outer ankle, all for the passage of tendons.

The additional cartilages of the tibia are two in number, called femilunar from their figure, and interarti-

culas from their fituation.

Each of these cartilages is in shape of a crescent or Roman C. Their convexity, or greatest curvature, is very thick; their concavity, or smallest curvature, very thin, something like the edge of a sickle. They lie on the two upper surfaces of the head of the tibia; their thickest part or convexity corresponding with the edges of the head, and their thin sharp edges to the middle of each surface; their extremities or cornua being turned toward each other.

Each cartilage is broad enough to cover about two thirds of the furface of the tibia on which it lies, leaving one third bare in the middle. Their under fides are flat, the upper fides hollow; and, together with the middle portons of the furfaces of the head of the tibia, form cavities proportionable to the convexities of

the condyles of the os femoris.

The fibula has two cartilages; one lying on the upper extremity of that bone, for its articulation with the small cartilaginous surface in the head of the tibia: the other cartilage covers the inside of the inferior extremity, or of the outer ankle; near the point of which, posteriorly, there is a superficial cartilaginous incrustation for the passage of the tendons of the musculi peronæi.

The cartilage at the upper extremity of the fibula feems to be thicker than that at the lower extremity.

The patella, which belongs properly to the tibia, has a pretty thick cartilage on its posterior side, divided by a superficial longitudinal rising into two parts, proportioned to the two portions of the pulley of the os semoris.

The

of the Tenton fupe ante

Part

T

the T thic med tren the

of the

os for the cial fior bet tior

of the bro in i

gar deg are fup The connections between the os femoris and bones

of the leg are as follow.

The innermost and broadest of the two lateral ligaments is fixed pretty low down on the inner side of the superior part of the tibia, between the beginnings of the anterior and internal angles. It is likewise joined to the edge of the internal semilunar cartilage.

The external lateral ligament, which is narrower and thicker than the former, is fixed partly in the tibia immediately above the fibula, and partly in the upper extremity of the fibula; and is joined also to the edge of

the external femilunar cartilage.

Both these ligaments lie a little behind the middle of the articulation.

The posterior ligament is fixed by several expansions

to the posterior part of the head of the tibia.

One of the crucial ligaments is fixed by one end to the internal fuperficial impression in the notch of the os semoris; and by the other to the notch in the head of the tibia, behind the cartilaginous tubercle which lies between the two superior surfaces. The other crucial ligament is fixed by one end to the external impression, in the notch of the os semoris; and by the other, between the anterior portions of the surfaces just mentioned, before the cartilaginous tubercle.

These two ligaments are composed of several series of sibres. The first, which is internal in respect to the os semoris, and posterior in respect to the tibia, is broader and stronger than the other; which is external in respect to the os semoris, and anterior in respect to

the tibia.

The femilunar cartilages have likewise particular ligaments besides their connections with the lateral ligaments of the tibia. Their cornua do, in some measure, degenerate into short strong ligaments; by which they are fixed to the cartilaginous tubercle between the two superior surfaces of the tibia, and likewise communicate by some portions with the crucial ligaments.

Ee 3 They

They have also a common ligament; which, like an arch, runs transversely between the anterior convexity of the one to that of the other.

These cartilages, therefore, have three sorts of connections. They are fixed to the tibia by the ligaments of the cornua; to each other by the transverse ligament; and to the os semoris by their communications with the crucial ligaments, and by their adhesions to the lateral and capsular ligaments.

The patella is fixed to the tubercle of the tibia by a broad and very strong ligament, which runs down directly from the apex of the patella, and is sometimes further strengthened by some sibres of a considerable tendon inserted into the upper part of that bone.

It has likewise small lateral ligaments fixed to the lower part of its edge on each side; which parting gradually from the great ligament as they run down, are inserted anteriorly, and a little laterally, to the edge of the head of the tibia.

The capfular ligament, formerly mentioned, is fixed round the edge of the head of the tibia, and in the edge of the patella; fo that the patella itself forms a portion of the mucilaginous capsula of the joint of the knee.

The crucial ligaments, and those of the semilunar cartilages, are included within this capsula; but the lateral and posterior ligaments, and those of the patella, lie without it, being closely joined to its outer surface.

This capfula is likewise joined to a considerable portion of the circumference of the semilunar cartilages; and it is also strengthened at different distances by several distinct series of ligamentous sibres. The inside of it is smooth and shining; and it is very thin where it is not covered by tendons. It not only contains and surrounds the ligaments already named, but also surnishes them with a very sine vagina.

There is likewise a very thin ligament fixed by one end to the lower part of the cartilaginous side of the patella;

fou

Par

pate

the

fat

the po the ing what the m

tic

w an lo

t

patella; and by the other to the anterior part of the great notch, between the condyles of the os femoris; the use of which seems to be, to prevent the articular fat from being compressed in the motions of the knee.

The fibula is joined to the tibia by nine ligaments; four at each end, and one in the middle called the in-

terosseous ligament.

The ligaments at the upper end of the fibula are short, very strong, more or less oblique, and compound. Two of them are anterior, two posterior; and they lie on each other, the superior ligaments surrounding the articulation more closely than the inferior, which leave a small void space, and are weaker than the former. They are all glued to the capsular ligament, which runs in between them and the articulation; and they are inserted round the edges of the cartilaginous surfaces in each bone.

The ligaments of the lower extremity of the fibula, which runs below the tibia, and forms the outer ankle, are much stronger, thicker, more complex, broader, longer, and more oblique, than those of the superior extremity; and are disposed much after the same man-

ner; that is, two before, and two behind.

They are fixed to the anterior and posterior edge of the lateral depression at the inferior extremity of the tibia; and from thence they run down on the lower end of the fibula. The two inferior ligaments are strongest; and they are fixed anteriorly and posteriorly at the lower end of the outer ankle.

The two fuperior are fixed more closely, and nearer each other; but still there is a small space between them filled with fat.

As the two bones touch each other only by the upper part of the cartilaginous furface of the outer ankle, and the small cartilaginous border on the lower edge of the depression of the tibia, the middle space between them is filled by a fort of capsular ligament, which lines each side of the bones; and is continued down to the

Ee4

true

true articulation of the external ankle with the inferior

edge of the basis of the tibia.

The middle or interoffeous ligament of the two bones of the leg, so called because it fills up all the space lest between them, being stretched from one to the other, is fixed along the posterior external angle of the tibia

and the neighbouring angle of the fibula.

It is made up chiefly of two planes of very oblique ligamentous fibres, which cross each other, and at different intervals seem to be multiplied. It is perforated both above and below, and sometimes in several places besides, for the passage of the blood-vessels and nerves. It serves chiefly as a ligamentous septum for the infertion of muscles; in which respect it supplies the place of bones, and seems partly to be a continuation of the periosteum of the bones of the leg.

At the lower part of each ankle there are commonly three strong ligaments for the connection of the bones of the tarsus with those of the leg; one that runs forwards, one that runs backwards, and one that runs

more or less directly downwards.

There are some other ligamentous expansions to be found in the leg, but these belong properly to the muscles.

The marrow of these bones lies in large masses in the great cavities, and in distinct moleculæ in the spon-

gy parts.

The fynovial fubstances lie in the small spaces, depressions, and superficial notches, near the edges of the cartilages, and within the capsular ligaments of the joints.

The fynovial fubstances of the knee, which lie near the edges of the patella, are the most considerable, being disposed in form of fringes, and supported by a

great quantity of fatty matter.

This common mass is contained within the capsular ligament; and on the side of the joint is covered by a

face ed l

Part

very

ed the more of t

low

in for ed me ligation

fur fur inf

firl

os th th

th

fu la or ti

o

very fine membrane, which likewife lines the inner fur-

face of the ligament.

The superior portion of this fat is as it were supported by the small ligament, fixed in the anterior part of the great common notch of the condyles of the os femoris, and which from thence runs to the upper part

of the patella.

There are other fatty substances, both above and below the edges of the semilunar cartilages, and likewise in the ham; some whereof serve for the joint, the rest for the crucial ligaments. These last lie in folds formed by the internal membrane of the capsular ligament, which give particular coverings to the crucial ligaments, and to the other bundles of ligamentous sibres near them.

The astragalus is covered by three cartilages. The first covers the three surfaces, which make the convex part and sides of the pulley; the second, the concave surface of its anterior part; and the third, the convex surface of its anterior part, being continued over the inferior part, so far as to form three other small surfaces, one of which can scarcely be called articular.

The first of these cartilages is for the articulation of this bone with the tibia and sibula; the second, for the os calcis; and the third, for the os scaphoides. Two of the inserior surfaces formed by the continuation of the third cartilage are for the articulation of this bone with the os calcis; the third contributes to the formation of

a channel for the passage of a tendon.

The os calcis has four cartilages; of which three are fuperior, one large and two small, for its triple articulation with the astragalus; the fourth is anterior, for the os cuboides. To these must be added a small thin cartilago-ligamentous substance, under the tubercle on the outside of this bone.

The os scaphoides has two cartilages; one posterior, for its articulation with the astragalus; and one anterior.

C

b d

tł

p

a

fi

0

n

f

t

rior, divided into three parts for the three offa cuneiformia.

The os cuboides has two remarkable cartilages; one posterior, for its articulation with the os calcis; and one anterior, lying in two planes for its articulation with the two last metatarsal bones. It has likewise a cartilage on the infide for the os cuneiforme, which is next to it; and one on the lower fide, covering a part of the ob-

lique eminence situated there.

The three offa cuneiformia have each of them a posterior cartilage for their articulation with the os fcaphoides; and one anterior for the three first metatarsal bones: they have likewise small cartilaginous surfaces, on their lateral fides, for their articulations with each other; and besides, the first and third bones are joined thereby to the lateral parts of the basis of the second metatarfal bone, and the third to the os cuboides.

The bases and heads of the metatarfal bones are co-

vered with cartilages.

The phalanges have cartilages in the fame manner at their bases and heads, excepting at the heads or extremities of the last.

The fefamoid bones are covered with cartilages on that

fide by which they slide on other bones.

The foot being made up of many bones, must, besides those ligaments by which it is tied to the bones of the leg, have feveral others, to connect not only the three parts of which it is composed, but also the particular

bones belonging to each part.

The ligaments connecting the ankle to the foot have been already mentioned. Those of the inner ankle are all fixed to the upper fide of the astragalus. The most anterior is pretty broad, and fometimes feems to be joined in one with the middle ligament. It often confifts of feveral distinct parts, like so many bands interlarded with fat.

The anterior and middle ligaments of the outer ankle being more or less broad, are fixed to the outside of the aftraastragalus; the posterior, which is narrowest, and pretty thick, is chiefly fixed to the outside of the great portion of the os calcis.

All these ligaments lie on the outside of the capsula which furrounds the articulation of the astragalus with

the bones of the leg.

The ligaments, by which the bones of the tarfus are connected with each other, are short, flat, of different breadths, and run from one bone to another in different directions. They are all superficial, except one by which the astragalus is tied to the os calcis; and for the most part are either fuperior or inferior, the lateral ligaments being but very few in number.

Some of them are partly common to several bones, and partly belong only to two: that is, the fuperficial strata of their fibres run over one bone into the following, and fometimes further; but the strata next the articulation are generally confined to two bones only.

The astragalus is tied to the other bones of the tarfus by feveral true ligaments:—To the infide of the os calcis, by a ligament which comes from the posterior internal tuberofity of the body of the astragalus; and is fixed in an inequality behind the laternal process of the os calcis:-To the infide of the fame bone, by a ligament which comes from the lateral process of the os calcis, to a fort of cartilaginous production on the infide of the neck of the astragalus:—To the outside of the fame bone by two ligaments, which come from the edge of the oblique inferior depression of the astragalus; and afterwards separating a little, are fixed to the outside of the great apophysis of the os calcis; one forward, which feems to fend off a small portion to the os cuboides; the other backward, of different breadths: -To the os fcaphoides superiorly, by a ligament, which goes from the neck of the aftragalus to the upper part of the os fcaphoides, and from thence is extended to the middle of the os cuneiforme: -To the same anteriorly, by two ligaments; one of which

which is a continuation of that which goes from the lateral process of the os calcis, to the cartilaginous production of the astragalus; the other is nearly the same production, being partly covered by the former, and fixed to the tuberosity of the os scaphoides:—To the os calcis, by a ligament, which comes from the oblique inferior depression of the astragalus, and is fixed to the oblique superior depression of the os calcis. If to these principal ligaments of the astragalus, we add several others less remarkable; and also those by which it is tied to the ankles, their number will be very considerable.

The capfular ligaments go very little farther than the edges of the articulations of this bone with the rest. They adhere very closely to the true ligaments, and are

covered and hid by them.

The os calcis is joined to the outer ankle and astragalus by the ligaments already described. It is likewise connected to the os scaphoides and os cuboides by se-

veral ligamentous planes.

It is connected to the os scaphoides, (1.) By a continuation of the ligament that goes from its lateral or internal process, to the cartilaginous production of the astragalus. (2.) By a ligamentous plane, which goes from the inferior tuberosity of its great process, and is fixed to the inferior part of the circumference of the os scaphoides. (3.) By a narrow ligament, which goes from the superior and internal part of the same process, and ends in the nearest part of the circumference of the os scaphoides.

It is connected to the os cuboides, (1.) By a ligament, or rather by several ligamentous fasciculi, which go from the extremity of its oblique superior depression to the contiguous angle of the os cuboides. (2.) By one lying between the first and the small external lateral tuberosity of the os calcis, and inserted to the os cuboides near the first. (2.) By one which is fixed to the exterior part of the great process of the os calcis, and to the contiguous part of the os cuboides. (4.) By

fit

OS

th

ed

ch

af

fo

ne

to

pa

th

it

fc

0

th

0

V

O

tl

ti

0

a

a

fi

a pretty broad plane, which covers the inferior part of the os calcis, and which, from the anterior tuberofity, spreads over the contiguous inferior part of the os cuboides, and ends in the oblique eminence of (5.) By a broader plane, which having fillthat bone. ed the lower part of the lateral cavity of the os calcis, is chiefly inferted to the contiguous angle of the os cuboides. The capfular ligaments agree with those of the

astragalus.

The os fcaphoides is tied to the aftragalus and os calcis in the manner already mentioned.—It is likewife joined to the os cuboides and the offa cuneiformia by feveral ligaments. On the outfide, or that next the os cuboides, there is one which connects it to the contiguous angle of that bone. On its upper part, two go from its circumference; one to the fecond, the other to the third os cuneiforme. On its inner fide, it is joined to the convex part of the great os cuneiforme by two ligaments. On its lower fide it has four; of which the first appears as it were double, going from the tuberofity of this bone to the basis of the first os cuneiforme. The fecond and third go obliquely to the other two offa cuneiformia; the fourth is a little transverse, being fixed to the inferior internal angle of the os cuboides.

The os cuboides, befides the ligaments which tie it to the aftragalus, os calcis, and os fcaphoides already mentioned, has others which connect it above, below, and on the outfide, with the third os cuneiforme, and two last bones of the metatarsus. The superior ligaments are almost equally flat; the inferior unequally thick, and stronger than the superior. The exterior goes from the os cuboides to the tuberofity in the basis of the last bone of the metatarsus, and seems likewise to communicate with the third bone by fome ligamentous fibres.

The three offa cuneiformia are fixed to the os scaphoides and os cuboides by the ligaments above mentioned tioned. They are connected on the upper part by particular ligamentous planes, which go more or less transversely from one bone to another, being all joined to one common ligamentous plane, which covers these three bones, and also the os cuboides. On the lower part, they are joined by stronger and thicker ligaments. They are likewise connected by ligaments to the three first bones of the metatarsus.

The great os cuneiforme is joined on its upper, lower, and interior or convex fide, to the basis of the first metatarsal bone, by ligamentous fibres, which form almost a continued plane, the inferior part of which is strong and thick, and appears to be double. It is likewise tied to the inside of the basis of the second metatarsal bone by a particular ligament.

It has likewise on the outside of its inferior part three considerable ligaments more or less oblique: the first and shortest of which goes to the basis of the second metatarsal bone; the second, to that of the third; and the third, which is the longest, to that of the

fourth.

The bones of the metatarfus are connected together by their bases and heads. The ligaments that go between the bases, are superior and inferior. The superior are flat and small; the inferior strong and thick, and as it were multiplied by entering the interstices between the bases.

The ligaments which go between the heads have nearly the same general disposition. The inferior have this peculiar to them, that by filling the spaces between the heads they keep them at some distance from each other. The inferior portions of these ligaments are fixed in the angles at the lower parts of each head. They are also strengthned by their union and intertexture with the aponeurosis plantaris.

The first phalanges of the toes are tied to the heads of the metatarfal bones, by a fort of orbicular ligament, set round the edges of the cartilaginous portions of the

head,

Pa

fo

lig

th

la

m

10

fc

tl

16

p

head, and those of the bases of the phalanges. In the four toes next the great toe, the inferior part of these ligaments is very thick, and crusted over as it were with the cartilaginous substance fixed to the basis of the phalanges, and from thence continued over the head of the metatarfal bone next it. This substance grows hard with

age like a fefamoid bone.

Of these sesamoid bones the great toe has two belonging to the first phalanx, which are the largest, the foonest formed, and most considerable of all that go by that name; they are about one-third of an inch in length, and about half as broad as long. They are joined, by their anterior extremities, to the basis of this phalanx, close to each other; and lie in the two depressions on the lower side of the head of the first metatarfal bone.

The fecond and third phalanges of all the toes being articulated by ginglymi, have lateral ligaments which go between the fides of the bases to the fides of the heads. At the inferior edges of all these bases there is a cartilaginous matter joined to the ligaments, which hardens with age, in the fame manner as those of the first phalanges.

The capfular ligaments of all these articulations are disposed in the same manner as in the first bones of the

tarfus already described.

The periosteum of all these bones is of the same kind

with that of the bones of the leg.

The marrow is fuitable to their internal structure; that is, in moleculæ in the cavernous portions, and in masses in those which have large cavities. Thus the marrow of all the tarfal bones is dispersed in moleculæ, because their internal structure is spongy. In the metatarfal bones and first phalanges of the toes, it is difposed in the same manner as in the tibia and sibula; that is, it lies in moleculæ in the extremities, the itructure of which is cavernous; but in the middle portions of them, it lies in masses greater or less, according

cording to the fize of the cavities. In the other phalanges, which are entirely spongy, it is accordingly disposed in moleculæ.

The fynovial fubstances answer in number and figure to the depressions between the cartilaginous edges and

ligaments.

INDEX.

# INDEX.

A BDOMEN, muscles situated on the anterior part of it, 299. within its cavity, 311. Abductor indicis manus, 346. - indicis pedis, 370. longus pollicis manus, 344 minimidigiti manus, 346. medii digiti pedis, 371. minimi digiti pedis, 369. oculi, 275. pollicis manus, 345. pollicis pedis, 368. tertii digiti pedis, 371. Acetabulum of offa innominata, 178. Accelerator urinæ, 305. Adductor brevis femoris, 351. ad minimum digitum, 346. · indicis pedis, 370. - longus femoris, ib. - medii digiti pedis, 370. - magnus femoris, 351. - medii digiti pedis, 370. - metacarpi minimi digiti manus, 346. minimi digiti pedis, 371. oculi, 275. - pollicis manus, 345. - pollicis pedis, 368. - tertii digiti pedis, 370. Amygdalz, 292. Analysis, chemical, of bones, 28. - phenomenon folved by it, ib. Anconæus, 336. Angularis, 325. Ani fphincter, 306, 309. - levator, 307, 309. Anterior auris, 269. Anterior mallei, 271. Antithenar, 369. Antitragicus, 270. Anus, mufcles of the, 306. Aperiens palpebrarum rectus, 274. Apophyses, their different forts and uses, 36. Many so named are epiphyses, ib. Appendices. Vid. Epiphyfes. Arm-bone. Vid. Humeri os. Arthrodia, 48, 51. Articulations, 47. Symphysis, synar-throsis, ib. Diarthrosis, 48. Dispute concerning them, 49. Arytenoideus minor, 297. - major, 298. - obliquus, ib.

VOL. I.

Arytenoideus transversus, 298. Aryteno-epiglottideus, ib. Aftragalus, 245. Its articulation, 246. Of children, 247. Atlas, or the first vertebra of the neck, 153. Of infants, 154. Attollens aurem, 269. - ocuti, 275. - elevator oculi, ib, Aurem attollens, 269. Auris anterior, 270. - retrahentes, ib. - posterior, ib. - fuperior, ib. - tranfverfus, 271. - obliquus, ib. Axis, or third vertebra colli, 155. Azygos uvulz, 194. Basio-cerato-chondro glossus, 288. Biceps brachii, 335 - flexor cubiti, ib. - flexor cruris, 359. - internus, 335. Biventer maxillæ inferioris, 286. Bones, what, 20. Their plates, 21. Fibres, claviculi, ib. Cancelli, 12. Arteries, 23. Veins, 24. Nerves, 25. Circulation of their liquors, nourishment, increase, decrease, ib. Phenomena and diseases, 26, 31, 32, 43. Transverse and longitudinal canals, 26. Analysis, 28. Phenomenon, uses, ib. Periosteum internum, ib. Marrow, 30. Distinguished into broad and round, 33. Strength increased by being hollow, 34. Proceffes, 35. Cavities, 16. Epiphyfes, ib. Offification, 39. Articulations, 47. Brachialis, 336. - internus, 335. externus, 336. Breaft-bone. Bregma, 84. Broad bones ; their fructure, 33. Buccinator, 280. Bulbo-cavernofus, 305. Of children, Calcis os, 244, 247. 248. Cancelli of bones, 22. Corrugati, crihriformes, reticulares, ufcs, 23. Canini dentes, 137.

Caninus,

Caninus, 278.

Capitis obliquus fuperior, 328. obliquus inferior, ib. - par tertium, 325. - rectus, 319. Carpus confifts of eight bones, 216. Its figure, substance, articulation, motions, 221. Uses, 222. Of infants, ib. Cartilages, what; their plates, fibres, &c. 55. Oflification, how prevented, 56. Their uses, diseases, 57. Cavities of bones, their different kinds, 36. Uses, 37. Cephalo-pharyngeus, 296. Cerato-gloffus, 287. Cervical vertebræ, their distinguishing marks, 151. Atlas, 153. Denta-ta, 155. Axis, 158. Seventh, 159. Cervicalis descendens, 324 Cheek-bones, Vid. Malarum offa. Circumflexus, or tenfor palati, 291. Clavicles, 196. Internal end, 197. Body, ib. External end, medullary veffels, substance, articulation, 198. Of children, 199. Ules, ib. Claviculi of hones, perpendicular, oblique, headed, crooked, 21. Cleido-mastoideus, 285. Coccygeus, 310. Coccygis os, 168. Its first, second, third, and fourth bones, ib. Substance, motion, diseases, uses, 170. Collar-hone. Vid. Clavicle. Colli latiffimus, 285. - longus, 318. - tranfverfalis, 327. -- femispinalis, ib. - interspinal.s, 319. - intertranfverfales, 330. Complexus, 324 minor, 325. Compressor naris, 277. Conftrictor cunni, 308. oris, 282. - ishmi faucium, 293. --- pharyngis, 295, 296. Coraco-brachialis, 333. - hvoideus, 189. - radialis, 335. Coronal future, 67 Corrugator fupercilii, 268. Coffæ. Vid. Ribs. Cranium, its figure, 64. Surfaces, 65. Tables, diploe, 66. Confifts of fix proper, ib and two common bones, 67. Its futures, ib. Muscles of its teguments, 267. Cremafter, 304.

Crico-arytenoideus lateralis, 207. Crico-arytenoideus posticus, ib. Crico-thyroideus, 289. Crico-pharyngeus, 295. Crotaphyte muscle, 283. Cruralis, 359. Cubit, or fore-arm. Vid. Fore-arm. Cubitarius Riolani, 336. Cuboides os, 249. At the birth, 250, Cucularis, 320. Cuneiforme os, of the wrift, 217. Cuneiformia offa of the foot; externum, 250 medium, internum, ib. 251. At the birth, ib. Cutaneus, 284. D. Dartos, 304. Deltoides, 333. Dentata, or fecond vertebra of the neck, 155. Its diffinguishing marks, ib. Of infants, 156. Dentes. Vide Teeth. Deprimens, 275. Depreffor angeli oris, 279. - alæ nali, 279. - labii inferioris, ib. labii fuperioris alæque nafi, ib. - labii superioris proprius, ib. - labiorum communis, 280. oculi, 275. Diaphragma, 311. 312. Diarthrofis, 48. Its three species, enarthrosis, arthrodia, 48, 50, 51. ginglimus, 49, 51. Digastricus, 286. Digiti. Vide Fingers and Toes. Diploe of the skull, 66. Dorfal vertebræ, their distinguishing structure, 159. Wherein they ditfer from each other, 161. Dorfi interspinales, 330. - latiffimus, 321. fpinalis, 323. - longissimus, ib. - femispinalis, 326. Ear, muscles of the, 269. Elevator, 275, 325. - oculi, 269. - labii fuperioris, 279. - labii inferioris, 280. - labiorum communis, 278. Enarthrolis, 48, 50. Epicranius, 268. Epiphyses, 36. Several have apophy-fes, 37. Their uses, 38. Epistaphilinus, 294.

Erector clitoridis, 308.

- penis, 305,

Ethmoidal future, 97.

Eth-

Eth

1

1

Ex

Ethmoides os, 97. Its cribriform plate, 98. Nasal plate, ib. Cellulæ, osla spongiosa, 99. Connection, uses, 100. Morbid phenomena, 101. Of a child, ib. Extensor brevis digitorum pedis, 365. - carpi radialis brevior, 339. - carpi radialis longior, ib. - carpi ulnaris, ib. digitorum communis, 341. - longus digitorum pedis, 364. - magnus, 361. - minor pollicis manus, ib. - ossis metacarpi pollicis manus, 344. - pollicis primus, 345. pollicis fecundus, ib. - primi internodii pollicis manus, - proprius pollicis pedis, 362. - fecundi internodii pollicis manus, major pollicis manus, ib. - pollicis fecundus, ib. - tertii internodii, 346, 347. - tarfi furalis, 361. - tarsi minor, 362. Externus mallei, 271. Extremities, inferior, consist each of thigh, 229. leg, 234. foot, 244. Muscles of, 350. Front view of, Table I. p. 378. View of their muscles, 381. - superior, consist each of shoulder, 196. arm, 205. fore-arm, 209. hand, 215. Ufes, 228. Muscles of, 331. Front view of the bones, Table I. p. 577. View of their muscles, 380, 383. Eye-ball, muscles of, 274. Eye-lids, muscles of, 273. Face, 109. Composed of upper jaw, ib. 115. Lower jaw, 127. Fallopii capitis par tertium, 325. Fasciæ latæ, 356. Fauces, muscles about the, 293. Faucium isthmi constrictor, 293. Female skeleton, distinguishing marks of it, 257. Femoris os, 229. Its upper end, 230. Body, 231. Lower end, 232. infants, its articulation and motions, Fibula, its upper end, 237. Body, 238. Inferior end, 239. Its arti-culation, uses, 140. Of infants, 241. Fingers, their bones, phalanges, first, fecond, third, 227.

Flexor accessorius digitorum pedis, 366.

Flexor brevis digitorum pedis, 365. - brevis minimi digiti pedis, 369. brevis pollicis manus, 343. brevis pollicis pedis, 368. longus digitorum pedis, 366. - primi internodii, 344. tertii internodii, 343. fecundi internodii, 344. carpi radialis, 338. carpi ulnaris, ib. · longus pollicis manus, 343. longus pollicis pedis, 368. offis metacarpi pollicis, 344. parvus minimi digiti manus, 347. profundus perforans, 340. - fublimis perforatus, ib. Fontanelle, 84. Foot, composed of tarfus, 244. metatarfus, 252. toes, '254. Its motion, 246. Muscles on it, 362. Fore-arm, 209. Confifts of ulna, 210. Radius, 212. Muscles on it, 334, Frontal bone, 74. Its external furface, ib. Processes, 75. Cavities, ib. Foramina, ib. Internal furface, 77. Foramen, 78. Substance, sinuses, ib. Connection, 81. Ufes, ib. Of children, ib. Frontalis, 268. Gastrocnemius externus, 360. - internus, 361. Gemellus, 360. Gemini, 354. Genæ quadratus, 285. Generation, organs of; muscles about them, 304 Genio-hyo-gloffus, 287. Genio-hyoideus, ib. Ginglimus, its species, 49, 51. Gloffo-flaphilinus, 293. - pharyngeus, 296. Glottis, muscles fituated about it, 296. Gluteus maximus, 352. - medius, 353. minimus, ib. Gomphosis, 48. Gracilis, 356. Gracilo-internus, ib. H. Hand, 215. Confifts of carpus, 216. Metacarpus, 222. Fingers, 225. Its mufcles, 343. Harmonia, 48. Haunch-bones. Vide Ilium offa. Head confifts of cranium, 64. Face, 109. Its motions, 156, 157, 158. View of muscles on it and neck, 379. 382.

Heel-bone. Vide Calcis os. Helicis major, 270.

minor, ib. Hip-bone, Vide Os ischium.

Humeri os, 205. Its upper end, ib. Body, 206. Lower end, ib. Substance, articulation, 208. Motion, ib. Of children, 209. Muscles situated on, 334.

Hyoides os, 142. Its body, ib. Cornua, ib. Appendices, 143. Ligaments, substance, connection, uses, of children, 143, 144. Muscles about

it, 286, 288, 190. Hyo-gloffus, 287. Hyo-pharyngens, 296. Hyo-thyroideus, 289. Hypothenar minor, 346.

- Riolani, 347.

Jaw, lower. Vide Maxilla inferior upper. Vide Maxilla fuperior.

Iliacus internus, 314. Ilium offa, 172. Spine, dorfum, ib. External furface, 173. Interior furface, 174. Medullary vessels, sub-Stance, of a child, 174, 175.

Incisivus inferior, 280. - lateralis, 279. - medius, ib. Incifores dentes, 136. Indicator, 346. Inferior proftate, 306. Infraspinatus, 331.

- costales, 318. Innominata offa, 171. Composed of os ilium, 172. Ischium, 175. Pubis, 177. Their great foramen, 178. Acetabulum, ib. In infants, 172, 180. Connection, phenomena, 180.

Uses, 181. Intercostales externi, 317.

- interni, ib. Internus auris, 272. - malle, ib.

Interosseus auricularis, 348.

pedis externi, 370. - pedis interni, 371. Interspinales colli, 329.

- dorfi et lumborum, 330. Intertransversales colli, ib.

- dorfi, ib.

- lumborum, ib.

Ischium os, 175. Process, depressions, tuber, ib. Substance, 176. Of infants, 177.

Ischio-cavernosus, 305.

Ishmi faucium constrictor, 293. L.

Lambdoid suture, 67. Its additaments, ib.

Latissimus colli, 285.

dorfi, 321. Lattice-work of bones. Vide Cancelli. Laxator tympani, 271.

Leg, composed of tibia, 234. Fibula, 237. Rotula, 241. Its articulation

and motions, 242, 243, 244. Levator palati mollis, 292.

- parvus mollis, 306.

menti, 280.

- palpebræ fuperioris, 147.

oculi, 275.

anguli oris, 278.

- labii superioris alæque nafi, ib.

- labii inferioris, 280.

- palati, 291.

- ani, 307, 309.

Ligaments, 52. Their fibres, ib. Veffels, nerves, diseases, 53. Ufes, phenomena, 54. Enumeration of them, 405.

Longus colli, 318. Longiffimus dorsi, 323.

Lumbar vertebræ, their particula structure, 162. Differences, 163. their particular

Lumbaris internus, 314. externus, 313. Lumbricales, 341.

- pedis, 366. Lunare os, 217.

M.

Magnum os of the wrift, 219. Malarum offa, their processes, 114. Holes, substance, connection, ib. Of

infants, 115.

Marrow, what, 30. Its chemical analyfis, its arteries, veins, nerves, ib. Diseases, 32. Transverse and longitudinal canals for conveying it, 26, 31. Uses, 31. Phenomena and difeafes, 32.

Maxilla inferior, 127. The chin, fides, base, 128. Phenomena, angles, processes, 129. Foramina, substance, 129, 130. Articulation, 130. Motions, 131. Of infants, uses, 132.

— superior, consists of 13 bones,

109. Their connection, ib. 120. Maxillaria offa, their processes, 115. Cavities, 117. Foramina, 118. Si-nuses, 119. Morbid phenomena, ib. Substance, connection, 120. Ufes, 121. Of children, ib.

Marsupialis, 310. Mastoideus, 285. -lateralis, 325.

Maffeter, 283. Metacarpus of the hand, composed of four bones, 222. Substance at the birth,

birth, articulation, 223. Figure, uses, digiti indicis, ib. Medii, 224. Annularis, minimi, 225. Metacarpus muscle, 347. Metatarfus, 369. Metatarfus, composed of five bones, 252. Pollicis, &c. ib. 253, 254. Molares, dentes, 138. Mucilaginous glands, 58. Liquor fecreted by them, 59. Their cellular sub-stance, vessels, nerves, diseases, ib. Uses, 60. Phenomena, ib. Multifidus ípinæ, 316. Muscles, tables of them, 375-403. Musculus cutaneus, 284, 285. - fupercilii, 268. frontalis, ib. - Stapedii, 272. tubæ novus, 291. - patientiæ, 325. aponeurosis, 356. Mylo-pharyngeus, 296. - hyoideus, 286. Nasi offa, their sides, 171. Connection, uses; of an infant, 121. Naviculare os, of the tarfus, 248. Of children, 249. Neck. Vide Cervical vertebra. 0. Obliques inferier, 276. - auris, 271. major, 276, 328. minor, 318, 176. externus abdominis, 301. descendens, ib. internus abdominis, 302. ascendens, ib. - fuperior, 318, 176. descendens externus, 200. - descendens internus, 301. - capitis fuperior, 328. - capitis inferior, ib. Obturator internus, 310. Occipito frontalis, 267. Occipitalis, 268. Occipitis os, gr. Its external furface, 92. Processes, ib. Internal surface, 94. Holes, 95. Substance, 96. 94. Holes, 95. Substance, 96. Connection, ib. Ufes; in infants, Omo-hyoideus, 288. Orbicularis palpebrarum, 273. - oris, 281. Offification of bones, 39. Accounted for from particular disposition of veffels, and from preffure, 40. Phenomena, 39, 43. Its history, 45.

Offeagenea, its history, 45. Of what

ufc, 46.

Palati os, rar. Its square palate plate. ib. Pterygoid process, nasal lamella, 122. Orbitar process, 123. Substance, connection, uses, 123, 124. Of children, 124. Its difeafes, ib. Palato-falphingeus, 291. - Staphilinus, 294. - pharyngeus, 293-Palmaris longus, 337. - brevis, 338. cutaneus, ib. Parathenar major, 369. minor, ib. Parietal bones, their external furfage, 82. Internal furface, 83. Substante, ib. Connection, uses, of children, 84. Practical observations concerning bregma, ib. Patella. Vide Rotula. Pelvis, 171, 181. Confifts of os facrum, 164. Coccygis, 168. Offa innominata, 171. Its cartilages, ligaments, &c. 415. Perichondrium, its vessels, 55. Periosteum externum, 17. Its fibres, ib. Arteries, veins, nerves, 19. Ufes, ib. Difeafes, 20. internum, its structure, 28. Ufes, 29. Petro-falpingo-ftaphilinus, 292. Pectoralis, 315. - major, ib. minor, 316. Pectinalis, 365. Pectineus, 350. Perforatus, 340, 365. Perforans, 341. Peroneus maximus, 363. - posterior, ib. primus, ib. medius, 364. fecundus, ib. longus, 363. brevis. ib. Phalanges of the fingers, 227. toes, 254. Piliforme os of the wrift, 218. Plates of bones, their fibres, 21. Platisma myoides, 284, 285. Plantaris, 362, 361. Pollicis manus offa. Vide Thumb. pedis offa. Vide Toes. Poplitæus, 359-Pores, transverse and longitudinal of bones, 16. Their ufes, 27. Posterior auris, 270. - indicis, 347. annularis, 349. Posterior Posterior medii, ib.

Posticus, 363. Processes of bones. Vide Apophyses. Profundus, 341. Pronator radii teres, 342. - quadratus, ib. Prior indicis, 347. - annularis, 348. - medii, ib. Pfoas, 314. - parvus, 313. Pterygoideus major, 284. - minor, ib. - internus, 283. - externus, 284. Pterygo staphilinus externus vulgo, 292. Pubis os, 177. Substance; of children, 178. Pyramidalis, 279, 303. Pyriformis, 354. Quadratus, 280, 313. - genæ, 285. · lumborum, 313. femoris, 335. Radius, its upper end, 212. Body, 213. Lower end, ib. Of children, articulation, 214. Radialis internus, 338. - externus longior, 339. - externus primus, ib. - externus brevior, ib. Rectus abdominis, 302. - major, 328. - anterior longus, 319. - anterior brevis, ib. - internus, 357. Rectus, ib. - capitis internus major, 319. - capitis internus minor, ib. - capitis lateralis, ib. - capitis posticus major, 327. - capitis posticus minor, 328. Retrahentes auris, 270. Retractor anguli oris, 281. Rhomboidens, 322.

major, ib. - minor, ib. Ribs, 182. end, 183. Their middle posterior Anterior end, substance, cartilages, 184. Articulation and motion, 185. Differ from each other, 186. Diftinguished into true and false, 187. Distinguishing marks

of the sft, 2d, 3d, 4th, 5th, 6th,

7th, 11th, 12th, 187, 188, 189. 0f infants, 190. Motion, 194. Their cartilages, ligaments, &c. 413. Rinæus vel nasalis, 277. Rotula described, 241. Its substance, ib. Articulation of children, 242. Round bones, their structure, 33. Strength, how increased, 34. Sacro-lumbalis, 324.
Sacrum os, its body, 165. Processes oblique, transverse, ib. Spinal, 166.
Foramina, 167. Substance, articuculation, ib. Uses; of children, 169. Sagittal future, 68. Salpingo-staphilinus, 292. - staphilinus internus, ib. Salpingo-pharyngeus, 294. Sartorius, 356. Scalenus prior, 329. fecond, ib. - third, ib. - anticus, 328. - medius, 329. posticus, ib. Scaphoid bone of the wrift, 216. Scapula, 199. Its bafe, inferior costa, 200. Superior costa, dorsum, anterior surface, 201. Processes, spine, acromion, coracoid process, 202. Third process, medullary vessels, 203. Substance, articulation, ib. Uses; of children, 204. Schyndelesis, 48. - minus, ib. - major, ib. Semispinalis, 326, 327. - internus, ib. externus, ib. Semi-interosseus, 346. Semi-membranofus, 358. Seminervofus, ib. Semi orbicularis, 282. Semi tendinofus, 358. Serratus anticus, 316. - magnus, 316.
-- posticus inferior, 321. posticus superior, 323. Sesamoid bones, how formed, 255. Where to be found, ib. Share-bone. Vide Offa pubis. Shoulder, confifts of clavicle, 196, and scapula, 199. Shoulder-blade. Vide Scapula. Sinus frontale, 78. Sphenoidal, 101. Maxillary 119. Skeleton, natural, artificial, 62. Situation of the bones, 63. Divided in-

to head, ib. trunk, 144. extreme-

ties

Ta

fro

Skele

Skull

Solev

Sphe

fac

In

10

SI

of

Sph Sph

Sph

Sp

ties, fuperior, 196. inferior, 219. Table of its bones, 375, 390. Skeleton of a female, its differences from that of the male, 257. Skull. Vide Cranium.

Soleus, 361. Sphenoid bone, 101. Its external furfaces, processes, ib. Cavities, 103. Internal furface, ib. Holes proper, Internal Intrace, 10. Sinuses, 107. Substance, connection, ib. of children, 108.

Sphenoidal future, 71.

Sphene-falpingo-staphilinus, 291.

- staphilinus, 292. Sphineter ani, 306, 309.

- vaginæ, 308. - labiorum, 282.

- externus, 307.
- cutaneus, ib.

Spinalis, 327.

- dorfi, 323.

Spine, its shape, 144. Consists of true vertebræ, ib. and falfe, 164.

Spongiosa inferiora ossa, 124. Their fubitance, connection, uses; of children, 125.

— superiora offa, 99.
Squamous sutures, 69. How formed,

Staphilinus, 294.

externus, 291. Stapedius, 272.

Sterno-mastoideus, 285. - cleido-mastoideus, ib.

- hyoideus, 288.

- thyroideus, 189. - costalis, 318.

Sternum, 190. Its fubstance, 191. First bone, fecond bone, 191, 192. Xiphoid cartilage, 193. Diseases, ib. Connection, of children, uses, motion, 194, 195, 196. Its cartilages, ligaments, &c. 413.

Stylo-gloffus, 290.

- hyoideus, ib. - pharyngeus, 291.

Subclavius, 315. Subscapularis, 334.

Sublimis, 340, 365. Superior auris, 269.

Supinator longus, 342.

- radii longus, 341. - brevis, 342.

Supra castoles, 318. Supraspinatus, 331.

Surure, 47. how formed, ib.

of the cranium, 67. Coronal, lambdoid, ib. Sagittal, 68. Squamous, 69. Ethmoidal, fphenoidal, transverse, zygomatic, 71. Uses, 72. 73, 74.

Suture, of the face distinguished by names, 109. Uses, 101.

Symphysis, divided into sychondrosis. fyneurolis, fysfarcolis, 47.

Synarthrofis, divided into future, harmonia, gomphosis, schyndelesis, 47, 48.

Synchondrosis, 47. Syneurosis, ib.

Synovia of joints, what composed of, 58. Its ufes, difeafes, 59.

Syffarcofis, 47.

Tarfus confifts of feven bones, 244. Its

articulation, uses, 251, 252.
Teeth, 132. Their base, roots, periosteum, substance, 133, 134. Canals, 134. Vessels, ib. Formation, 135. Shedding, 136. Connection, ufes, ib.

Distinguished into incifores, Canini, molares, 137, 138. Pheno-

mena, 140.
Temporal hones, 85. Their external furface, processes, ib. Cavities, 87. Holes, ib. Internal furface, 89. Processes, cavities, holes, 89, 90. Substance, 90. Connection, ib. Uses

of infants, 91. Temporalis, 282.

Tenfor tympani, 271. - palati, 291.

- vaginæ femoris, 356.

Teres minor, 332. - major, ib.

Thenar, 368.

Thigh bone. Vide Femoris os.
Thorax, 182. Consists of vertebræ, 144. ribs, 182. Sternum, 190. Its motion, 194, 195. 196.

Thumb, consists of three bones; the first, its articulation, 225. At the time of birth, 226. Second bone, its articulation, ib. Third bone, ib.

Thyro-staphilinus, 294. - pharyngo-staphilinus, ib. - pharyngeus, 295.

Thyreo-arytænoideus, 297.

- epiglottideus, 298. Tibia, its upper end, cartilages, 234. Body, ib. Lower end, 236. Articulation; of children, 237.

Tibialis gracilis, 362. - anticus, ib.

- posticus, ib.

Toes, 254. Differ from the fingers; uses; of children, ib. 255.

Tranfverse suture, 71.

Trapezium \*

Transversus abdominis, 302. Trapezium os, 218. Trapezoides os, 219. Tragicus, 270. Transversus auris, 271. Transversalis, 302. Transversus perinei, 305, 306, 309. - alter, 306. Trapezius, 320. Trachelo-mastoideus, 325. Transversalis colli, 327. - pedis, 371. urethræ, 306. - lumborum, vulgo facer, 327. - dorfi, ib. --- colli, ib. Transverso-spinalis dorsi, 326. - lumborum, veterib. facer. ib. - spinalis colli pars interna, ib. - fpinalis colli pars externa, 327. Triangularis, 280, 318. Triceps minus, 351. - tertius, 352. - extensor cubiti, 336. - adductor femoris, 351. Triquetra offa, 68. Trochlearis, 276. Trochoides ginglimus, 51. Trunk of the skeleton, confists of the fpine, 144. pelvis, 171, 181. tho-rax, 182. Its cartilages and ligaments, 409. Turbinata offa. Vide Spongiofa. Vastus externus, 357. - internus, ib.

Vertebræ cervical. Vide Cervical.

Vertebræ dorfal. Vide Dorfal. false, composed of os facrum. 165. Coccygis, 168.

— lumbar. Vide Lumbar. es, 144. Their bodies, cartilages, 145. foramina, 166. Articulations, phenomena, 222. Of infants, 149. Dif. eases, 150. Divided into cervical, 151. dorsal, 159. lumbar, 162. Ufes, fecured from luxations, 164. Vessels of periosteum, 19. Of the bones, 23. Of the marrow, 31. Of ligaments, 53. Cartilages, 55. Mucilaginous glands, 59. Vomer, 125. Its figure, situation, ib. 126. Substance, connection, uses, 127. Ulna, its upper end, 210. Body, 248. Lower end, 211. Articulation, 212. Of children, 114. Ulnaris gracilis, 337. - internus, 339. externus, 340. Women, the differences of their bones from thefe of men, 257. Wormiana offa. Vide Triquetra. Wrist, Vide Carpus. Xiphoid cartilage, 193. Its discases, Zygoma,, 86, 114. Zygomatic suture, 71.

### END OF VOLUME FIRST.

Zygomaticus, 281.

- minor, ib.

## DIRECTIONS for placing the PLATES of Vol. I.

Table I. to fro	nt	•		P. 378
II.				381
III.				384
IV.				389
v.				390
VI.				392
VII.				396
VIII.				403

